



**Australian Government**

**Department of Agriculture,  
Fisheries and Forestry**

# **FINAL**

Review of policy: importation of grapevine  
(*Vitis* species) propagative material into  
Australia



April 2013

© Commonwealth of Australia

### **Ownership of intellectual property rights**

Unless otherwise noted, copyright (and any other intellectual property rights, if any) in this publication is owned by the Commonwealth of Australia (referred to as the Commonwealth).

### **Creative Commons licence**

All material in this publication is licensed under a Creative Commons Attribution 3.0 Australia Licence, save for content supplied by third parties, photographic images, logos, and the Commonwealth Coat of Arms.



Creative Commons Attribution 3.0 Australia Licence is a standard form licence agreement that allows you to copy, distribute, transmit and adapt this publication provided that you attribute the work. A summary of the licence terms is available from [creativecommons.org/licenses/by/3.0/au/deed.en](http://creativecommons.org/licenses/by/3.0/au/deed.en). The full licence terms are available from [creativecommons.org/licenses/by/3.0/au/legalcode](http://creativecommons.org/licenses/by/3.0/au/legalcode).

This publication (and any material sourced from it) should be attributed as:

Department of Agriculture, Fisheries and Forestry (2013) *Final review of policy: importation of grapevine (Vitis species) propagative material into Australia*. CC BY 3.0.

### **Cataloguing data**

Department of Agriculture, Fisheries and Forestry (2013) *Final review of policy: importation of grapevine (Vitis species) propagative material into Australia*. Department of Agriculture, Fisheries and Forestry, Canberra.

### **Internet**

*Final review of policy: importation of grapevine (Vitis species) propagative material into Australia* is available at [daff.gov.au](http://daff.gov.au).

Inquiries regarding the licence and any use of this document should be sent to: [copyright@daff.gov.au](mailto:copyright@daff.gov.au).

### **Disclaimer**

The Australian Government acting through the Department of Agriculture, Fisheries and Forestry has exercised due care and skill in the preparation and compilation of the information in this publication. Notwithstanding, the Department of Agriculture, Fisheries and Forestry, its employees and advisers disclaim all liability, including liability for negligence, for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying upon any of the information in this publication to the maximum extent permitted by law.

### **Biosecurity-Plant**

Department of Agriculture, Fisheries and Forestry  
GPO Box 858  
CANBERRA ACT 2601  
AUSTRALIA  
Telephone +61 2 6272 3933  
Facsimile +61 2 6272 3307  
Email [plant@daff.gov.au](mailto:plant@daff.gov.au)  
Website [daff.gov.au/ba](http://daff.gov.au/ba)

# Contents

---

<b>Tables and figures .....</b>	<b>4</b>
<b>Acronyms and abbreviations .....</b>	<b>5</b>
<b>Summary .....</b>	<b>6</b>
<b>1 Introduction .....</b>	<b>8</b>
1.1 AUSTRALIA'S BIOSECURITY POLICY FRAMEWORK.....	8
1.2 THIS REVIEW OF EXISTING POLICY .....	8
1.2.1 Background .....	9
1.2.2 Scope .....	9
1.2.3 Permitted species of grapevine.....	10
<b>2 Pest risk analysis.....</b>	<b>11</b>
2.1 STAGE 1: INITIATION .....	11
2.2 STAGE 2: PEST RISK ASSESSMENT.....	11
2.2.1 Pest categorisation .....	12
2.2.2 Assessment of the probability of entry, establishment and spread.....	15
2.2.3 Assessment of potential consequences.....	17
2.3 STAGE 3: PEST RISK MANAGEMENT.....	18
2.3.1 Identification and selection of appropriate risk management options .....	18
<b>3 Recommended risk management measures for grapevine propagative material .....</b>	<b>20</b>
3.1 PROPAGATIVE MATERIAL FROM ALL SOURCES (NON-APPROVED SOURCES) .....	20
3.1.1 Dormant cuttings .....	20
3.1.2 Tissue cultures (microplantlets) .....	26
3.1.3 Seed for sowing (non-approved sources).....	26
3.2 PROPAGATIVE MATERIAL FROM APPROVED SOURCES).....	27
3.2.1 Seed for sowing (approved sources) .....	27
<b>4 Framework for approval of high health sources and production requirements .....</b>	<b>29</b>
4.1 FRAMEWORK FOR APPROVAL OF HIGH HEALTH SOURCES .....	29
<b>5 Conclusion .....</b>	<b>30</b>
<b>Appendix A: Initiation and pest categorisation of pests associated with <i>Vitis</i> species worldwide .....</b>	<b>32</b>
<b>Appendix B: Additional quarantine pest data.....</b>	<b>226</b>
<b>Glossary .....</b>	<b>239</b>
<b>References .....</b>	<b>241</b>

## Tables and figures

---

Table 1.1 List of <i>Vitis</i> species permitted entry into Australia from all sources .....	10
Table 2.1 Quarantine pests for grapevine propagative material .....	13
Table 3.1 Recommended screening procedures for bacteria, fungi and phytoplasma.....	22
Table 3.2 Recommended grapevine virus indexing procedures.....	25

## Acronyms and abbreviations

---

Term or abbreviation	Definition
<b>ALOP</b>	Appropriate level of protection
<b>APPPC</b>	Asia and Pacific Plant Protection Commission
<b>APPD</b>	Australian Plant Pest Database
<b>CABI</b>	CAB International
<b>CMI</b>	Commonwealth Mycological Institute
<b>COSAVE</b>	Comité de Sanidad Vegetal del Cono Sur
<b>CPPC</b>	Caribbean Plant Protection Commission
<b>DAFF</b>	Australian Government Department of Agriculture, Fisheries and Forestry
<b>EPPO</b>	European and Mediterranean Plant Protection Organisation
<b>FAO</b>	Food and Agriculture Organization of the United Nations
<b>IAPSC</b>	Inter African Phytosanitary Council
<b>IMF</b>	Immunofluorescence
<b>IPC</b>	International Phytosanitary Certificate
<b>IPM</b>	Integrated Pest Management
<b>IPPC</b>	International Plant Protection Convention
<b>IRA</b>	Import Risk Analysis
<b>ISPM</b>	International Standard for Phytosanitary Measures
<b>JUNAC</b>	Comisión del Acuerdo de Cartagena
<b>NAPPO</b>	North American Plant Protection Organization
<b>NPPO</b>	National Plant Protection Organization
<b>OEPP</b>	Organisation Européenne et Méditerranéenne pour la Protection des Plantes
<b>PCR</b>	Polymerase chain reaction
<b>PEQ</b>	Post-entry quarantine
<b>PRA</b>	Pest risk analysis
<b>RT-PCR</b>	Reverse-transcription polymerase chain reaction
<b>SPS</b>	Sanitary and phytosanitary
<b>TEM</b>	Transmission electron microscopy
<b>WTO</b>	World Trade Organisation

## Summary

---

Australia initiated this review as new pathogens have been identified on grapevines (*Vitis* species) and several pathogens have extended their global range. Uncontrolled movement of infected propagative material has helped to spread these pathogens into new areas.

Additionally, the Grape and Wine Research and Development Corporation requested Plant Biosecurity to review and develop PEQ protocols for *Vitis* nursery stock that will minimise the time imported cultivars spend in quarantine. The review recommends several changes to the existing policy that will reduce the PEQ period for dormant cuttings and tissue cultures (microplantlets), while maintaining quarantine integrity.

### Recommended risk management measures

The recommended risk management measures for propagative material are detailed below.

#### All sources (unknown health status)

##### Dormant cuttings

- Mandatory on-arrival inspection; fumigation; hot water treatment; and surface sterilisation;
- Mandatory growth in a closed government PEQ facility for a minimum period of 16 months for pathogen screening (visual observation; culturing; and electron microscopy); and
- Active pathogen testing through herbaceous host indexing and molecular tests including, but not limited to, PCR or ELISA.

##### Tissue cultures (microplantlets)

- Mandatory on-arrival inspection;
- Mandatory growth in a closed government PEQ facility for a minimum period of 12 months for pathogen screening (visual observation; culturing; and electron microscopy); and
- Active pathogen testing through herbaceous host indexing and molecular tests including, but not limited to, PCR or ELISA.

##### Seed

- Mandatory on-arrival inspection, surface sterilisation, fungicidal treatment, and growth in a closed government PEQ facility for a minimum period of nine months for pathogen screening (visual observation and electron microscopy); and
- Active pathogen testing through herbaceous host indexing and molecular tests including, but not limited to, PCR.

#### Approved sources (High health sources)

Foundation Plant Services, California, USA is currently an approved source to supply pathogen tested grapevine propagative material to Australia. However, Plant Biosecurity will consider requests for approval of other overseas sources (e.g. institutions, NPPOs), based on the framework recommended in this review.

The recommended changes to import requirements for dormant cuttings and tissue cultures from non-approved sources will also apply to material from approved sources (e.g. the PEQ

period will be reduced to 16 months for dormant cuttings and 12 months for tissue cultures). Seed for sowing from approved sources is currently not subject to PEQ and this is recommended to continue.

Plant Biosecurity has made several changes following consideration of stakeholder comments on the *Draft review of policy – importation of grapevine (Vitis spp.) propagative material into Australia*. However, these changes have no impact on recommended risk management measures. The major changes made in the finalisation of this policy include:

- the addition of three viruses (*Grapevine Pinot gris virus*, *Grapevine red blotch-associated virus* and *Grapevine virus F*) as pathogens of quarantine concern; and
- the inclusion of *Daktulosphaira vitifoliae* and *Pseudococcus maritimus* as quarantine pests for grapevine propagative material.

# 1 Introduction

---

## 1.1 Australia's biosecurity policy framework

Australia's biosecurity policies aim to protect Australia against the risks that may arise from exotic pests<sup>1</sup> entering, establishing and spreading in Australia, thereby threatening Australia's unique flora and fauna, as well as those agricultural industries that are relatively free from serious pests.

The pest risk analysis (PRA) process is an important part of Australia's biosecurity policies. It enables the Australian Government to formally consider the risks that could be associated with proposals to import products into Australia. If the risks are found to exceed Australia's appropriate level of protection (ALOP), risk management measures are proposed to reduce the risk to an acceptable level. If it is not possible to reduce the risks to an acceptable level, then no trade will be allowed.

Successive Australian governments have maintained a conservative, but not a zero risk, approach to the management of biosecurity risks. This approach is expressed in terms of Australia's ALOP, which reflects community expectations through government policy and is currently described as providing a high level of protection aimed at reducing risk to a very low level, but not to zero.

Australia's PRAs are undertaken by Plant Biosecurity and Animal Biosecurity (formerly conjointly known as Biosecurity Australia), within the Department of Agriculture, Fisheries and Forestry (DAFF), using teams of technical and scientific experts in relevant fields. PRAs involve consultation with stakeholders at various stages during the process. Plant Biosecurity and Animal Biosecurity provide recommendations for animal and plant quarantine policy to Australia's Director of Animal and Plant Quarantine (the Secretary of the Australian Department of Agriculture, Fisheries and Forestry). The Director or delegate is responsible for determining whether or not an importation can be permitted under the *Quarantine Act 1908*, and if so, under what conditions. Plant Import Operations, within DAFF (formerly the Australian Quarantine and Inspection Service), is responsible for implementing appropriate risk management measures.

More information about Australia's biosecurity framework is provided in the *Import Risk Analysis Handbook 2007* (update 2009) located on the DAFF website [www.daff.gov.au/ba](http://www.daff.gov.au/ba).

## 1.2 This review of existing policy

Australia has an existing policy to import grapevine propagative material from all countries. However, this policy has not been reviewed for some time. Propagative material represents one of the highest plant quarantine risks, as it can harbour various forms of pathogens and arthropod pests. The introduction of plant pathogens, especially with latent infection, is of particular concern in propagative material. A range of exotic arthropod pests and pathogens can be introduced and established via propagative material when imported in a viable state for ongoing propagation.

---

<sup>1</sup> A pest is any species, strain or biotype of plant, animal, or pathogenic agent injurious to plants or plant products (FAO 2009).



### 1.2.1 Background

Many pathogens are associated with the production of grapevines worldwide. Like other vegetatively propagated crops, grapevines are infected by numerous pathogens, among which viroids, viruses and phytoplasmas play a major role, causing degenerative diseases, heavy losses and sometimes plant death. As grapevines are propagated mainly by vegetative means, there is a considerable risk of introducing and spreading these pathogens through international trade of grapevine propagative material. The introduction of economically important grapevine pests into Australia could result in substantial costs in eradication, containment or control. Pest establishment and spread could also lead to an increase in the use of chemical controls and could jeopardize export markets.

Australia's existing policy allows importation of grapevine propagative material (dormant cuttings, tissue culture and seed) from any source. The policy includes on-arrival inspection and mandatory treatment and growth in a government post-entry plant quarantine (PEQ) facility, with appropriate disease screening. Separate conditions also exist for dormant cuttings, tissue culture and seeds from approved sources.

Plant Biosecurity initiated this review as new pathogens have been identified on grapevines (*Vitis* species) and several pathogens have extended their global range. For instance, *Grapevine virus F* (Al-Rwahnih *et al.* 2012b), *Grapevine Pinot gris virus* (Giampetruzzi *et al.* 2012) and *Grapevine red blotch-associated virus* (Al-Rwahnih *et al.* 2012a) have recently been identified. *Grapevine leafroll-associated virus 5* (GLRaV-5) has been reported to have spread to vineyards in China, Chile, Portugal, Spain, Turkey; *Arabis mosaic virus* in Spain; GLRaV-4 in China; GSYV-1 in Washington state, Italy and France; GLRaV-2 and GVB in Croatia; *Citrus exocortis viroid* in China and *Grapevine yellow speckle viroid 1* and *Hop stunt viroid* in New Zealand (Martelli 2012). Uncontrolled movement of infected propagative material has helped to spread these pathogens into new areas. Additionally, the Grape and Wine Research and Development Corporation requested Plant Biosecurity to review and develop PEQ protocols for *Vitis* nursery stock that will minimise the time imported cultivars spend in quarantine, while maintaining quarantine integrity.

### 1.2.2 Scope

*Vitis* propagative material can currently be imported as dormant cuttings, tissue cultures (microplantlets) or seed. Whole plants (other than tissue cultures) of *Vitis* are not allowed entry into Australia, due to their significantly higher risk in comparison to other types of nursery stock commodities. Therefore, whole plants are not considered in this review. The scope of this review of existing policy is limited to:

- the identification of biosecurity risks associated with grapevine propagative material (dormant cuttings, tissue cultures and seed) from all sources;
- the evaluation of existing risk management measures for the identified risks; and
- the proposal of additional measures where appropriate.

This review does not consider existing phytosanitary measures during the pest risk assessment. Existing phytosanitary measures are only considered during the development of risk management measures, if they are required, following the pest risk analysis.

This policy review is limited to recommending appropriate phytosanitary measures to address the risk of introducing quarantine pests of grapevine propagative material into Australia. It is the importer's responsibility to ensure compliance with the requirements of all other regulatory and advisory bodies associated with importing commodities to Australia. Among others, these could include the Australian Customs Service, Department of Health and

Ageing, Therapeutic Goods Administration, Australian Pesticides and Veterinary Medicines Authority, Department of the Environment, Water, Heritage and the Arts and State Departments of Agriculture.

### 1.2.3 Permitted species of grapevine

There are a number of grapevine species (*Vitis* species) that are currently permitted entry into Australia as propagative material (dormant cuttings, tissue cultures and seed), subject to specific import conditions. These conditions are available on the Import CONditions database (ICON) at <http://www.aqis.gov.au/icon>. The list of *Vitis* species currently permitted entry into Australia (C 16904) from all sources is provided in Table 1.1. ‘Grapevine propagative material’ will hereafter refer to the dormant cuttings, tissue cultures and/or seed of these permitted species only.

**Table 1.1 List of *Vitis* species permitted entry into Australia from all sources**

Scientific names	Synonyms
<i>Vitis aestivalis</i> x ( <i>labrusca</i> x <i>vinifera</i> )	-
<i>Vitis aestivalis</i> x <i>Vitis vinifera</i>	-
<i>Vitis brevipedunculata</i> (Maxim.) Dippel	<i>Ampelopsis glandulosa</i> (Wall.) Momiy. var. <i>brevipedunculata</i> (Maxim.) Momiy, <i>Ampelopsis brevipedunculata</i> (Maxim.) Trautv
<i>Vitis glandulosa</i> Wall.	<i>Ampelopsis glandulosa</i> (Wall.) Momiy. var. <i>glandulosa</i>
<i>Vitis heterophylla</i> Thunb	<i>Ampelopsis glandulosa</i> (Wall.) Momiy. var. <i>heterophylla</i> (Thunb.) Momiy.
<i>Vitis himalayana</i> (Royle) Brandis	<i>Parthenocissus semicordata</i> (Wall.) Planch. var. <i>roylei</i> (King) Raizada & H. O. Saxena
<i>Vitis hypoglauca</i> (A. Gray) F. Mueller	<i>Cissus hypoglauca</i> A. Gray
<i>Vitis quadrangularis</i> (L.) Wall. ex Wight	<i>Cissus quadrangularis</i> L.
<i>Vitis rhombifolia</i> (Vahl) Baker	<i>Cissus alata</i> Jacq.; <i>Cissus rhombifolia</i> Vahl
<i>Vitis riparia</i> Michx.	-
<i>Vitis rupestris</i> Scheele	-
<i>Vitis sicyoides</i> (L.) Miq.	<i>Cissus verticillata</i> (L.) Nicolson & C. E. Jarvis subsp. <i>verti</i> ; <i>Cissus sicyoides</i> L.; <i>Viscum verticillatum</i> L.
<i>Vitis striata</i> (Ruiz & Pav.) Miq.	<i>Cissus striata</i> Ruiz & Pav. subsp. <i>striata</i>
<i>Vitis vinifera</i> L.	<i>Vitis vinifera</i> L. subsp. <i>vinifera</i> , <i>Vitis vinifera</i> L. subsp. <i>sativa</i> (DC.) Hegi; <i>Vitis vinifera</i> L. subsp. <i>sylvestris</i> (CC Gmel.) Hegi; <i>Vitis sylvestris</i> CC Gmel.

## 2 Pest risk analysis

---

Plant Biosecurity has conducted this pest risk analysis (PRA) in accordance with the International Standards for Phytosanitary Measures (ISPMs), including ISPM 2: *Framework for pest risk analysis* (FAO 2007) and ISPM 11: *Pest risk analysis for quarantine pests, including analysis of environmental risks and living modified organisms* (FAO 2004). The standards provide a broad rationale for the analysis of the scientific evidence to be taken into consideration when identifying and assessing the risk posed by quarantine pests.

Following ISPM 11, this pest risk analysis process comprises of three discrete stages:

- Stage 1: Initiation of the PRA
- Stage 2: Pest Risk Assessment
- Stage 3: Pest Risk Management

Phytosanitary terms used in this PRA are defined in ISPM 5 (FAO 2009).

### 2.1 Stage 1: Initiation

The *initiation* of a risk analysis involves identifying the reason for the PRA and the identification of the pest(s) and pathway(s) that should be considered for risk analysis in relation to the identified PRA area.

This commodity-based pest risk assessment was initiated by Plant Biosecurity as a basis for a review and possible revision of the existing phytosanitary regulations to import grapevine propagative material into Australia. Additionally, the Grape and Wine Research and Development Corporation requested Plant Biosecurity to review and develop PEQ protocols for *Vitis* nursery stock that will minimise the time imported cultivars spend in quarantine, while maintaining an appropriate level of protection from the threat of exotic pests and diseases. The review was also necessary as new pathogens have been identified on grapevine and several pathogens have extended their global range.

In the context of this PRA, grapevine propagative material (dormant cuttings, tissue culture and seed) is a potential import ‘pathway’ by which a pest can enter Australia.

A list of pests associated with grapevines worldwide was tabulated from published scientific literature, such as reference books, journals and database searches. This information is set out in Appendix A and forms the basis of the pest categorisation.

For this PRA, the ‘PRA area’ is defined as Australia for pests that are absent from Australia or of limited distribution and under official control in Australia.

### 2.2 Stage 2: Pest Risk Assessment

A pest risk assessment is the ‘evaluation of the probability of the introduction and spread of a pest and of the magnitude of the associated potential economic consequences’ (FAO 2009, p. 13). The pest risk assessment provides technical justification for identifying quarantine pests and for establishing phytosanitary import requirements.

This is a commodity-initiated pest risk analysis and risk is estimated through a standard set of factors that contribute to the introduction, establishment, spread or potential economic impact of pests. This pest risk assessment was conducted using three consecutive steps: pest categorisation; assessment of the probability of entry, establishment and spread; and assessment of potential consequences.

### 2.2.1 Pest categorisation

Pest categorisation is a process to examine, for each pest identified in Stage 1 (*Initiation of the PRA process*), whether the criteria for a quarantine pest is satisfied. In the context of propagative material, pest categorisation includes all the main elements of a full pest risk assessment. However, assessment of entry, establishment and spread is done in less detail for propagative material as pests are already with, or within, a suitable, living host that will be grown under favourable conditions to ensure the survival of the host plant. In addition, pests can spread from infected propagative material not only by natural dispersal, but also by domestic trade of infected nursery stock. The process of pest categorisation is summarised by ISPM 11 (FAO 2004) as a screening procedure based on the following criteria:

- identity of the pest;
- presence or absence in the endangered area;
- regulatory status;
- potential for establishment and spread in the PRA area; and
- potential for economic consequences in the PRA area.

Pests are categorised according to their association with the pathway; their presence or absence or regulatory status; their potential to establish or spread; and their potential for economic consequences. Pests associated with grapevines listed in Appendix A were used to develop a pathway-specific pest list for all pathways (dormant cuttings, tissue cultures and seed). This list identifies the pathway association of pests recorded on grapevines and their status in Australia; their potential to establish or spread; and their potential for economic consequences. Pests likely to be associated with grapevine propagative material, and absent or under official control in Australia, may be capable of establishment or spread within Australia if suitable ecological and climatic conditions exist.

The quarantine pests of grapevines from all sources identified in the pest categorisation are listed in Table 2.1. These pathogens fulfil the IPPC criteria for a quarantine pest, specifically:

- these pests are economically important (as they cause a variety of direct and indirect economic impacts, such as reduced yield, reduced commodity value and/or loss of foreign or domestic markets); and
- these pests are not present in Australia or have a limited distribution and are under official control.

Pests under official control in Australia have been taken into account in this review. If regional pests<sup>2</sup> are identified on plants during PEQ, DAFF will notify relevant State Departments of Agriculture.

---

<sup>2</sup> Regional pests are quarantine pests for specific Australian states and territories, but may be present in other Australian states.

**Table 2.1 Quarantine pests for grapevine propagative material**

Pest type	Pathway association <sup>3</sup>		
	Dormant cuttings	Tissue cultures	Seed
<b>ARTHROPODS</b>			
<b>ACARI (mites)</b>			
<i>Brevipalpus chilensis</i> Baker [Acari: Tenuipalpidae]	✓		
<i>Colomerus vitis</i> Pagenstecher strain c [Acari: Eriophyidae]	✓		
<b>COLEOPTERA (beetles, weevils)</b>			
<i>Sinoxylon perforans</i> Schrank [Coleoptera: Bostrichidae]	✓		
<i>Sinoxylon sexdentatum</i> Olivier [Coleoptera: Bostrichidae]	✓		
<b>HEMIPTERA (aphids, leafhoppers, mealybugs, psyllids, scales, true bugs, whiteflies)</b>			
<i>Daktulosphaira vitifoliae</i> Fitch [Hemiptera: Phylloxeridae]	✓		
<i>Planococcus ficus</i> Signoret [Hemiptera: Pseudococcidae]	✓		
<i>Planococcus lilacinus</i> Cockerell [Hemiptera: Pseudococcidae]	✓		
<i>Planococcus kraunhiae</i> Kuwana [Hemiptera: Pseudococcidae]	✓		
<i>Pseudococcus maritimus</i> Ehrhorn [Hemiptera: Pseudococcidae]	✓		
<i>Targionia vitis</i> Signoret [Hemiptera: Diaspididae]	✓		
<b>LEPIDOPTERA (moths, butterflies)</b>			
<i>Paranthrene regalis</i> Butler [Lepidoptera: Sesiidae]	✓		
<i>Zeuzera coffeae</i> Nietner [Lepidoptera: Cossidae]	✓		
<b>PATHOGENS</b>			
<b>BACTERIA</b>			
<i>Xanthomonas campestris</i> pv. <i>viticola</i> (Nayudu) Dye	✓	✓	
<i>Xylella fastidiosa</i> (Wells et al.) – grapevine strain	✓	✓	
<i>Xylophilus ampelinus</i> (Panagopoulos) Willems et al.	✓	✓	
<b>FUNGI</b>			
<i>Alternaria viticola</i> Brunaud	✓		
<i>Cadophora luteo-olivacea</i> (J.F.H Beyma) T.C. Harr. & McNew	✓		
<i>Cadophora melinii</i> Nannf.	✓		
<i>Eutypella leprosa</i> (Pers.) Berl.	✓		
<i>Eutypella vitis</i> (Schwein.:Fr.) Ellis & Everhart	✓		
<i>Fomitiporia mediterranea</i> M. Fischer	✓		
<i>Fomitiporia polymorpha</i> M. Fischer	✓		
<i>Guignardia</i> species ( <i>Guignardia bidwellii</i> , <i>Guignardia bidwellii</i> f. <i>euvitis</i> , <i>Guignardia bidwellii</i> f. <i>muscadinii</i> )	✓		
<i>Inocutis jamaicensis</i> (Murrill) Gottlieb et al.	✓		
<i>Monilinia fructigena</i> Honey	✓		
<i>Phaeoacremonium</i> species ( <i>P. alvesii</i> , <i>P. angustius</i> , <i>P. argentinense</i> , <i>P. armeniacum</i> , <i>P. austroafricanum</i> , <i>P. cinereum</i> , <i>P. croatiense</i> , <i>P. globosum</i> , <i>P. griseorubrum</i> , <i>P. hispanicum</i> , <i>P. hungaricum</i> , <i>P. inflatipes</i> , <i>P. iraniamum</i> , <i>P. krajdinii</i> , <i>P. mortoniae</i> , <i>P. occidentale</i> , <i>P. rubrigenum</i> , <i>P. scoltyi</i> , <i>P. sicilianum</i> , <i>P. subulatum</i> , <i>P. tuscanum</i> , <i>P. venezuelense</i> , <i>P. viticola</i> )	✓		
<i>Phakopsora</i> species ( <i>Phakopsora euvitis</i> , <i>Phakopsora muscadinae</i> , <i>Phakopsora uva</i> )	✓		

<sup>3</sup> This review considers that certain pathogens (bacteria, phytoplasma, viroids and viruses) may not be excluded from the pathway and remains associated with micropropagated plantlets (tissue culture). In contrast, it considers that fungal or fungal-like pathogens are not on the pathway of micropropagated plantlets.

Pest type	Pathway association <sup>3</sup>		
	Dormant cuttings	Tissue cultures	Seed
<b>PHYTOPLASMA</b>			
<i>Candidatus</i> Phytoplasma asteris [16SrI – Aster yellows group]	✓	✓	
<i>Candidatus</i> Phytoplasma fraxini [16SrVII-A] (Ash yellows group)	✓	✓	
<i>Candidatus</i> Phytoplasma phoenicium [16SrIX]	✓	✓	
<i>Candidatus</i> Phytoplasma pruni [16SrIII – peach X-disease phytoplasmas group]	✓	✓	
<i>Candidatus</i> Phytoplasma solani [16 SrXII-A] (Stolbur group)	✓	✓	
<i>Candidatus</i> Phytoplasma ulmi [16SrV–A] (Elm yellows group EY group)	✓	✓	
<i>Candidatus</i> Phytoplasma vitis [16SrV] (Elm yellows group)	✓	✓	
European stone fruit yellows Phytoplasma 16SrX-B (Apple proliferation group)	✓	✓	
<b>VIRUSES</b>			
<i>Arabis mosaic virus</i> (ArMV) – grape strain	✓	✓	✓
<i>Artichoke Italian latent virus</i> (AILV)	✓	✓	
<i>Blueberry leaf mottle virus</i> (BLMoV) New York (NY) strain	✓	✓	✓
<i>Cherry leafroll virus</i> (CLRV) – grape isolate	✓	✓	
<i>Grapevine ajinashika virus</i> (GAgV)	✓	✓	
<i>Grapevine Anatolian ringspot virus</i> (GARSV)	✓	✓	
<i>Grapevine angular mosaic-associated virus</i> (GAMaV)	✓	✓	✓
<i>Grapevine asteroid mosaic associated virus</i> (GAMV)	✓	✓	
<i>Grapevine berry inner necrosis virus</i> (GINV)	✓	✓	
<i>Grapevine Bulgarian latent virus</i> (GBLV)	✓	✓	✓
<i>Grapevine chrome mosaic virus</i> (GCMV)	✓	✓	✓
<i>Grapevine deformation virus</i> (GDefV)	✓	✓	
<i>Grapevine fanleaf virus</i> (GFLV)	✓	✓	✓
<i>Grapevine leafroll associated virus</i> (GLRaV – 6,7,10, 11)	✓	✓	
<i>Grapevine line pattern virus</i> (GLPV)	✓	✓	✓
<i>Grapevine Pinot gris virus</i> (GPGV)	✓	✓	
<i>Grapevine red blotch-associated virus</i> (GRBaV)	✓	✓	
<i>Grapevine red globe virus</i> (GRGV)	✓	✓	
<i>Grapevine rupestris vein feathering virus</i> (GRVFV)	✓	✓	
<i>Grapevine syrah virus-I</i> (GSyV-I)	✓	✓	
<i>Grapevine Tunisian ringspot virus</i> (GTRSV)	✓	✓	
<i>Grapevine virus B</i> (strains associated with corky bark) (GVB)	✓	✓	
<i>Grapevine virus E</i> (GVE)	✓	✓	
<i>Grapevine virus F</i> (GVF)	✓	✓	
<i>Peach rosette mosaic virus</i> (PRMV)	✓	✓	✓
<i>Petunia asteroid mosaic virus</i> (PeAMV)	✓	✓	
<i>Raspberry ringspot virus</i> (RpRSV) – grapevine strain	✓	✓	
<i>Sowbane mosaic virus</i> (SoMV) – grape infecting strain	✓	✓	
<i>Strawberry latent ringspot virus</i> (SLRSV)	✓	✓	
<i>Tobacco necrosis virus</i> (TNV) – grape strain	✓	✓	
<i>Tomato black ring virus</i> (TBRV)	✓	✓	✓
<i>Tomato ringspot virus</i> (ToRSV)	✓	✓	✓

## **2.2.2 Assessment of the probability of entry, establishment and spread**

Details of how to assess the ‘probability of entry’, ‘probability of establishment’ and ‘probability of spread’ of a pest are given in ISPM 11 (FAO 2004).

In the case of propagative material imports, the concepts of entry, establishment and spread have to be considered differently. Propagative material intended for ongoing propagation purposes is deliberately introduced, distributed and aided to establish and spread. This material will enter and then be maintained in a suitable habitat, potentially in substantial numbers and for an indeterminate period. Significant resources are utilised to ensure the continued welfare of imported propagative material. Therefore, the introduction and establishment of plants from imported propagative material in essence establishes the pests and pathogens associated with the propagative material. Pathogens, in particular, may not need to leave the host to complete their life cycles, further enabling them to establish in the PRA area. Furthermore, propagative material is expected to be shipped at moderate temperatures and humidity, which is unlikely to adversely affect any pest that is present during shipment.

Several key factors contribute to the increased ability of pests and pathogens associated with propagative material to enter, establish and spread in Australia.

### **Probability of entry**

- Association with host commodities provides the opportunity for the pest to enter Australia. Their ability to survive on, or in, propagative material acts to ensure their viability on route to, and during distribution across, Australia.
- Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected. Therefore, propagative material provides a pathway for viruses.
- Propagative material is assumed to come from areas where these pests specifically occur and no phytosanitary measures have been applied. The primary conditions for survival of pests are fulfilled by the presence of the live propagative material and the associated environmental conditions. Therefore, association with propagative material can provide long term survival for the pests.
- Infected propagative material is the main pathway for the introduction of the pests into new areas. This mode of introduction is greatly enhanced because of latency periods before conspicuous symptoms develop. Long latency periods can lead to the propagation and distribution of infected propagative material and can therefore assist in the introduction of these pests into Australia.
- The pests associated with propagative material may be systemic or are associated with the vascular system (or occur internally in the nursery stock) and they are unlikely to be dislodged during standard harvesting, handling and shipping operations. Therefore, pests associated with propagative material are likely to survive during transport.
- Seeds will be maintained at a suitable temperature and humidity to maintain seed viability. Seed-borne and seed-transmissible pathogens will therefore be maintained within the seed for subsequent propagation.

### **Probability of establishment**

- Association with the host will facilitate the establishment of pests of propagative material, as they are already established with, or within, a suitable host. As host plant material is likely to be maintained in places with similar climates to the area of production, climatic

conditions are expected to favour the pest's establishment.

- Some pest specific factors are likely to impact upon a pest's ability to establish in Australia. For example, the likelihood of establishment will vary if an alternative host is required for the pest to complete its life cycle or if multiple individuals are required to form a founder population. Where appropriate, these considerations are addressed in the potential for establishment and spread field of the pest categorisation.
- Propagative material, including grapevine cuttings, tissue culture and seed, is intended for ongoing propagation or horticultural purposes and therefore is deliberately introduced, distributed and aided to establish. This material will enter and then be maintained in a suitable habitat, potentially in substantial numbers and for an indeterminate period. Therefore, the introduction and establishment of plants from imported propagative material in essence establishes the pests and pathogens associated with the propagative material.
- The latent period of infection before visible symptoms appear may result in non-detection of these pathogens; therefore, the pathogens will have ample time to establish into new areas.

### **Probability of spread**

- The ability of the pest to be introduced and distributed throughout Australia on propagative material through human mediated spread is a high risk for continued spread post-border in Australia. Pest related factors that aid the spread of the pest once it has established in Australia (such as wind, water or mechanical transmission) will increase the pest's ability to spread from an already high baseline.
- In the absence of statutory control, there is a high probability that the pests will spread quickly in Australia by trade of propagative material. Planting of infected propagative material will bring the pests into the environment. Climatic conditions, such as those found in propagation houses, may be sufficient for pest survival and spread.
- The systemic nature of some of the pests associated with propagative material is a major pathway for dispersal. Accordingly, local and long-distance spread of these pathogens has been associated with the movement of infected propagative material.
- The symptomless nature of several pathogens may contribute to the inadvertent propagation and distribution of infected material that will help spread these pathogens within Australia. Additionally, insect vectors present in Australia will help spread viruses from infected plants to healthy plants.
  - Viruses may differ in particle morphologies, disease symptoms induced and means of natural spread by insect or nematode vectors. However, each virus is readily carried and dispersed in nursery stock.
  - In some instances, pathogens may be introduced via infected plants into a viticulture region where native vector species reside resulting in secondary spread to neighbouring grapevines or to surrounding vineyards.

As a result of these pathway specific factors, it would be inappropriate to assess the probability of entry, establishment and spread using the processes described in ISPM 11 (FAO 2004). For the purposes of this PRA, the overall likelihood for the probability of entry, establishment and spread is considered to be high for pests entering Australia on grapevine propagative material.



### 2.2.3 Assessment of potential consequences

The purpose of assessment of potential consequences in the pest risk assessment process is to identify and quantify, as much as possible, the potential impacts that could be expected to result from a pest's introduction and spread.

The basic requirements for the assessment of consequences are described in the SPS Agreement, in particular Article 5.3 and Annex A. Further detail on assessing consequences is given in the “potential economic consequences” section of ISPM 11. This ISPM separates the consequences into “direct” and “indirect” and provides examples of factors to consider within each.

The introduction of pests which meet the criteria of a quarantine pest will have unacceptable economic consequences in Australia as these pests will cause a variety of direct and indirect economic impacts. The identified pests are of economic concern and do not occur in Australia or are under official control. A summary and justification is provided below:

- Direct impacts of the introduction and spread of multi-host pests in Australia will not only affect the imported host but also other hosts.
- Introduction and establishment of quarantine pests in Australia would not only result in phytosanitary regulations imposed by foreign or domestic trading partners, but also in increased costs of production, including pathogen control costs.
- Quarantine pest introduction and establishment would also be likely to result in industry adjustment. The potential economic impact for the nursery trade is high. Without controls, these pests have the potential to spread further in the trade network and could potentially expand their host range.
- Grapevines that are vegetatively propagated may be exposed to attack by a variety of pests and pathogens. Of these pests, infectious intracellular agents (viruses, viroids, bacteria and phytoplasmas) play a major role, causing heavy yield loss, shortening the productive life of vineyards and endangering the survival of affected vines (Martelli and Boudon-Padieu 2006).
- Both phytoplasmas and viruses are able to affect fruit development and ripening, possibly as a result of phloem disruption. This blockage can hinder berry sugar accumulation and delay ripening.
- Grapevine viruses cause yield loss, reduced fruit quality, reduced vine growth, vine decline and vine death. For example, leafroll viruses and rugose wood viruses are associated with yield losses (Guidoni *et al.* 2000; Mannini and Credi 2000; Kovacs *et al.* 2000, 2001; Tomazic *et al.* 2000, 2005; Komar *et al.* 2007). Leafroll viruses also cause poor fruit quality (Woodham *et al.* 1983; Komar *et al.* 2007). Grapevine fanleaf virus and Arabis mosaic virus are associated with significant yield loss, reduced fruit quality, reduced vine vigour, vine decline and vine death (Auger *et al.* 1992; Martelli 1993; Walter and Martelli 1998; Golino *et al.* 2003; Legorburu *et al.* 2009; Santini *et al.* 2009). Rugose wood complex viruses are associated with vine death (Tomazic *et al.* 2005).
- The identified pests are considered important as they cause a variety of direct and indirect economic impacts, such as reduced yield, reduced commodity value and loss of foreign or domestic markets. Therefore, these pests have a potential for economic consequences in Australia. For example, some of these pathogens are identified by COSAVE, EPPO, NAPPO and other countries as pests of quarantine concern. The presence of these pests and pathogens in Australia would impact upon Australia's ability to access overseas markets.

Pests and pathogens listed in Table 2.1 are of economic significance and are either absent from Australia, or if present, are under official control. Therefore, they meet the IPPC criteria for a quarantine pest and phytosanitary measures are justified to manage these pests and pathogens.

## 2.3 Stage 3: Pest Risk Management

ISPM 11 (FAO 2004) provides details on the identification and selection of appropriate risk management options. Pest risk management describes the process of identifying and implementing phytosanitary measures to manage risks posed by identified quarantine pests, while ensuring that any negative effects on trade are minimised.

Pest risk management evaluates and selects risk management options to reduce the risk of entry, establishment or spread of identified pests for the identified import pathways. To effectively prevent the introduction of pests associated with an identified pathway, a series of important safeguards, conditions or phytosanitary measures must be in place. Propagative material represents a direct pathway for pests identified by the pest categorisation. This pathway is direct since the end-use is the planting of a known host plant.

### 2.3.1 Identification and selection of appropriate risk management options

Phytosanitary measures to prevent the establishment and spread of quarantine pests may include any combination of measures, including pre- or post-harvest treatments, inspection at various points between production and final distribution, surveillance, official control, documentation, or certification. A measure or combination of measures may be applied at any one or more points along the continuum between the point of origin and the final destination. Pest risk management explores options that can be implemented (i) in the exporting country, (ii) at the point of entry or (iii) within the importing country. The ultimate goal is to protect plants and prevent the introduction of identified quarantine pests.

Examples of phytosanitary measures which may be applied to propagative material consignments include:

- **Import from pest free areas only (ISPM 4, 10)**—the establishment and use of a pest free area by an NPPO provides for the export of plants from the exporting country to the importing country without the need for application of additional phytosanitary measures when certain requirements are met.
- **Inspections or testing for freedom from regulated pests**—this is a practical measure for visible pests or for pests which produce visible symptoms on plants.
- **Inspection and certification (ISPM 7, 12, 23)**—the exporting country may be asked to inspect the shipment and certify that the shipment is free from regulated pests before export.
- **Specified conditions for preparation of the consignment**—the importing country may specify steps that must be followed in order to prepare the consignment for shipment. These conditions can include the requirement for plants to be produced from appropriately tested parent material.
- **Pre-entry or post-entry quarantine**—the importing country may define certain control conditions, inspection and possible treatment of shipments upon their entry into the country. Post-entry quarantine (PEQ) of dormant cuttings, seed and even *in vitro* plantlets

can help avoid introduction of new viruses or allied pathogens into the importing countries.

- **Removal of the pest from the consignment by treatment or other methods**—the importing country may specify chemical or physical treatments that must be applied to the consignment before it may be imported.

Measures can range from total prohibition to permitting import subject to visual inspection. In some cases more than one phytosanitary measure may be required in order to reduce the pest risk to an acceptable level.

### 3 Recommended risk management measures for grapevine propagative material

---

The ultimate goal of phytosanitary measures is to protect plant health and prevent the introduction of identified quarantine pests associated with grapevine propagative material. Plant Biosecurity considers that the risk management measures recommended in this final review of policy will be adequate to mitigate the risks posed by the identified quarantine pests and pathogens.

#### 3.1 Propagative material from all sources (non-approved sources)

The review recommends pro-active testing and a reduction in the growth period in PEQ for dormant cuttings and tissue cultures from all sources. Recommended testing procedures are based on active testing for quarantine pathogens, using traditional and modern techniques. This approach allows dormant cutting imports to be screened for a minimum period of 16 months in PEQ instead of the current 24 months and tissue cultures to be screened for a minimum period of 12 months in PEQ instead of the current 24 months.

##### 3.1.1 Dormant cuttings

The restriction of grapevine to one year old dormant cuttings with 2–3 internodes from all sources (approved or non-approved sources) is recommended to continue. Fully dormant canes should be imported during January to February from the Northern Hemisphere and July to September from the Southern Hemisphere. If this does not occur, there may be delays in the release of planting material because the growth period may be too short to obtain sufficient material to conduct required testing.

##### Mandatory on-arrival inspection

Imported dormant cuttings must be subject to mandatory on-arrival inspection to verify freedom from disease symptoms, live insects, soil and other extraneous contaminants of quarantine concern.

##### Mandatory on-arrival fumigation

It is recommended that imported dormant cuttings be subject to mandatory on-arrival methyl bromide fumigation (T9060) to manage the risk posed by arthropod pests from all sources.

Alternative treatments to methyl-bromide fumigation for grapevine dormant cuttings, if requested by an exporting country, will be considered by Plant Biosecurity on a case by case basis. Prior to the acceptance of an alternative treatment for grapevine dormant cuttings, Plant Biosecurity would need to assess the efficacy of that fumigant to ensure it gives an equal level of protection to methyl-bromide for all pests likely to be associated with the commodity.

##### Mandatory hot water treatment

It is recommended that dormant cuttings be subjected to hot water treatment at 50 °C for 30 minutes to minimise the risk of phytoplasmas.

- Hot water treatment at 50 °C for 30 minutes is effective against some phytoplasmas (Caudwell *et al.* 1997) and in eliminating most known fungal pathogens and endophytes from grapevine cuttings, including pathogens associated with young grapevine decline (Crous *et al.* 2001).
- After hot water treatment, dormant cuttings must be plunged into cold water to quickly

lower the temperature and minimise heat damage to the tissue (Waite *et al.* 2005).

### **Mandatory sodium hypochlorite treatment**

It is recommended that dormant cuttings be subjected to sodium hypochlorite treatment (1% NaOCl for 5 minutes) for surface sterilisation. Sodium hypochlorite treatment of dormant grapevine cuttings has been recommended to facilitate the safe introduction of grapevine propagative material (Frison and Ikin 1991). Treatment with sodium hypochlorite should be undertaken after the hot water treatment outlined above; this should allow some residual effect and increase the efficacy of the sodium hypochlorite treatment.

### **Mandatory culturing**

It is recommended that following hot water and sodium hypochlorite treatments, macerated buds from dormant cuttings be cultured to detect bacterial and fungal pathogens. This broad spectrum culturing test is useful to screen imported dormant cuttings for fungal and bacterial pathogens.

### **Mandatory growth in PEQ facilities**

It is recommended that imported grapevine cuttings be grown in a closed government PEQ facility for a minimum period of 16 months. The purpose of growth in PEQ facilities is to screen imported grapevine propagative material for pathogens in order to prevent the introduction of quarantine pests into Australia. It is recommended that newly established plants are maintained at 20–25 °C for 12 months in closed quarantine followed by four months growth in screen houses. During growth in PEQ, plants must be subject to pathogen screening, visual inspection and pathogen testing, as outlined below.

#### **Pathogen screening**

It is recommended that during PEQ growth period, plants and plantlets are subjected to visual inspection, electron microscopy and active testing, including biological indexing and molecular testing.

#### **Visual inspection**

Pathogen screening (visual screening) during growth in PEQ is recommended to continue for the detection of symptomatic pathogens. Fungal and bacterial pathogens associated with grapevines may produce distinct symptoms that make them easy to identify by visual inspection during growth period in PEQ.

#### **Pathogen testing**

The recommended pathogen testing during growth in PEQ will include active testing for quarantine pathogens, using traditional and modern techniques. Laboratory methods; including culturing, biological indicators, electron microscopy and molecular tests (PCR); may be used to detect grapevine pathogens.

#### **Bacterial pathogens**

- Active pathogen testing including molecular tests for *Xylella fastidiosa*, in addition to hot water treatment and visual inspection is recommended.
- Diagnostic tests, including culturing and microscopy, are recommended for *Xanthomonas campestris* pv. *viticola* and *Xylophilus ampelinus*. However, if symptoms develop during growth in PEQ, molecular testing (including PCR) for *Xanthomonas campestris* pv. *viticola* (Trindade *et al.* 2005) and *Xylophilus ampelinus* (Botha *et al.* 2001) is recommended.

**Fungal pathogens**

- Newly established plants (from imported propagative material) will be subject to growing season inspection and if symptoms develop during the PEQ period, further diagnostic testing; including culturing, microscopy and molecular tests; is recommended.

**Phytoplasmas**

- Newly established plants (from imported propagative material) will be subject to growing season inspection and active pathogen testing, including a generic PCR.

Recommended pathogen testing procedures are summarised in Table 3.1.

**Table 3.1 Recommended screening procedures for bacteria, fungi and phytoplasma**

Pathogen type	Mandatory screening			Additional tests <sup>4</sup>	Reference(s)
	Growing season inspection	Culture & microscopy	PCR		
BACTERIA					
<i>Xanthomonas campestris</i> pv. <i>viticola</i>	✓	✓		PCR	Trindade <i>et al.</i> 2005
<i>Xylella fastidiosa</i>	✓		✓		Luck <i>et al.</i> 2012
<i>Xylophilus ampelinus</i>	✓	✓		PCR	Botha <i>et al.</i> 2001
Fungi					
<i>Alternaria viticola</i>	✓	✓			
<i>Cadophora luteo-olivacea</i>	✓	✓			
<i>Cadophora melinii</i>	✓	✓			
<i>Eutypella leprosa</i>	✓	✓			
<i>Eutypella vitis</i>					
<i>Fomitiporia mediterranea</i>	✓	✓		PCR	Pilotti <i>et al.</i> 2010
<i>Fomitiporia polymorpha</i>					
<i>Guignardia</i> species	✓	✓			
<i>Inocutis jamaicensis</i>	✓	✓			
<i>Monilinia fructigena</i>	✓	✓			
<i>Phaeoacremonium</i> species	✓	✓	PCR	Aroca and Raposo 2007	
<i>Phakopsora</i> species	✓	✓			
Phytoplasma					
<i>Candidatus</i> Phytoplasma <i>asteris</i>	✓		✓		Deng and Hiruki 1991; Lee <i>et al.</i> 1995; Schneider <i>et al.</i> 1995
<i>Candidatus</i> Phytoplasma <i>fraxini</i>					
<i>Candidatus</i> Phytoplasma <i>phoenicium</i>					
<i>Candidatus</i> Phytoplasma <i>pruni</i>					
<i>Candidatus</i> Phytoplasma <i>solani</i>					
<i>Candidatus</i> Phytoplasma <i>ulmi</i>					
<i>Candidatus</i> Phytoplasma <i>vitis</i>					
European stone fruit yellows Phyttoplasma					

<sup>4</sup> If disease symptoms develop

## Viruses

Grapevine viruses are transmissible entities; they can be detected and identified on herbaceous and woody indicator plants. Herbaceous host indexing assays may be completed in a matter of weeks whereas woody indicator assays require a lengthier incubation period (up to two years) to complete (Rowhani *et al.* 2005). Herbaceous hosts are used to test for sap transmissible nepoviruses, whereas woody indicator plants are used to test for phloem limited viruses (Rowhani *et al.* 2005). Laboratory methods; including electron microscopy and molecular tests (PCR); can also be used to detect grapevine infecting viruses.

- As woody indexing is time consuming, molecular tests are recommended to replace woody indexing, thereby leading to a reduction of the PEQ growth period from a minimum of 24 months to a minimum of 16 months.
- Molecular tests (PCR, RT-PCR, and qPCR) target the genetic material of plant pathogens and specifically test for molecular sequences that are unique to a particular pathogen. Molecular tests can be used for the detection of grapevine pathogens because each pathogen has its own unique genetic code (Van Guilder *et al.* 2008). However, these molecular tests may not detect different strains or variants of a particular virus. Therefore, a combination of biological indexing and molecular tests is recommended to increase the likelihood of detecting viruses and their variants.

Effective and robust diagnostic methods based on a well established combination of biological, serological, and/or molecular tests are required to detect viruses. Recommended mandatory general methods for viruses include:

- Electron microscopy for the identified viruses.
- Herbaceous host indexing for nepoviruses (*Chenopodium quinoa*, *Chenopodium amaranticolor*, *Cucumis sativus* and other species may be used as herbaceous indicators).
- Generic molecular tests for *Ampelovirus*, *Ilarvirus*, *Maculavirus*, *Nepovirus* and *Vitivirus*.
- Specific RT-PCR for GVB (strains associated with corky bark).
- Specific RT-PCR for GRBaV (Al Rwahnih *et al.* 2012a).

Recommended diagnostic methods for virus groups are detailed below.

### Ampeloviruses

- Detection of ampeloviruses will include, but will not be limited to, the following tests:
  - Mandatory generic PCR for GLRaV-6, 10, 11 using the dHSP-nest2 / LR5 clusdoL primers (Maliogka *et al.* 2008b); and
  - Mandatory specific one step RT-PCR for GLRaV-7 using the primer pair LR7-F/ LR7-R (Engel *et al.* 2008).

### Ilarviruses

- Detection of ilarviruses will include, but will not be limited to, the following tests:
  - Herbaceous host indexing, including *Cucumis sativus* or *Nicotiana glutinosa* (*Grapevine line pattern virus*); and
  - Mandatory genus specific nested PCR for ilarviruses (GAMV, GLPV) using the Ilar2F5/Ilar2R9 primer pair (Untiveros *et al.* 2010).

### Maculaviruses

- Detection of maculaviruses will include, but will not be limited to, the following test:
  - Mandatory genus specific nested PCR for maculaviruses (GAMaV, GRGV) using the primer pair RD1/RGAP (Sabanadzovic *et al.* 2000).

Nepoviruses

- Herbaceous host indexing using a range of herbaceous indicators, that include but are not limited to:
  - *Chenopodium quinoa* (ArMV, BLMoV, CLRV, GARMV, GBLV, GCMV, GDefV, GFLV, GTRSV, PRMV, RpRSV, SLRV, TBRV, ToRSV);
  - *Chenopodium amaranticolor* (ArMV, BLMoV, CLRV, GARMV, GBLV, GCMV, GDeF, GFLV, PRMV, RpRSV, SLRV, TBRV, ToRSV);
  - *Cucumis sativus* (AILV, SLRV, TBRV, ToRSV); and
- Generic PCR testing for nepoviruses (Digiario *et al.* 2007; Wei and Clover 2008). If nepoviruses are detected, then virus specific tests must be performed. Virus specific tests may include (but are not limited to):
  - ArMV and GFLV using the primer pair M2/M3 (Wetzel *et al.* 2002);
  - CLRV using the primer pair CLRV-5/CLRV-3 (Werner *et al.* 1997);
  - GARSV using the primer pair A34-1/ A34-2 (Gokalp *et al.* 2003);
  - GCMV and TBRV using the primer pair P1/P2 (Le Gall *et al.* 1995);
  - GDefV using the primer pair N66-1/ N66-2 (Cigsar *et al.* 2003);
  - PRMV using the primer pair PRMVV1/ PRMVC1 (Kheder *et al.* 2004);
  - RpRSV using the primer pair RpRSVF1/ RpRSVR1 (Ochoa-Corona *et al.* 2006);
  - SLRSV using the primer pair SLRSV-5D / SLRSV-3D (Faggioli *et al.* 2002); and
  - ToRSV using the primer pair D1/U1 (Griesbach 1995).

Vitiviruses

- Detection of vitiviruses will include, but will not be limited to, the following test:
  - Mandatory specific RT-PCR for GVB (strains associated with corky bark) (Minafra and Hadidi 1994).

Tombusviruses

- Detection of tombusviruses will include, but will not be limited to, the following tests:
  - Mandatory genus specific nested PCR for *Tombusvirus* (PetAMV) using the pairs TomCPR/TomCPR (Russo *et al.* 2002) or TBSVGralF1/ TBSVGralR1 (Harris *et al.* 2006).

Plant material will be tested for other viruses using pathogen specific PCR tests if symptoms develop during growth in PEQ.

A summary of recommended grapevine virus indexing procedures is provided in Table 3.2.



**Table 3.2 Recommended grapevine virus indexing procedures**

Pathogen type	Mandatory tests			Additional tests <sup>5</sup>	Reference(s)
	Electron microscopy	Herbaceous indexing	PCR or RT-PCR		
<i>Arabidopsis mosaic virus</i> (ArMV) – grape strain	✓	✓	◆	RT-PCR	Wetzel <i>et al.</i> 2002
<i>Artichoke Italian latent virus</i> (AILV)		✓	◆	RT-PCR	Minafra <i>et al.</i> 1994
<i>Blueberry leaf mottle virus</i> (BLMoV) New York strain		✓	◆		Digiario <i>et al.</i> 2007
<i>Cherry leafroll virus</i> (CLRV) – grape isolate		✓	◆	RT-PCR	Werner <i>et al.</i> 1997
<i>Grapevine ajinashika virus</i> (GAgV) <sup>6</sup>					
<i>Grapevine Anatolian ringspot virus</i> (GARSV)		✓	◆	RT-PCR	Gokalp <i>et al.</i> 2003
<i>Grapevine angular mosaic-associated virus</i> (GAMaV)			★		Sabanadzovic <i>et al.</i> 2000
<i>Grapevine asteroid mosaic associated virus</i> (GAMV)			★		Untiveros <i>et al.</i> 2010
<i>Grapevine berry inner necrosis virus</i> (GINV)		✓	□		Yoshikawa <i>et al.</i> 1997
<i>Grapevine Bulgarian latent virus</i> (GBLV)		✓	◆		Digiario <i>et al.</i> 2007
<i>Grapevine chrome mosaic virus</i> (GCMV)		✓	◆		Le Gall <i>et al.</i> 1995
<i>Grapevine deformation virus</i> (GDefV)		✓	◆	RT-PCR	Cigsar <i>et al.</i> 2003
<i>Grapevine fanleaf virus</i> (GFLV)		✓	◆	RT-PCR	Wetzel <i>et al.</i> 2002
<i>Grapevine leafroll associated virus</i> (GLRaV–6,10, 11)			●		Maliogka <i>et al.</i> 2008b
<i>Grapevine leafroll associated virus</i> (GLRaV–7)			□		Engel <i>et al.</i> 2008
<i>Grapevine line pattern virus</i> (GLPV)		✓	★		Untiveros <i>et al.</i> 2010
<i>Grapevine Pinot gris virus</i> (GPGV)				RT-PCR	Cho <i>et al.</i> 2013
<i>Grapevine red blotch-associated virus</i> (GRBaV)			✓		Al Rwahnih <i>et al.</i> 2012a
<i>Grapevine red globe virus</i> (GRGV)			★		Sabanadzovic <i>et al.</i> 2000
<i>Grapevine rupestris vein feathering virus</i> (GRVFV)		✓	□		Abou Ghanem-Sabanadzovic <i>et al.</i> 2003
<i>Grapevine syrah virus-I</i> (GSyV-I)			★		Sabanadzovic <i>et al.</i> 2000
<i>Grapevine Tunisian ringspot virus</i> (GTRSV)		✓	◆		Digiario <i>et al.</i> 2007
<i>Grapevine virus B</i> (corky bark strains) (GVB)			Δ		Minafra and Hadidi 1994
<i>Grapevine virus E</i> (GVE)			⬆		Dovas and Katis 2003
<i>Grapevine virus F</i> (GVF)				RT-PCR	Al Rwahnih <i>et al.</i> 2012b
<i>Peach rosette mosaic virus</i> (PRMV)		✓	◆		Kheder <i>et al.</i> 2004
<i>Petunia asteroid mosaic virus</i> (PeAMV)			⊠		Russo <i>et al.</i> 2002; Harris <i>et al.</i> 2006
<i>Raspberry ringspot virus</i> (RpRSV) – grapevine strain		✓	◆		Ochoa-Corona <i>et al.</i> 2006
<i>Sowbane mosaic virus</i> (SoMV) – grape infecting strain		✓			
<i>Strawberry latent ringspot virus</i> (SLRSV)		✓	◆		Faggioli <i>et al.</i> 2002
<i>Tobacco necrosis virus</i> (TNV) – grape strain		✓	◆		Digiario <i>et al.</i> 2007
<i>Tomato black ring virus</i> (TBRV)		✓	◆		Le Gall <i>et al.</i> 1995
<i>Tomato ringspot virus</i> (ToRSV)		✓	◆		Griesbach 1995

<sup>5</sup> If disease symptoms develop<sup>6</sup> No PCR or commercial ELISA test are available, but the disease could possibly be diagnosed based on electron microscopy if the virus is detected then biological indexing onto the cultivar Koshu (Martelli 1993) will be required.

- ◆ Generic *Nepovirus* PCR (Digiario *et al.* 2007)
- ☒ Genus specific nested PCR for *Tombusvirus* (Russo *et al.* 2002 or Harris *et al.* 2006)
- Specific RT-PCR test (Yoshikawa *et al.* 1997; Abou Ghanem-Sabanadzovic *et al.* 2003; or Engel *et al.* 2008)
- Genus specific PCR for *Ampelovirus* (Maliogka *et al.* 2008b)
- ★ Genus specific nested PCR for ilarviruses (Untiveros *et al.* 2010)
- ☆ Genus specific nested PCR for maculaviruses (Sabanadzovic *et al.* 2000)
- Δ Strain specific PCR (Minfra and Hadidi 1994)
- ⌚ Generic PCR test for *Vitivirus* (Dovas and Katis 2003)
- Virus specific test for *Grapevine red blotch-associated virus* (Al Rwahnih *et al.* 2012a)

Plant Biosecurity acknowledges that advances in serological or molecular techniques is an on-going process and therefore the recommended PCR tests can be replaced when more up-to-date testing procedures are validated.

### 3.1.2 Tissue cultures (microplantlets)

It is recommended that imported tissue cultures (microplantlets) should be well rooted prior to arrival as this helps in their establishment out of agar into the growth media.

#### Mandatory on-arrival inspection

The imported tissue cultures (microplantlets) must be subject to mandatory on-arrival inspection to verify freedom from bacterial and fungal infection, disease symptoms, live insects and other extraneous contamination of quarantine concern.

The agar culture media must be clear and not contain antibiotics. If diseased material is detected during on-arrival inspection, the material must be held and referred to a plant pathologist for identification/risk assessment.

#### Mandatory culturing

It is recommended that direct culturing be undertaken to screen imported tissue cultures (microplantlets) for bacterial pathogens.

#### Mandatory growth in PEQ facilities and pathogen screening

The imported tissue culture (microplantlets) must be grown in a closed government PEQ facility for a minimum of 12 months for pathogen screening.

It is recommended that mandatory hot water treatment of plants established from tissue cultures (that requires two years) be replaced by mandatory PCR for detecting *Xylella fastidiosa*. Additionally, mandatory indexing for corky bark associated virus using LN 33 is replaced by a mandatory PCR.

The introduction of mandatory molecular testing leads to a reduction of the PEQ period. Therefore, it is recommended tissue cultures (microplantlets) be grown in a PEQ facility for a minimum of 12 months for pathogen screening, including biological indexing and molecular tests (Table 3.1 [bacteria and phytoplasma] and 3.2 [virus indexing]).

### 3.1.3 Seed for sowing (non-approved sources)

Although several nepoviruses are recorded on grapevines, not all of them are seed-borne (Richardson 1990). Seed-borne viruses of grapevine include ArMV, BLMoV-NY, GAMaV, GCMV, GBLV, GFLV, GLPV, GRSPaV, PRMV, TBRV and ToRSV (Uyemoto 1975; Uyemoto *et al.* 1977; Martelli 1978; Lazar *et al.* 1990; Richardson 1990; Lehoczyk *et al.* 1992; Girgis *et al.* 2009). Therefore, during growth in PEQ, seedlings must be visually inspected for symptoms of viruses.

**Mandatory on arrival inspection**

The imported grapevine seed must be subject to mandatory on-arrival inspection to verify freedom from live insects, soil, disease symptoms, prohibited seeds, other plant material (e.g. leaf, stem material, fruit pulp, pod material etc.), animal material (e.g. animal faeces, feathers etc.) and any other extraneous contamination of quarantine concern.

**Mandatory sodium hypochlorite treatment**

The imported grapevine seed must be subject to mandatory surface sterilisation with sodium hypochlorite treatment (1% NaOCl for 10 minutes).

**Mandatory seed fungicide treatment**

The imported grapevine seed must be subject to mandatory fungicidal treatment (Thiram) prior to sowing.

**Mandatory growth in PEQ facilities**

The imported grapevine seed must be grown in a closed government PEQ facility for a minimum of 9 months as growth in the PEQ facility for three months may not be sufficient for plant establishment from seed and to complete pathogen screening.

**Mandatory virus testing**

It is recommended that in addition to visual inspection for symptoms during growth in PEQ facility, the following procedures are required to detect viruses:

- Electron microscopy is mandatory for the identified seed-borne viruses.
- Herbaceous host indexing and generic PCR for nepoviruses is mandatory. Detection of nepoviruses on indicator plants will require further testing, including virus specific PCR, RT-PCR, or qPCR (Table 3.2).
- Detection of ilarviruses will include, but will not be limited to, the following tests:
  - Herbaceous host indexing; and
  - Mandatory molecular testing PCR (Table 3.2).

## **3.2 Propagative material from approved sources)**

Existing measures for grapevine propagative material from approved sources are recommended to continue and additional requirements are not recommended. However, recommended changes to import requirements for material from non-approved sources will also apply to material from approved sources (e.g. the PEQ period will be reduced from 24 months to 16 months for dormant cuttings and 12 months for tissue cultures).

If the required pathogen screening is completed at an overseas approved source then Plant Biosecurity may further reduce the recommended PEQ growth requirement.

### **3.2.1 Seed for sowing (approved sources)**

Currently, seeds sourced from approved sources (Foundation Plant Services, University of California, USA) are permitted entry into Australia. The existing policy requires certification that the seeds were sourced from mother plants grown in the USA that were tested and found free of *Arabis mosaic nepovirus* (ArMV), *Blueberry leaf mottle nepovirus* (BLMV), *Grapevine Bulgarian latent nepovirus* (GBLN), *Peach rosette mosaic nepovirus* (PRMN), *Raspberry ringspot nepovirus* (RpRSV), *Strawberry latent ringspot nepovirus* (SLRSV), *Tomato black ring nepovirus* (TBRV) and *Tomato ringspot nepovirus* (ToRSV).

As part of the review of policy, the current seed-borne list of viruses associated with grapevine seed was revised and updated.

- *Grapevine angular mosaic-associated virus* (GAMaV) and *Grapevine fanleaf virus* (GFLV) were added to the list as these seed-borne viruses are present in the USA;
- *Raspberry ringspot nepovirus* (RpRSV) and *Strawberry latent ringspot nepovirus* (SLRSV) were removed from the list as there is no published evidence that these viruses are seed-borne in grapevine; and
- *Tobacco ringspot nepovirus* (TRSV) was removed from the list as it is present in Australia.

Based on this review, the new recommended conditions for grapevine seeds from Foundation Plant Services, University of California, USA includes:

- an import permit;
- a Phytosanitary Certificate (seed was sourced from virus tested mother plants free of '*Arabis mosaic nepovirus*(ArMV), *Blueberry leaf mottle nepovirus* (BLMV), *Grapevine angular mosaic-associated virus* (GAMaV), *Grapevine Bulgarian latent nepovirus* (GBLN), *Grapevine fanleaf virus* (GFLV), *Peach rosette mosaic nepovirus* (PRMN), *Tomato black ring nepovirus* (TBRV) and *Tomato ringspot nepovirus* (ToRSV)'; and
- on-arrival inspection of seed to verify freedom from soil, disease symptoms and other extraneous contamination of quarantine concern.

Specific conditions; including surface sterilization (T9371), fungicidal treatment (T9420) and release from quarantine; are recommended to continue.

## 4 Framework for approval of high health sources and production requirements

---

### 4.1 Framework for approval of high health sources

Foundation Plant Services, California, USA is currently the only source approved to supply pathogen tested grapevine propagative material to Australia. However, Plant Biosecurity will consider requests for approval of other overseas sources (e.g. institutions, NPPOs) based on their compliance with international standards and a rigorous examination of the recommended facilities. The key factors for approval of high health sources include:

- **Capacity for National Authority oversight**—facilities producing pathogen tested propagative material must be authorized/approved or operated directly by the National Plant Protection Organization (NPPO), as import conditions routinely require phytosanitary certification to be provided by the NPPO.
- **Capacity to produce pathogen tested propagative material**—facilities must demonstrate their capacity to produce and maintain high health plant material through appropriate disease screening/testing and monitoring.
- **Capacity to meet containment and security requirements**—facilities for the establishment of pest-free propagative material and testing for pest freedom must be subject to strict physical containment and operational requirements to prevent contamination or infestation of material.
- **Audits and inspections**—all facilities producing pathogen tested propagative material should be officially audited by DAFF to ensure that they continue to meet Australia's requirements.
- **Identity preservation systems**—all facilities must be able to demonstrate their ability to maintain adequate and verifiable safeguards to ensure that propagative material undergoing post-entry quarantine procedures are not diverted, contaminated or intermingled with other material during and following completion of the quarantine measures.
- **On arrival verification**—the requirement for the health status of all consignments of high health propagative material to be verified on-arrival through supporting documentation (e.g. Phytosanitary Certificate, NPPO reports, audit report etc.) and testing as required.

Based on this framework, Australia will consider replacing the conditions for on-arrival pathogen screening with an equivalent set of conditions for approved sources. The key elements of material produced in approved sources are:

- Pathogen screening/testing must be equivalent to Australia's post-entry quarantine screening/testing;
- Each consignment must have a certificate of testing with results, dates and details of the testing methods used issued by the approved source and certified by the NPPO of the exporting country;
- Imported propagative material may be subjected to verification testing for a range of quarantine pathogens during growth in a closed government PEQ facility; and
- Where any accredited source does not undertake the complete range of pathogen screening/testing required, those missing tests will be performed during growth in a closed government PEQ facility in Australia.

## 5 Conclusion

---

The findings of this final review of policy are based on a comprehensive analysis of the scientific literature. As part of this revision, the quarantine status of grapevine pathogens was reviewed and several new pests of quarantine concern were identified. Consequently, Plant Biosecurity evaluated the appropriateness of existing risk management measures for the identified risks and recommended additional measures where required.

### **Recommended risk management measures**

The recommended risk management measures for propagative material are detailed below.

#### **All sources (unknown health status)**

##### **Dormant cuttings**

- Mandatory on-arrival inspection fumigation; hot water treatment; and surface sterilisation;
- Mandatory growth in a closed government PEQ facility for a minimum period of 16 months for pathogen screening (visual observation; culturing; and electron microscopy); and
- Active pathogen testing through herbaceous host indexing and molecular tests including, but not limited to, PCR or ELISA.

##### **Tissue cultures (microplantlets)**

- Mandatory on-arrival inspection;
- Mandatory growth in a closed government PEQ facility for a minimum period of 12 months for pathogen screening (visual observation; culturing; and electron microscopy); and
- Active pathogen testing through herbaceous host indexing and molecular tests including, but not limited to, PCR or ELISA.

##### **Seed**

- Mandatory on-arrival inspection, surface sterilisation; fungicidal treatment; and growth in a closed government PEQ facility for a minimum period of nine months for pathogen screening (visual observation and electron microscopy); and
- Active pathogen testing through herbaceous host indexing and molecular tests including, but not limited to, PCR.

#### **Approved sources (high health sources)**

Foundation Plant Services, California, USA is currently the only source approved to supply pathogen tested grapevine propagative material to Australia. However, Plant Biosecurity will consider requests for approval of other overseas sources (e.g. institutions, NPPOs etc), based on the framework recommended in this review. If the requirements of the framework are met, Plant Biosecurity will consider replacing the existing conditions with an alternative set of conditions for approved sources.

The recommended changes to import requirements for dormant cuttings and tissue cultures from non-approved sources will also apply to material from approved sources (e.g. the PEQ period will be reduced to 16 months for dormant cuttings and 12 months for tissue cultures). Seed for sowing from approved sources is currently not subject to PEQ and this is recommended to continue.

# Appendices

## Appendix A: Initiation and pest categorisation of pests associated with *Vitis* species worldwide

Initiation identifies the pests that occur on *Vitis* species, their status in Australia and their pathway association. In this assessment, **pathway** is defined as *Vitis* propagative material (one-year-old dormant cuttings, seed and tissue culture). Restricting budwood to one-year-old material reduces the risk of opportunistic wound pathogens and wood rots. In addition, dormant cuttings are semi-hardwood and have not developed mature bark. Therefore, pests associated with the hardwood and mature bark of older grapevines is not considered to be on the pathway. As grapevine cuttings are harvested when they are dormant, pests associated with new plant growth (e.g. developing buds, new shoots, tendrils and fruit) do not occur on the pathway. Dormant grapevine cuttings are also free of roots and leaves, consequently pests associated with roots and leaves are not considered to be on the pathway. Please note that the 'Potential to be on pathway' column usually specifies the association of pests with dormant cuttings. Bacteria, phytoplasmas and viruses occurring on tissue culture are considered to be the same as those occurring on dormant cuttings. Seeds are only referred to in the pathway column if the pest is known to be associated with seeds.

Pest categorisation identifies the potential for pests associated with grapevine propagative material to enter, establish, spread and cause economic consequences in Australia, to determine if they qualify as quarantine pests.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<b>ARTHROPODS</b>					
<b>ACARI (mites)</b>					
<i>Brevipalpus californicus</i> Banks 1904 [Acari: Tenuipalpidae]	Yes (Naumann 1993)	Assessment not required			
<i>Brevipalpus chilensis</i> Baker 1949 [Acari: Tenuipalpidae]	Not known to occur	<b>Yes:</b> Mites lay eggs on the young shoots and leaves or in the unopened buds of grapevines (González 1968, 1983). This mite overwinters as fertilised females, usually in colonies under the bark crevices of host plants (Jeppson <i>et al.</i> 1975). Therefore, semi-hardwood dormant cuttings provide a pathway for this mite.	<b>Yes:</b> This mite has established in areas with a wide range of climatic conditions (Waterhouse and Sands 2001). It has a wide host range (Waterhouse and Sands 2001), has four to five generations per year (González 1968) and can spread naturally in infested propagative material. Therefore, this mite has the potential for establishment and spread in Australia.	<b>Yes:</b> This mite is recognised as a significant pest of grapes in Chile and causes as much as 30% crop loss (González 1983). This mite is regarded as a quarantine pest by trading partners. Therefore, this mite may potentially increase production costs by triggering trading partners to issue specific control measures. As such, this mite has the potential for significant economic consequences in Australia.	<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Brevipalpus lewisi</i> McGregor 1949 [Acari: Tenuipalpidae]	Yes (Naumann 1993)	Assessment not required			
<i>Brevipalpus obovatus</i> Donnadieu 1875 [Acari: Tenuipalpidae]	Yes (Naumann 1993)	Assessment not required			
<i>Brevipalpus phoenicis</i> Geijskes 1936 [Acari: Tenuipalpidae]	Yes (Naumann 1993)	Assessment not required			
<i>Calepitrimerus vitis</i> Nalepa 1905 [Acari: Eriophyidae]	Yes (Naumann 1993)	Assessment not required			
<i>Colomerus vitis</i> Pagenstecher 1857 strain a [Acari: Eriophyidae]	Yes (James and Whiteney 1993)	Assessment not required			
<i>Colomerus vitis</i> Pagenstecher 1857 strain b [Acari: Eriophyidae]	Yes (James and Whiteney 1993)	Assessment not required			
<i>Colomerus vitis</i> Pagenstecher 1857 strain c [Acari: Eriophyidae]	Not known to occur	<b>Yes:</b> Mites lay eggs on dormant buds (Carew <i>et al.</i> 2004) and overwinter as adults inside grapevine buds (Jeppson <i>et al.</i> 1975). Therefore, dormant cuttings may provide a pathway for this mite.	<b>Yes:</b> This mite has established in areas with a wide range of climatic conditions (Afonin <i>et al.</i> 2008). It has several generations per year (Jepson <i>et al.</i> 1975; Carew <i>et al.</i> 2004) and can independently spread in infested plant material and by human activities (Jeppson <i>et al.</i> 1975; Gonzalez 1983). Therefore, this mite has the potential for establishment and spread in Australia.	<b>Yes:</b> <i>Colomerus vitis</i> is associated with short shoot syndrome of grape vines (Bernard <i>et al.</i> 2005). This mite causes deformation of the primordial bud cluster, distortion of the basal leaves, stunting of the main growing point of the buds and often death of the overwintering buds (Pfeiffer and Schultz 1986). Bud burst failure and high yield losses have been attributed to this mite (Walton <i>et al.</i> 2007). This mite may cause yield losses of up to 56% when uncontrolled (Dennill 1991). Therefore, this mite has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Bryobia praetiosa</i> Koch 1836 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Bryobia rubrioculus</i> Scheuten 1857 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Eotetranychus carpini</i> Oudemans 1905 [Acari: Tetranychidae]	Not known to occur	No: Tetranychid mites belonging to the genus <i>Eotetranychus</i> are foliage feeders and lay eggs on leaves (Jeppson <i>et al.</i> 1975); Karban <i>et al.</i> 1991; EPPO 2006). These mites overwinter as females under the bark and become active with the new plant growth (HYPPZ 1998). Therefore, foliage free, semi-hardwood dormant cuttings do not provide a pathway for these mites.	Assessment not required		
<i>Eotetranychus lewisi</i> (McGregor 1943) [Acari: Tetranychidae]	Not known to occur		Assessment not required		
<i>Eotetranychus sexmaculatus</i> Riley (1890) [Acari: Tetranychidae]	Yes (CSIRO 2005)		Assessment not required		
<i>Eotetranychus willametti</i> McGregor 1917 [Acari: Tetranychidae]	Not known to occur		Assessment not required		
<i>Eutetranychus orientalis</i> Klein (1936) [Acari: Tetranychidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Oligonychus coffeae</i> Nietner 1861 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Oligonychus mangiferus</i> Rahman & Sapro 1940 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Oligonychus punicae</i> Hirst 1926 [Acari: Tetranychidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Oligonychus vitis</i> Zaher & Shehata 1965 [Acari: Tetranychidae]	Not known to occur	No: Tetranychid mites belonging to <i>Oligonychus</i> genus are foliage feeders (Jeppson <i>et al.</i> 1975; Gonzalez 1983; Gutierrez and Schicha 1983). Therefore, foliage free dormant cuttings do not provide a pathway for these mites.	Assessment not required		
<i>Oligonychus yothersi</i> McGregor 1914 [Acari: Tetranychidae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Panonychus citri</i> McGregor 1916 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Panonychus ulmi</i> Koch 1836 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Petrobia latens</i> Müller 1776 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Polyphagotarsonemus latus</i> Banks 1904 [Acari: Tarsonemidae]	Yes (Naumann 1993)	Assessment not required			
<i>Tetranychus cinnabarinus</i> (Boisduval) Boudreaux 1956 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Tetranychus desertorum</i> Banks 1900 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Tetranychus kanzawai</i> Kishida 1927 [Acari: Tetranychidae]	Yes (Seeman and Beard 2011)	Assessment not required			
<i>Tetranychus ludeni</i> Zacher 1913 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<i>Tetranychus pacificus</i> McGregor 1919 [Acari: Tetranychidae]	Not known to occur	No: These species feed and oviposit on the under surface of leaves (Jeppson <i>et al.</i> 1975; McLaren <i>et al.</i> 1999; Rieger 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this mite.	Assessment not required		
<i>Tetranychus telarius</i> (Linnaeus 1758) [Acari: Tetranychidae]	Yes (PHA 2001)	Assessment not required			
<i>Tetranychus urticae</i> Koch 1836 [Acari: Tetranychidae]	Yes (Naumann 1993)	Assessment not required			
<b>COLEOPTERA (beetles, weevils)</b>					
<i>Acalolepta vastator</i> Newman 1847 [Coleoptera: Cerambycidae]	Yes (Naumann 1993)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Acrothinium gaschkevitschii</i> (Motschulsky 1860) [Coleoptera: Chrysomelidae]	Not known to occur	No: This species feeds externally on the buds, leaves and flowers of grapevines (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Adoretus sinicus</i> Burmeister 1855 [Coleoptera: Scarabaeidae]	Not known to occur	No: These scarabaeid beetles lay eggs in the soil, larvae feed on roots and adults feed on the leaves of grapevines (NIIR 2004; Zhang 2005). Therefore, root free and foliage free dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Adoretus versutus</i> Harold 1869 [Coleoptera: Scarabaeidae]	Not known to occur		Assessment not required		
<i>Agriotes lineatus</i> Linnaeus 1767 [Coleoptera: Elateridae]	Not known to occur	No: This species lays eggs on or in the soil and larvae feed on roots (Bournier 1976). Therefore, root free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Anomala corpulenta</i> Motschulsky 1854 [Coleoptera: Scarabaeidae]	Not known to occur	No: These scarabaeid beetles lay eggs in the soil, larvae feed on the roots and adults feed on leaves and flowers (Bhuiyan and Nishigaki 1997; Larsson <i>et al.</i> 2001; Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Anomala cuprea</i> Hope 1839 [Coleoptera: Scarabaeidae]	Not known to occur		Assessment not required		
<i>Altica grvida</i> Blackburn 1896 [Coleoptera: Chrysomelidae]	Yes (AFD 2008)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Altica ampelophaga</i> Guérin-Meneville-Menevil 1858 [Coleoptera: Chrysomelidae]	Not known to occur	No: Grape flea beetles overwinter as adults under the soil surface, in wood crevices, under stones, sticks or logs and in or around vineyards (Galvan <i>et al.</i> 2007). These beetles emerge in early spring when grapevine buds begin to swell and lay eggs either at the base of the buds, on the buds and bark crevices of grapevines (Benvenuti and Lucchi 2005; Galvan <i>et al.</i> 2007), or on the underside of leaves (Alford 2007). Therefore, foliage free, semi-hardwood dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Altica chalybea</i> Illiger 1807 [Coleoptera: Chrysomelidae]	Not known to occur		Assessment not required		
<i>Altica torquata</i> (LeConte 1859) [Coleoptera: Chrysomelidae]	Not known to occur		Assessment not required		
<i>Altica woodsi</i> Isely 1920 [Coleoptera: Chrysomelidae]	Not known to occur		Assessment not required		
<i>Ampelogypter ampelopsis</i> Riley 1869 [Coleoptera: Curculionidae]	Not known to occur	No: Grape cane gallmakers overwinter as adults in the debris on the ground (Riedl and Taschenberg 2008). Grape cane gallmakers lay eggs in new canes in spring and adults start emerging in midsummer (Riedl and Taschenberg 2008). No life stage is associated with dormant cuttings. Therefore, these species are not on the pathway.	Assessment not required		
<i>Ampelogypter ater</i> LeConte 1876 [Coleoptera: Curculionidae]	Not known to occur		Assessment not required		
<i>Ampelogypter sesostris</i> LeConte 1876 [Coleoptera: Curculionidae]	Not known to occur		Assessment not required		
<i>Anoplistes halodendri</i> Kozlovi (Semenov & Znojdo 1934) [Coleoptera:]	Not known to occur	No: The wood-boring larvae of this beetle damage	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Cerambycidae]		grapevines and other woody plants (Luo <i>et al.</i> 2005). However, semi-hardwood dormant cuttings are not preferred sites for egg laying for this species. Therefore, dormant cuttings do not provide a pathway for this species.			
<i>Anoplophora glabripennis</i> Motschulsky 1853 [Coleoptera: Cerambycidae]	Not known to occur	No: The wood-boring larvae of this beetle damage grapevines and other woody plants (Lingafelter and Hoebeke 2002). Adults also feed on the leaves, stems and bark of many woody plant species (Yang <i>et al.</i> 1995). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, foliage free, semi-hardwood dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Asynonychus cervinus</i> Boheman (1840) [Coleoptera: Curculionidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Athlia rustica</i> Erichson 1835 [Coleoptera: Scarabaeidae]	Not known to occur	No: This species feeds externally on leaves, buds and flowers of host plants (Gonzalez 1983). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Atrichonotus taeniatulus</i> Berg (1881) [Coleoptera: Curculionidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Aulacophora femoralis chinensis</i> Weise 1923 [Coleoptera: Chrysomelidae]	Not known to occur	No: Adults feed on the leaves of grapes, pears, apples and leaf vegetables while the larvae live in the soil and feed on young plant roots (Li 2004). Therefore, foliage and root free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Bostrychopsis jesuita</i> Fabricius 1775 [Coleoptera: Bostrichidae]	Yes (PHA 2001)	Assessment not required			
<i>Bromius obscurus</i> Linnaeus 1758 [Coleoptera: Chrysomelidae]	Not known to occur	No: Larvae of this species feed on grapevine roots while adults feed on leaves, green bark of canes and cut shallow grooves in berries (Peacock 1992). Eggs are laid in clusters on old loose bark in crevices (BCMAL 2010). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Byctiscus betulae</i> (Linnaeus, 1758) [Coleoptera: Rhynchitidae]	Not known to occur	No: These weevils mostly feed on leaves (Bournier 1976; Zhang 2005) and lay eggs inside of rolled leaves (Trdani and Valič 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Byctiscus lacunipennis</i> (Jekel 1860) [Coleoptera: Rhynchitidae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Callideriphus laetus</i> Blanchard 1851 [Coleoptera: Cerambycidae]	Not known to occur	No: This species primarily feeds on downed logs, stumps and dead or dying branches (Klein-Koch and Waterhouse 2000). Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Carpophilus dimidiatus</i> Fabricius 1992 [Coleoptera: Nitidulidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Carpophilus hemipterus</i> Linnaeus 1758 [Coleoptera: Nitidulidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Carpophilus humeralis</i> Fabricius 1758 [Coleoptera: Nitidulidae]	Yes (Hossain and Williams 2003)	Assessment not required			
<i>Cerasphorus albofasciatus</i> (Laporte and Gory 1835) [Coleoptera: Cerambycidae]	Not known to occur	No: This species is a trunk borer (MAF 2009). Borers generally require thick wood for egg laying and development (Goodwin 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Ceresium sinicum</i> ornaticolle Pic 1907 [Coleoptera: Cerambycidae]	Not known to occur	No: Larvae of this species attack woody parts of grapevines as internal feeders (Luo <i>et al.</i> 2005). Borers generally require thick wood for egg laying and development (Goodwin 2005). Semi-hardwood dormant cuttings are not preferred	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.			
<i>Cerosterna scabrator</i> Fabricius 1781 [Coleoptera: Cerambycidae]	Not known to occur	No: This longicorn beetle attacks the main stem and branches of host plants. The female oviposits in the stem and larvae feed inside the stems (Ranga Rao <i>et al.</i> 1979). Borers generally require thick wood for egg laying and development (Goodwin 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Chlorophorus annularis</i> Fabricius 1787 [Coleoptera: Cerambycidae]	Yes (PHA 2001)	Assessment not required			
<i>Chlorophorus quatuordecimmaculatus</i> (Chevrolat 1863) [Coleoptera: Cerambycidae]	Not known to occur	No: Larvae bore through larger stems of grapevines while adults eat the flowers (Zhang 2005). Borers generally require thick wood for egg laying and development (Goodwin 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Colaspis brunnea</i> Fabricius 1798 [Coleoptera: Chrysomelidae]	Not known to occur	No: Larvae of this beetle feed on roots and the adults feed on foliage (Pfeiffer and Schultz 1986). Eggs are laid in the soil (Eaton 1978). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Colaspoides foveiventris</i> Lea 1926 [Coleoptera: Chrysomelidae]	Yes (Naumann 1993)	Assessment not required			
<i>Colaspoides heroni</i> Lea 1915 [Coleoptera: Chrysomelidae]	Yes (AFD 2008)	Assessment not required			
<i>Colaspoides picticornis</i> Lea 1915 [Coleoptera: Chrysomelidae]	Yes (PHA 2001)	Assessment not required			
<i>Coniontis parviceps</i> Casey 1890 [Coleoptera: Tenebrionidae]	Not known to occur	No: The larvae of soil-dwelling tenebrionid are root feeders (Allsopp 1980) and adults are bud and foliage feeders (Flaherty <i>et al.</i> 1992). Therefore, foliage and root free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Corticaria japonica</i> Reitter (1877) [Coleoptera: Latridiidae]	Yes (PHA 2001)	Assessment not required			
<i>Cotinis nitida</i> Linnaeus 1764 [Coleoptera: Scarabaeidae]	Not known to occur	No: Adults of this species feed on grape berries (Brown and Hudson 2005). Eggs are laid in the soil, where the hatching larvae then feed on decaying organic matter (OSU 2010). Therefore, dormant cuttings do not provide a pathway for	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		this species.			
<i>Craponius inaequalis</i> Say 1831 [Coleoptera: Curculionidae]	Not known to occur	No: The grape curculionid lays eggs in the fruit and developing larvae feed on seed and pulp (Bournier 1976). Therefore, dormant cuttings do not provide a pathway for this species. Larvae are not reported to feed internally in seeds; therefore seeds also do not provide a pathway for this species.	Assessment not required		
<i>Didymocantha obliqua</i> Newman (1840) [Coleoptera: Bostrichidae]	Yes (PHA 2001)	Assessment not required			
<i>Dilochrosis atripennis</i> MacLeay 1863 [Coleoptera: Scarabaeidae]	Yes (PHA 2001)	Assessment not required			
<i>Dryocoetops coffeae</i> (Eggers 1923) [Coleoptera: Scolytinae]	Not known to occur	No: Scolytine beetles are associated with woody plant products (Luo <i>et al.</i> 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.			
<i>Egiona viticola</i> Luo [Coleoptera: Curculionidae]	Not known to occur	No: This wood-boring pest of grapevines requires hardwood to lay eggs (Luo <i>et al.</i> 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		this species.			
<i>Fidia viticida</i> Walsh 1867 [Coleoptera: Chrysomelidae]	Not known to occur	No: Grape rootworm beetles lay eggs under the bark of grapevine trunks. Immature grubs feed on the roots and adults feed on grape foliage (Dennehy and Clark 2008). Therefore, semi-hardwood foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Gametis jucunda</i> (Faldermann 1835) [Coleoptera: Scarabaeidae]	Not known to occur	No: Larvae of this species feed on roots while adults feed on grapevine flowers (Zhang 2005). Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Glyptoscelis squamulata</i> Crotch 1873 [Coleoptera: Chrysomelidae]	Not known to occur	No: Adult beetles feed on buds, immature leaves and young flowers (Flint 2006). Eggs are laid in cracks and under the bark (Stern and Johnson 1984). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Hayashiclytus acutivittis</i> (Kraatz 1879) [Coleoptera: Cerambycidae]	Not known to occur	No: This cerambycid beetle is associated with grapevines (Zhang 2005). Cerambycid larvae generally feed internally on woody plant material, while adults feed on flowers or foliage (CSIRO	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		1991). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.			
<i>Heteronychus arator</i> Fabricius 1775 [Coleoptera: Scarabaeidae]	Yes (Naumann 1993)	Assessment not required			
<i>Holotrichia diomphalia</i> (Bates 1888) [Coleoptera: Scarabaeidae]	Not known to occur	No: The larvae of this Scarabaeid beetle feed on roots while adults feed on shoots, young leaves and flowers (AQSIQ 2007). Therefore, foliage and root free dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Holotrichia oblita</i> (Faldermann 1835) [Coleoptera: Scarabaeidae]	Not known to occur		Assessment not required		
<i>Hoplia callipyge</i> (LeConte 1856) [Coleoptera: Scarabaeidae]	Not known to occur	No: This species lays eggs in the soil (Perry 2002), larvae are root feeders and adults feed on leaves and flowers (Evans and Hogue 2006). Therefore, foliage and root free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Hypothenemus javanus</i> (Eggers 1908) [Coleoptera: Scolytinae]	Not known to occur	No: Scolytine beetles are associated with woody plant products (Luo <i>et al.</i> 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for			
<i>Hypothenemus erectus</i> Leconte 1876 [Coleoptera: Scolytinae]	Not known to occur				

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		this species.			
<i>Hypothenemus eruditus</i> Westwood 1836 [Coleoptera: Scolytinae]	Yes (PHA 2001)	Assessment not required			
<i>Leptopius robustus</i> (Olivier 1807) [Coleoptera: Curculionidae]	Yes (Naumann 1993)	Assessment not required			
<i>Limonius canus</i> Leconte 1853 [Coleoptera: Elateridae]	Not known to occur	No: Click beetle lays eggs in soil and newly hatched larvae feed on roots (Berry 1998). These pests overwinter as larvae or as recently developed adults in the soil. Adults feed on buds in spring (Bentley <i>et al.</i> 2008). Therefore, root free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Linda fraterna</i> Chevrolat 1852 [Coleoptera: Cerambycidae]	Not known to occur	No: Longicorn beetles attack mature trees (Smith 1996). Adult beetles lay eggs into crevices or cracks in the bark on the trunk or main branches of host plants (Smith 1996). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Liparetrus atriceps</i> Macleay 1864 [Coleoptera: Scarabaeidae]	Yes (PHA 2001)	Assessment not required			
<i>Listroderes difficilis</i> Germain 1895 [Coleoptera: Curculionidae]	Yes (Ronald and Jayma 1992)	Assessment not required			
<i>Listroderes costirostris</i> Schönherr 1826 [Coleoptera: Curculionidae]	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Macrodactylus subspinosus</i> Fabricius 1775 [Coleoptera: Scarabaeidae]	Not known to occur	No: Adults feed externally on flowers, buds, foliage and fruits (OARDC 2008) and eggs of this species are laid in the soil (McLeod and Williams 1990). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Maladera orientalis</i> (Motschulsky 1857) [Coleoptera: Scarabaeidae]	Not known to occur	No: Larvae feed on the roots of grapevines while adults feed on the young shoots, leaves, and flowers of grapes (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Melalqus confertus</i> LeConte [Coleoptera: Bostrichidae]	Not known to occur	No: This wood-boring beetle lays eggs in protected areas and emerging larvae bore into dead wood where they continue to feed (Flaherty <i>et al.</i> 1992). During bud swell, adults feed on buds and bore into canes directly through buds or burrow into the canes at the bud axils destroying the bud and weakening the twig (Flaherty <i>et al.</i> 1992). Strong wind can cause infested canes to twist and break at feeding sites (Flaherty <i>et al.</i> 1992). Dormant canes are not preferred for adult feeding and	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		larval boring, and therefore do not provide a pathway for this species.			
<i>Melolontha melolontha</i> Fabricius 1775 [Coleoptera: Scarabaeidae]	Not known to occur	No: This species lays eggs in the soil (AgroAtlas 2009c) and larvae feed on roots and other underground plant parts (Bournier 1976). Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Micrapate humeralis</i> (Blanchard 1851) [Coleoptera: Bostrichidae]	Not known to occur	No: Bostrichids are associated with hardwoods, shrubs and woody vines (Booth <i>et al.</i> 1990). Eggs are laid in vine trunks and hatching larvae penetrate into the wood and construct a gallery in which they live and feed (Gonzalez 1983). These species overwinter as larvae, pupae and adults. Semi-hardwood dormant cuttings are not the preferred site for adult feeding and larval boring. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Micrapate scabrata</i> (Erichson 1847) [Coleoptera: Bostrichidae]	Not known to occur		Assessment not required		
<i>Monolepta australis</i> Jacoby 1882 [Coleoptera: Chrysomelidae]	Yes (PHA 2001)	Assessment not required			
<i>Monolepta divisa</i> Blackburn 1888 [Coleoptera: Chrysomelidae]	Yes (PHA 2001)	Assessment not required			
<i>Naupactus leucoloma</i> Boheman 1840 [Coleoptera: Curculionidae]	Yes (PHA 2001)	Assessment not required			



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Naupactus xanthographus</i> Germar Sturm 1826 [Coleoptera: Curculionidae]	Not known to occur	No: Larvae of this species damage the roots and adults feed on the foliage of grapevines (Gonzalez 1983). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Neoclytus caprea</i> Say 1824 [Coleoptera: Cerambycidae]	Not known to occur	No: This species is associated with dead wood (Hovore 1983). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Neoterius mystax</i> Blanchard 1851 [Coleoptera: Bostrichidae]	Not known to occur	No: This opportunistic borer is found in trunks and branches of host plants (Gonzalez 1983). Bostrichids require hard wood for egg laying (Madge 2007). Semi-hardwood dormant cuttings are not preferred sites for egg laying and development. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Oides decempunctata</i> Billberg 1808 [Coleoptera: Chrysomelidae]	Not known to occur	No: <i>Oides</i> species lay eggs either in the soil or on the soil surface (Park <i>et al.</i> 2001) or beneath the bark (Joshi and Gupta 1988). Chrysomelid adults and larvae feed on	Assessment not required		
<i>Oides scutellata</i> Hope 1831 [Coleoptera: Chrysomelidae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		young foliage, flowers and roots (Booth <i>et al.</i> 1990). Therefore, root and foliage free dormant cuttings do not provide a pathway for these species.			
<i>Orthorhinus cylindrirostris</i> Schoenherr 1825 [Coleoptera: Curculionidae]	Yes (PHA 2001)	Assessment not required			
<i>Orthorhinus klugi</i> Boheman 1835 [Coleoptera: Curculionidae]	Yes (Farquhar and Williams 2000)	Assessment not required			
<i>Oryzaephilus surinamensis</i> Linnaeus 1758 [Coleoptera: Silvanidae]	Yes (PHA 2001)	Assessment not required			
<i>Otiorhynchus cribricollis</i> Gyllenhal 1834 [Coleoptera: Curculionidae]	Yes (Farquhar and Williams 2000)	Assessment not required			
<i>Otiorhynchus rugosostriatus</i> Goeze 1777 [Coleoptera: Curculionidae]	Yes (Naumann 1993)	Assessment not required			
<i>Otiorhynchus singularis</i> Linnaeus 1829 [Coleoptera: Curculionidae]	Not known to occur	No: This species lays eggs at a shallow depth in the soil and hatching larvae feed on roots (Alford 2007). Adults of this species feed externally on buds (Alford 2007). Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Otiorhynchus sulcatus</i> Germar 1824 [Coleoptera: Curculionidae]	Yes (Naumann 1993)	Assessment not required			
<i>Paracotalpa ursina</i> Horn 1867 [Coleoptera: Scarabaeidae]	Not known to occur	No: This species is an external feeder of buds and very young shoots (Pimentel 2007). Therefore, foliage free	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		dormant cuttings do not provide a pathway for this species.			
<i>Paraphloeostiba gayndahensis</i> MacLeay 1873 [Coleoptera: Staphylinidae]	Yes (Thayer 2001)	Assessment not required			
<i>Pelidnota punctata</i> (Linnaeus 1758) [Coleoptera: Scarabaeidae]	Not known to occur	No: Larvae of grapevine beetles feed and live in decaying hardwood stumps, roots and logs, and adults feed on foliage (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Peritelus sphaeroides</i> (Germar 1824) [Coleoptera: Curculionidae]	Not known to occur	No: This bud weevil lays eggs in the soil and hatching larvae feed on roots. Adults attack buds, young foliage and flowers (Alford 2007). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Phlyctinus callosus</i> Boheman 1834 [Coleoptera: Curculionidae]	Yes (Farquhar and Williams 2000)	Assessment not required			
<i>Phymatodes albicinctus</i> Bates 1873 [Coleoptera: Cerambycidae]	Not known to occur	No: Larvae of this species feed internally on woody parts of the grapevine (Luo <i>et al.</i> 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		this species.			
<i>Phymatodes mediofasciatus</i> Pic 1933 [Coleoptera: Cerambycidae]	Not known to occur	No: Larvae of this species feed internally on woody parts of the grapevine (Cherepanov 1991). Semi-hardwood dormant cuttings are not preferred sites for egg laying. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Popillia japonica</i> Newman 1841 [Coleoptera: Scarabaeidae]	Not known to occur	No: Scarabaeid larvae feed on plant roots and adults feed on foliage or flowers (Flaherty <i>et al.</i> 1992). <i>Popillia</i> species lay eggs in the soil and emerging larvae feed on the roots (Zhang 2005; EPPO 2006, Tan <i>et al.</i> 1998). Feeding adults skeletonise plant leaves and can cause complete defoliation (Regniere <i>et al.</i> 1983). Therefore, foliage free and root free dormant cuttings do not provide a pathway for <i>Popillia</i> species.	Assessment not required		
<i>Popillia mutans</i> Newman 1838 [Coleoptera: Scarabaeidae]	Not known to occur		Assessment not required		
<i>Popillia quadriguttata</i> Fabricius 1787 [Coleoptera: Scarabaeidae]	Not known to occur		Assessment not required		
<i>Proagopertha lucidula</i> Faldermann 1835 [Coleoptera: Scarabaeidae]	Not known to occur	No: This scarabaeid beetle lays eggs in the soil, larvae feed on roots and adults feed on leaves and flowers (Lee <i>et al.</i> 1973). Therefore, foliage free and root free dormant cuttings do not provide a pathway for this beetle.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Protaetia brevitarsis</i> Lewis 1879 [Coleoptera: Scarabaeidae]	Not known to occur	No: Larvae feed on roots of grapevines while adults feed on buds, leaves, flowers and fruit (Zhang 2005). Therefore, foliage free and root free dormant cuttings do not provide a pathway for this beetle.	Assessment not required		
<i>Rhyparida dimidiata</i> Baly 1861 [Coleoptera: Chrysomelidae]	Yes (Naumann 1993)	Assessment not required			
<i>Rhyparida polymorpha</i> Lea 1915 [Coleoptera: Chrysomelidae]	Yes (AFD 2008)	Assessment not required			
<i>Scelodonta brevipilis</i> Lea 1915 [Coleoptera: Chrysomelidae]	Yes (PHA 2001)	Assessment not required			
<i>Scelodonta lewisii</i> Baly 1874 [Coleoptera: Chrysomelidae]	Not known to occur	No: These chrysomelid beetles damage the sprouting buds and also feed on tender shoots, pedicels, leaves and tendrils (Sun <i>et al.</i> 1992; NHB 2007). Therefore, foliage free dormant cuttings do not provide a pathway for these beetles.	Assessment not required		
<i>Scelodonta strigicollis</i> Motschulsky 1866 [Coleoptera: Chrysomelidae]	Not known to occur		Assessment not required		
<i>Sinoxylon perforans</i> Schrank 1789 [Coleoptera: Bostrichidae]	Not known to occur	<b>Yes:</b> These species lay eggs into branches and emerging larvae tunnel into new shoots (Filip 1986; Moleas 1988). Adults are twig borers and feeds on shoots and branches and have a long life cycle. Therefore, dormant cuttings may harbour larvae and may provide a pathway for these bostrichids.	<b>Yes:</b> These species occur naturally in temperate climates (Filip 1986; Moleas 1988) and would find both climatic conditions and host plants suitable for survival and establishment in Australia. Independent spread is facilitated by active flying (Fettig 2005). Therefore, these species have the	<b>Yes:</b> <i>Sinoxylon perforans</i> is recorded as infesting 30–40% of grapevines in Romania (Filip 1986) and is becoming a serious pest in Italy (Ragazzini 1996). <i>Sinoxylon sexdentatum</i> has been recorded as causing severe infestations (28%) in a two year old vineyard in Italy (Moleas 1988). Therefore, these pests have the potential for economic	<b>Yes</b>
<i>Sinoxylon sexdentatum</i> Olivier 1790 [Coleoptera: Bostrichidae]	Not known to occur				<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
			potential to establish and spread in Australia.	consequences in Australia.	
<i>Sitona discoideus</i> Gyllenhal 1834 [Coleoptera: Curculionidae]	Yes (PHA 2001)	Assessment not required			
<i>Stenygrinum quadrinotatum</i> Bates 1873 [Coleoptera: Cerambycidae]	Not known to occur	No: Larvae of this species attack woody parts of grapevines as internal borers (Luo <i>et al.</i> 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying and development. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Trichoferus campestris</i> Faldermann 1835 [Coleoptera: Cerambycidae]	Not known to occur	No: This species is a timber borer that has been intercepted in dunnage (Iwata and Yamada 1990; Grebennikov <i>et al.</i> 2010). Borers require thick wood for egg laying and development (Goodwin 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying and development. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Tristaria grouvellei</i> Reitter 1878 [Coleoptera: Bostrichidae]	Yes (PHA 2001)	Assessment not required			
<i>Trogoxylon impressum</i> Comolli 1837 [Coleoptera: Lyctidae]	Yes (PHA 2001)	Assessment not required			
<i>Xyleborus cristatus</i> Schedl 1953	Not known to	No: Scolytine beetles are			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Coleoptera: Curculionidae]	occur	associated with woody plant products (Luo <i>et al.</i> 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying and development. Therefore, dormant cuttings do not provide a pathway for this species.			
<i>Xylobosca bispinosa</i> MacLeay 1872 [Coleoptera: Bostrichidae]	Yes (Naumann 1993)	Assessment not required			
<i>Xylopsocus gibbicollis</i> MacLeay 1872 [Coleoptera: Bostrichidae]	Yes (Naumann 1993)	Assessment not required			
<i>Xylotrechus pyrrhoderus</i> Bates 1873 [Coleoptera: Cerambycidae]	Not known to occur	No: Larvae bore into the roots, stems and branches of grapevines (Zhang 2005). Borers require thick wood for egg laying and development (Goodwin 2005). Semi-hardwood dormant cuttings are not preferred sites for egg laying and development. Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<b>DERMAPTERA (earwigs)</b>					
<i>Forficula auricularia</i> Linnaeus 1758 [Dermaptera: Forficulidae]	Yes (Weiss and McDonald 1998)	Assessment not required			
<b>DIPTERA (flies)</b>					
<i>Bactrocera dorsalis</i> (Hendel 1912) [Diptera: Tephritidae]	Not known to occur	No: This species damages fruit (White and Elson-Harris 1992). None of the life stages are associated with dormant			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		cuttings; therefore this pest is not on the pathway.			
<i>Bactrocera tryoni</i> Froggatt 1897 [Diptera: Tephritidae]	Yes (Malipail et al. 1996), under official control.	No: <i>Bactrocera tryoni</i> larvae feed internally in fruit (Botha et al. 2000). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Ceratitis capitata</i> Wiedemann 1824 [Diptera: Tephritidae]	Yes (Smith et al. 1997), under official control	No: <i>Ceratitis capitata</i> adults lay eggs within host fruits and the larvae feed internally (Thomas et al. 2001). Therefore, dormant grapevine cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Ceratitis rosa</i> Karsch 1887 [Diptera: Tephritidae]	Not known to occur	No: This species damages fruits (White and Elson-Harris 1992; Smith et al. 1997). None of the life stages are associated with dormant cuttings; therefore this pest is not on the pathway.	Assessment not required		
<i>Contarinia johnsoni</i> Felt 1909 [Diptera: Cecidomyiidae]	Not known to occur	No: Grape blossom midges lay eggs in unopened grape flower buds and hatching larvae eat the inner portions of the flower (Williams et al. 2011). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Drosophila melanogaster</i> Meigen 1830 [Diptera: Drosophilidae]	Yes (Naumann 1993)	Assessment not required			
<i>Drosophila simulans</i> Sturtevant 1919	Yes (PHA 2001)	Assessment not required			



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Diptera: Drosophilidae]					
<i>Drosophila suzukii</i> matsumura 1931 [Diptera: Drosophilidae]	Not known to occur	No: This species preferentially lays eggs on fully ripened fruits of host plants (Kanzawa 1936). Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Lasioptera vitis</i> Osten Sacken 1862 [Diptera: Cecidomyiidae]	Not known to occur	No: Eggs are laid either on or in leaves, leaf petioles, tendrils or cluster stems, causing gall formation on these plant parts (Williams <i>et al.</i> 2011). The emerging larvae feed within the gall and later on larvae leave the galls, fall to the soil and pupate. Dormant cuttings are not egg laying sites for gall-forming flies and therefore do not provide a pathway for this pest.	Assessment not required		
<b>HEMIPTERA (aphids, leafhoppers, mealybugs, psyllids, scales, true bugs, whiteflies)</b>					
<i>Acia lineatifrons</i> Naude 1926 [Hemiptera]	Not known to occur	No: This species lays eggs on the underside of leaves (Marais 1997) and adults feed on leaves and suck sap from the phloem (Marais 1997). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Aleurolobus taeonabe</i> (Kuwana 1911) [Hemiptera: Aleyrodidae]	Not known to occur	No: Adults and nymphs suck plant juice from the leaves and grape berries (Li 2004).	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		Therefore, foliage free dormant cuttings do not provide a pathway for this species.			
<i>Amblypelta lutescens lutescens</i> Distant 1911 [Hemiptera: Coreidae]	Yes (Naumann 1993)	Assessment not required			
<i>Aonidiella aurantii</i> Maskell 1879 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Aonidiella orientalis</i> Newstead 1894 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Aphis craccivora</i> Koch 1854 [Hemiptera: Aphididae]	Yes (Naumann 1993)	Assessment not required			
<i>Aphis fabae</i> Scopoli 1763 [Hemiptera: Aphididae]	Not known to occur	No: These aphid species overwinter as eggs on their primary hosts (Cammell 1981; Mackenzie and Dixon 1990; OHU 2010) and adults move to secondary hosts, in the summer months and attack the foliage, flowers and twigs of host plants (Mackenzie 1996; Liburd <i>et al.</i> 2004; Graham 2007). Therefore, foliage free dormant cuttings do not provide a pathway for these aphids.	Assessment not required		
<i>Aphis illinoisensis</i> Shimer 1866 [Hemiptera: Aphididae]	Not known to occur		Assessment not required		
<i>Aphis spiraeicola</i> Patch 1914 [Hemiptera: Aphididae]	Yes (Naumann 1993)	Assessment not required			
<i>Arboridia adanae</i> Diabola 1957 [Hemiptera: Cicadellidae]	Not known to occur	No: These leafhoppers feed on the leaf-mesophyll tissue of <i>Vitis</i> species (Bournier 1976). <i>Arboridia adanae</i> eggs are laid on the leaves	Assessment not required		
<i>Arboridia apicalis</i> Nawa 1913 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Arboridia hussaini</i> Ghauri 1963	Not known to		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Hemiptera: Cicadellidae]	occur	(Kharizanov 1969). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.			
<i>Arboridia Kermanshah</i> Dlabola 1963 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Arboridia viniferata</i> Sohi & Sandhu 1971 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Aspidiotus destructor</i> Signoret 1869 [Hemiptera: Diaspididae]	Yes (PHA 2001)	Assessment not required			
<i>Aspidiotus nerii</i> Bouché 1966 [Hemiptera: Diaspididae]	Yes (PHA 2001)	Assessment not required			
<i>Asterolecanium pustulans</i> Cockerell 1892 [Hemiptera: Asterolecaniidae] (synonym: <i>Russellaspis pustulans pustulans</i> Cockerell)	Not known to occur	No: Oleander pit scale is found on the leaves, bark, stems and fruit of host plants (Hamon 1977), including grapevines (Ben-Dov <i>et al.</i> 2012). The severity of pit development around the scale is dependent on the susceptibility of host plants (Hamon 1977). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Balclutha hebe</i> Kirkaldy 1976 [Hemiptera: Cicadellidae]	Not known to occur	No: This species feeds and lays eggs on the leaves of host plants (Abu-Yaman 1967). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Ceroplastes rusci</i> Linnaeus 1758 [Hemiptera: Coccidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Chrysomphalus aonidum</i> Linnaeus 1758 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Chrysomphalus dictyospermi</i> Morgan 1889 [Hemiptera: Diaspididae]	Yes ( Naumann 1993)	Assessment not required			
<i>Cicadella viridis</i> Linnaeus 1758 [Hemiptera: Cicadellidae]	Not known to occur	No: This species often occurs in fens and marshes (Nickel and Remane 2002) and in vineyards (Mazzoni <i>et al.</i> 2001). This leafhopper species feeds on the leaves of host plants (Silverside 2006). Therefore, foliage free dormant cuttings do not provide a pathway for this leafhopper.	Assessment not required		
<i>Coccus hesperidum</i> Linnaeus 1758 [Hemiptera: Coccidae]	Yes (Naumann 1993)	Assessment not required			
<i>Colgar peracutum</i> Walker 1858 [Hemiptera: Flatidae]	Yes (Naumann 1993)	Assessment not required			
<i>Creontiades dilutus</i> Stal 1859 [Hemiptera: Miridae]	Yes (Naumann 1993)	Assessment not required			
<i>Daktulosphaira vitifoliae</i> Fitch 1855 [Hemiptera: Phylloxeridae]	Yes (Restricted distribution [Loch and Slack 2007] and under official control [NVHSC 2005]).	<b>Yes:</b> <i>Daktulosphaira vitifoliae</i> has a complex lifecycle that involves migration from the roots to the leaves and back to the roots (Granett <i>et al.</i> 2001; Forneck and Huber 2009). It is possible that migrating crawlers or overwintering hibernants could be present on grapevine stems. Radicoles may also feed on stems (Granett <i>et al.</i> 2001) and winged forms lay overwintering eggs beneath the bark on grapevine stems	<b>Yes:</b> <i>Daktulosphaira vitifoliae</i> is already established in small areas of Victoria and New South Wales in Australia (Loch and Slack 2007), indicating its potential to establish and spread in similar environments in other parts of Australia. It can spread naturally by active crawling of from vine to vine, by wind and and through human assisted means (NVHSC 2005). Therefore, this pest has the potential for establishment	<b>Yes:</b> Feeding activity of this insect reduces productivity of vineyards (Granett <i>et al.</i> 2001; Lock and Slack 2007) and infestation render vineyards uneconomic within 3–10 years (Buchanan and Whiting 1991). Therefore, it has the potential for economic consequences in Australia. This insect is considered of quarantine concern by several countries. Presence and spread of this insect in other grapevine growing areas of Australia would impact	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Buchanan <i>et al.</i> 1994; Granett <i>et al.</i> 2001). Therefore, this pest may be associated with grapevine dormant cuttings.	and spread in Australia.	upon Australia's ability to access overseas markets. Therefore, this insect has the potential for economic consequences in Australia.	
<i>Diaspidiotus ancyclus</i> Putnam 1878 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Diaspidiotus perniciosus</i> (Comstock) Cockerell 1899 [Hemiptera: Diaspididae]	Yes (PHA 2001)	Assessment not required			
<i>Diaspidiotus uvae</i> Comstock 1881 [Hemiptera: Diaspididae]	Not known to occur	No: Grape scale is associated with two year old wood and spends most of its life under the protection of its waxy scale cover (Williams <i>et al.</i> 2011). Therefore, one year old dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Dolycoris baccarum</i> (Linnaeus 1758) [Hemiptera: Pentatomidae]	Not known to occur	No: Nymphs and adults suck sap from young buds, leaves, young shoots and fruit of grapevines (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this insect.	Assessment not required		
<i>Draeculacephala minerva</i> Ball 1927 [Hemiptera: Cicadellidae]	Not known to occur	No: Sharpshooters are xylophagous and feed on leaves, buds, shoots and stems of grapevine (Hill and Purcell 1997; Feil <i>et al.</i> 2000; Irvin and Hoddle 2005; Flint 2006). Egg masses are laid under the lower leaf epidermis	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		of host plants (Bentley <i>et al.</i> 2008). Adults are mobile and are highly unlikely to remain on shoots and stems following harvest, while egg masses are not associated with shoots or stems. Therefore, foliage free dormant cuttings do not provide a pathway for this insect.			
<i>Dysdercus sidae</i> Montrouzier 1861 [Hemiptera: Pentatomidae]	Yes (Naumann 1993)	Assessment not required			
<i>Empoasca decipiens</i> Paoli 1930 [Hemiptera: Cicadellidae]	Not known to occur	No: <i>Empoasca</i> leafhoppers lay eggs on leaves and adults feed on leaves (Boll and Herrmann 2001; Backus <i>et al.</i> 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Empoasca fabae</i> Harris 1841 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Empoasca punjabensis</i> Singh-Pruthi 1940 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Empoasca vitis</i> Gothe 1875 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura bistrata</i> McAtee 1920 [Hemiptera: Cicadellidae]	Not known to occur	No: <i>Erythroneura</i> leafhoppers lay eggs on foliage (MacGill 1932; Paxton and Thorvilson 1996) and feed primarily on the leaf (Martinson <i>et al.</i> 1997; Flint 2006). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Erythroneura calycula</i> McAtee 1920 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura coloradensis</i> Gillette 1892 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura comes</i> Say 1825 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura elegantula</i> Osborn 1828 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura pallidifrons</i> Edwards 1924 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura tricincta</i> Fitch 1851	Not known to		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Hemiptera: Cicadellidae]	occur				
<i>Erythroneura variabilis</i> Beamer 1929 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura vitifex</i> Fitch 1856 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura vitis</i> Harris 1831 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura vulnerata</i> Fitch 1851 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erythroneura ziczac</i> Walsh 1862 [Hemiptera: Cicadellidae]	Not known to occur		Assessment not required		
<i>Erthesina fullo</i> (Thunberg 1783) [Hemiptera: Pentatomidae]	Not known to occur	No: Nymphs and adults feed on young buds, leaves and young shoots of grapevines (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this insect.	Assessment not required		
<i>Eulecanium pruinsum</i> Coquillett [Hemiptera: Coccidae]	Yes (Malipatil <i>et al.</i> 1996)	Assessment not required			
<i>Eulecanium tiliae</i> (Linnaeus 1758) [Hemiptera: Coccidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Euschistus conspersus</i> Uhler 1979 [Hemiptera: Pentatomidae]	Not known to occur	No: Pentatomine stink bugs are mostly associated with fruits but also feed on stems and leaves (Weaver 1976; McPherson and McPherson 2000). This pest lays eggs on groundcover crops and occasionally on the leaves of host fruit trees (Borden and Madsen 1951). Therefore, foliage free dormant cuttings	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		do not provide a pathway for this stink bug.			
<i>Ferrisia virgata</i> Cockerell 1893 [Hemiptera: Pseudococcidae]	Yes (Naumann 1993)	Assessment not required			
<i>Graphocephala atropunctata</i> Signoret 1854 [Hemiptera: Cicadellidae]	Not known to occur	No: Sharpshooters are xylophagous and feed on leaves, buds, shoots and stems of grapevines (Hill and Purcell 1997; Feil <i>et al.</i> 2000; Irvin and Hoddle 2005; Flint 2006). Eggs are laid under the lower leaf epidermis (CABI 2012a). Adults are mobile and are highly unlikely to remain on shoots and stems following harvest, while egg masses are not associated with shoots or stems. Therefore, foliage free dormant cuttings do not provide a pathway for this insect.	Assessment not required		
<i>Halyomorpha halys</i> (Stål 1855) [Hemiptera: Pentatomidae]	Not known to occur	No: Nymphs and adults suck sap from young buds, leaves, young shoots and fruit of grapevines (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this insect.	Assessment not required		
<i>Heliococcus bohemicus</i> Sulc 1912 [Hemiptera: Pseudococcidae]	Not known to occur	No: This species feeds on the leaves of herbaceous plants and on the bark of woody plants (Ben-Dov <i>et al.</i> 2012). This insect has been recorded	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		on the foliage of grapes (Zorloni <i>et al.</i> 2006). Therefore, foliage free dormant cuttings do not provide a pathway for this mealybug.			
<i>Helopeltis antonii</i> Signoret 1858 [Hemiptera: Miridae]	Not known to occur	No: This insect sucks the sap of young plants and the tender new growth of host plants (Siswanto <i>et al.</i> 2008). Tender shoots, leaves, petioles and immature fruits of new growth flushes are the sites of egg laying as well as feeding (Sundararaju and Babu 2000; Siswanto <i>et al.</i> 2008). Dormant cuttings are not the preferred site for egg laying and therefore do not provide a pathway for this insect.	Assessment not required		
<i>Hemiberlesia lataniae</i> Signoret 1869 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Hemiberlesia rapax</i> Comstock 1881 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Homalodisca coagulata</i> Say 1832 [Hemiptera: Cicadellidae]	Not known to occur	No: Sharpshooters are xylophagous and feed on leaves, buds, shoots and stems of grapevine (Hill and Purcell 1997; Feil <i>et al.</i> 2000; Irvin and Hoddle 2005; Flint 2006). Eggs are laid under the lower leaf epidermis (CABI 2012a). Therefore, foliage	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		free dormant cuttings do not provide a pathway for this insect.			
<i>Hyalesthes obsoletus</i> Signoret 1865 [Hemiptera: Cixiidae]	Not known to occur	No: The eggs and larvae of this species are associated with roots while adults feed on foliage (Riolo <i>et al.</i> 2007; Forte <i>et al.</i> 2010). Therefore, root and foliage free dormant cuttings do not provide a pathway for this planthopper.	Assessment not required		
<i>Icerya palmeri</i> Riley & Howard 1890 [Hemiptera: Margarodidae]	Not known to occur	No: This species is associated with <i>Vitis</i> species (Morales 1991). Other members of this genus lay eggs within an egg sac and crawlers move to and settle on the underside of the leaves. The older nymphs continue to feed but migrate to the larger twigs, and finally, as adults, settle on the larger branches and trunk (Fasulo and Brooks 2010). Therefore, foliage free, semi-hardwood dormant cuttings do not provide a pathway for this scale.	Assessment not required		
<i>Icerya purchasi</i> Maskell 1878 [Hemiptera: Margarodidae]	Yes (Naumann 1993)	Assessment not required			
<i>Icerya seychellarum</i> Westwood 1855 [Hemiptera: Margarodidae]	Yes (Smith <i>et al.</i> 1997)	Assessment not required			
<i>Jacobiasca lybica</i> Bergevin & Zanon 1922 [Hemiptera: Cicadellidae]	Not known to occur	No: The green leaf hopper lays eggs on the underside of leaves and adults are foliage	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		feeders (Gonzalez-Andujar <i>et al.</i> 2006). Therefore, foliage free dormant cuttings do not provide a pathway for this leaf hopper.			
<i>Lepidosaphes ulmi</i> Linnaeus 1758 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Lygus lineolaris</i> (Palisot de Beauvois 1818) [Hemiptera: Miridae]	Not known to occur	No: The nymphs and adults feed on leaves and flowers of grapevines (Bostanian <i>et al.</i> 2003; Fleury <i>et al.</i> 2006) from early spring until grape harvest (Bostanian <i>et al.</i> 2003). This bug overwinters in fallen plant litter, including dead leaves (Cleveland 1982; Wheeler and Stimmel 1988). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Lygus lucorum</i> Meyer-Duer 1843 [Hemiptera: Miridae]	Not known to occur	No: This species lays eggs on the tips of vegetative branches of host plants (Guo <i>et al.</i> 2005). Both nymphs and the adults damage young shoots and leaves causing withering and perforation in grapes (Lee <i>et al.</i> 2002; Liu <i>et al.</i> 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Maconellicoccus hirsutus</i> Green 1908	Yes (Gullan 2000)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Hemiptera: Pseudococcidae]					
<i>Macrosiphum euphorbiae</i> Thomas 1778 [Hemiptera: Aphididae]	Yes (Naumann 1993)	Assessment not required			
<i>Magicicada septendecim</i> (Linnaeus 1758) [Hemiptera: Cicadidae]	Not known to occur	No: Adult female cicadas injure grapevines by making ovipositional slits in the young canes. The canes then may break at the slits (Williams <i>et al.</i> 2011). Dormant cuttings are not preferred sites for egg laying and therefore do not provide a pathway for these pests.	Assessment not required		
<i>Magicicada cassinii</i> (Fisher 1851) [Hemiptera: Cicadidae]	Not known to occur		Assessment not required		
<i>Magicicada septendecula</i> Alexander & Moore 1962 [Hemiptera: Cicadidae]	Not known to occur		Assessment not required		
<i>Margarodes brasiliensis</i> Wille 1922 [Hemiptera: Margarodidae]	Not known to occur	No: Adult ground pearls lay eggs in the vicinity of roots and hatching larvae feed on root tissues (de Klerk 2010). Pupation also occurs in the soil (de Klerk 1987; de Klerk 2010). Therefore, root free dormant cuttings do not provide a pathway for these ground pearls.	Assessment not required		
<i>Margarodes capensis</i> Giard 1897 [Hemiptera: Margarodidae]	Not known to occur		Assessment not required		
<i>Margarodes greeni</i> Brain 1915 [Hemiptera: Margarodidae]	Not known to occur		Assessment not required		
<i>Margarodes meridionalis</i> Morrison 1927 [Hemiptera: Margarodidae]	Not known to occur		Assessment not required		
<i>Margarodes prieskaensis</i> (Jakubski 1965) [Hemiptera: Margarodidae]	Not known to occur		Assessment not required		
<i>Margarodes vitis</i> (Philippi 1884) [Hemiptera: Margarodidae]	Not known to occur		Assessment not required		
<i>Margarodes vredendalensis</i> De Klerk 1980 [Hemiptera: Margarodidae]	Not known to occur		Assessment not required		
<i>Metcalfa pruinosa</i> Say 1830 [Hemiptera: Flatidae]	Not known to occur	No: Frosted moth-bugs lay eggs and overwinter in the corky parts of the bark or under the bark of host plants (Lucchi and Santini 1993; Kahrer 2005). Therefore,	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		semi-hardwood dormant cuttings do not provide a pathway for this pest.			
<i>Mictis profana</i> Fabricius 1803 [Hemiptera: Coreidae]	Yes (PHA 2001)	Assessment not required			
<i>Myzus persicae</i> Sulzer 1776 [Hemiptera: Aphididae]	Yes (Naumann 1993)	Assessment not required			
<i>Nezara viridula</i> Linnaeus 1758 [Hemiptera: Aphididae]	Yes (Smith <i>et al.</i> 1997)	Assessment not required			
<i>Nipaecoccus viridis</i> Newstead 1894 [Hemiptera: Pseudococcidae]	Yes (Gullan 2000)	Assessment not required			
<i>Nysius ericae</i> (Schilling 1829) [Hemiptera: Lygaeidae]	Not known to occur	No: Nymphs and adults of <i>Nysius</i> species attack the leaves of host plants (Malipatil <i>et al.</i> 1996; Flint 2006). Adults suck sap from the leaves and fruits of host plants (Malipatil <i>et al.</i> 1996; Flint 2006). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Nysius niger</i> Baker 1906 [Hemiptera: Lygaeidae]	Not known to occur		Assessment not required		
<i>Nysius raphanus</i> Howard 1872 [Hemiptera: Lygaeidae]	Not known to occur		Assessment not required		
<i>Nysius vinitor</i> Bergroth 1891 [Hemiptera: Lygaeidae]	Yes (Naumann 1993)	Assessment not required			
<i>Oxycarenus arctatus</i> Walker 1872 [Hemiptera: Oxycarenidae]	Yes (Naumann 1993)	Assessment not required			
<i>Parasaissetia nigra</i> Nietner 1861 [Hemiptera: Coccidae]	Yes (PHA 2001)	Assessment not required			
<i>Parlatoria oleae</i> Colvée 1880 [Hemiptera: Diaspididae]	Yes (Naumann 1993)	Assessment not required			
<i>Parthenolecanium corni</i> Bouché 1844 [Hemiptera: Coccidae]	Yes (Naumann 1993)	Assessment not required			
<i>Parthenolecanium pruinosum</i> Coquillett	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
1891 [Hemiptera: Coccidae]					
<i>Perissopneumon ferox</i> Newstead 1900 [Hemiptera: Margarodidae]	Not known to occur	No: This species lays eggs in the soil (Srivastava and Verghese 1985) and adults feed on fruit stalks, inflorescences and fruit (Srivastava 1997). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Pinnaspis strachani</i> Cooley 1899 [Hemiptera: Diaspididae]	Yes (PHA 2001)	Assessment not required			
<i>Planococcus citri</i> Risso 1813 [Hemiptera: Pseudococcidae]	Yes (Gullan 2000)	Assessment not required			
<i>Planococcus ficus</i> Signoret 1875 [Hemiptera: Pseudococcidae]	Not known to occur	<b>Yes:</b> The vine mealybug is capable of feeding on many different parts of grapevines, including trunks, canes, leaves, clusters and sometimes the roots (Bournier 1976; Fuchs 2007; Bentley <i>et al.</i> 2008). Therefore, dormant cuttings may provide a pathway for vine mealybugs.	<b>Yes:</b> <i>Planococcus ficus</i> is polyphagous and has established in areas with a wide range of climatic conditions (Walton and Pringle 2004) and can spread naturally in infested propagative material. (Haviland <i>et al.</i> 2005). Therefore, this species has the potential for establishment and spread in Australia.	<b>Yes:</b> This species is reported as one of the most important pests of grape industries in South Africa (Walton <i>et al.</i> 2004). It causes progressive weakening of vines through early leaf loss, yield loss and reduced crop quality (Walton and Pringle 2004; Walton <i>et al.</i> 2004). Therefore, vine mealybugs have the potential for economic consequences in Australia.	<b>Yes</b>
<i>Planococcus lilacinus</i> Cockerell 1905 [Hemiptera: Pseudococcidae]	Not known to occur	<b>Yes:</b> Mealybugs may be concealed under the bark or may be spread over different parts of the host plant (Flint 2006). This mealybug has been intercepted on host cuttings (MacLeod 2006). Therefore, dormant cuttings	<b>Yes:</b> Coffee mealybug is polyphagous (Ben-Dov 1994) and has established in areas with a wide range of climatic conditions (Williams 1982; Ben-Dov 1994). It can spread naturally in infested propagative material (Williams	<b>Yes:</b> This species causes damage to a wide variety of economically important crops. It is considered a potential threat to citrus, grapes, guavas and mangoes (Tandon and Verghese 1987; Cox 1989). This species causes severe damage to young	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		may provide a pathway for this mealybug.	1982) as it has been intercepted on host cuttings (MacLeod 2006). Therefore, coffee mealybug has the potential for establishment and spread in Australia.	trees by killing the tips of branches and roots of many economically important species (Tandon and Verghese 1987). Therefore, it has the potential for economic consequences in Australia.	
<i>Planococcus kraunhiae</i> (Kuwana 1902) [Hemiptera: Pseudococcidae]	Not known to occur	<b>Yes:</b> This mealybug is reported on grapes (Narai and Murai 2002) and is found on leaves and branches of grapes (NPQS 2007). Therefore, dormant cuttings may provide a pathway for this mealybug.	<b>Yes:</b> This mealybug is polyphagous (Ben-Dov 1994) and has established in areas with a wide range of climatic conditions (Ben-Dov 1994). It can spread naturally in infested propagative material. Therefore, this mealybug has the potential for establishment and spread in Australia.	<b>Yes:</b> This sap sucking insect reduces productivity and quality and promotes the growth of sooty mould through production of honeydew (CABI 2012a). Although the mouth parts of mealybugs rarely penetrate beyond the fruit epidermis, their feeding activities can also cause fruit spotting and distortion (CABI 2012a). Therefore, it has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Plautia affinis</i> Dallas 1851 [Hemiptera: Pentatomidae]	Yes (Coombs and Khan 1998)	Assessment not required			
<i>Plautia stali</i> Scott 1874 [Hemiptera: Pentatomidae]	Not known to occur	No: Adults feed on fruit when ripe or near ripe (Mau and Mitchell 1978; Schaefer and Panizzi 2000). Therefore, fruit free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Pseudococcus calceolariae</i> Maskell 1879 [Hemiptera: Pseudococcidae]	Yes (Gullan 2000)	Assessment not required			
<i>Pseudococcus longispinus</i> Targioni-Tozzetti 1867 [Hemiptera: Pseudococcidae]	Yes (Gullan 2000)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Pseudococcus maritimus</i> Ehrhorn 1900 [Hemiptera: Pseudococcidae]	Not known to occur <sup>7</sup>	<b>Yes:</b> <i>Pseudococcus maritimus</i> overwinters under bark as eggs or first-instar crawlers (Burts and Dunley 1993; Grasswitz and James 2008). Adults and nymphs feed on the bases of buds, stalks, young branches, vines and young roots (Burts and Dunley 1993; AQSIQ 2007). Therefore, dormant grapevine cuttings may provide a pathway for this pest.	<b>Yes:</b> This mealybug is polyphagous and has established in areas with a wide range of climatic conditions (Ben-Dov 1994). <i>Pseudococcus maritimus</i> can spread naturally in infested propagative material and by the movement of crawlers and winged males (Burts and Dunley 1993; Grasswitz and James 2008). Therefore, this mealybug has the potential for establishment and spread in Australia.	<b>Yes:</b> This sap sucking insect reduces productivity and quality and promotes the growth of sooty mould through the production of honeydew (Geiger and Daane 2001). The damage caused by sooty mould growth is purely cosmetic but lowers grape quality and marketability. <i>Pseudococcus maritimus</i> is also a vector of Grapevine leafroll-associated virus 3 (Grasswitz and James 2008). Therefore, this mealybug is of economic significance to Australia.	<b>Yes</b>
<i>Pseudococcus viburni</i> Signoret 1875 [Hemiptera: Pseudococcidae] (synonym: <i>Pseudococcus affinis</i> Maskell 1894)	Yes (Gullan 2000)	Assessment not required			
<i>Pulvinaria innumerabilis</i> Putnam 1880 [Hemiptera: Coccidae]	Not known to occur	<b>Yes:</b> <i>Pulvinaria</i> species overwinter as immature females attached to the twigs and small branches of host plants (University of Illinois 2004). Therefore, dormant cuttings provide a pathway for these scales.	<b>Yes:</b> These scales have established in areas with a wide range of climatic conditions (Ben-Dov <i>et al.</i> 2012) and can spread naturally in infested propagative material. Establishment will be favoured by the wide host range in Australia. Therefore, these scales have the potential to establish and spread in Australia.	No: <i>Pulvinaria</i> species damage shoots and foliage by sucking sap (Bournier 1976; Fuchs 2007). Although <i>Pulvinaria innumerabilis</i> and <i>P. vitis</i> are vectors of Grapevine leafroll-associated virus 1 and 3 (Fuchs <i>et al.</i> 2007), these viruses are already present in Australia. Therefore, these scales are not of economic significance to Australia.	
<i>Pulvinaria vitis</i> Linnaeus 1758 [Hemiptera: Coccidae]	Not known to occur				

<sup>7</sup> Reports of this species in Australia are based on misidentifications of *P. affinis*, *P. caleolariae* and *P. longispinus* (Williams 1985).



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Quadraspidiotus perniciosus</i> Comstock 1881 [Hemiptera: Diaspididae]	Yes (PHA 2001)	Assessment not required			
<i>Rastrococcus iceryoides</i> Green 1908 [Hemiptera: Pseudococcidae]	Not known to occur	No: This species is reported to occur on grapevine (Ben-Dov <i>et al.</i> 2012). Adults usually feed on the tender terminal shoots, inflorescences and fruit whereas first instars nymphs feed on the underside of leaves (Rawat and Jakhmola 1970). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Rhizoecus falcifer</i> Kunchel d'Herculis 1878 [Hemiptera: Pseudococcidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Rhizoecus kondonis</i> Kuwana 1923 [Hemiptera: Pseudococcidae]	Not known to occur	No: The citrus ground mealybug exists entirely below the soil surface and sucks the liquid from small feeder roots (Blodgett 1992). Therefore, root free dormant cuttings do not provide a pathway for citrus ground mealybugs.	Assessment not required		
<i>Saissetia coffeae</i> Walker 1852 [Hemiptera: Coccidae]	Yes (Naumann 1993)	Assessment not required			
<i>Saissetia oleae</i> Olivier 1791 [Hemiptera: Coccidae]	Yes (Naumann 1993)	Assessment not required			
<i>Scaphoideus titanus</i> Ball 1931 [Hemiptera: Cicadellidae]	Not known to occur	No: Juveniles and adults feed on shoots near the root stock and leaves. Females lay eggs beneath the bark of two year	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		old wood where they overwinter (Lessio and Alma 2004a, b; Boudon-Padieu and Maixner 2007). Therefore, one year old dormant cuttings do not provide a pathway for vine leafhoppers.			
<i>Scolypopa australis</i> Walker 1851 [Hemiptera: Ricaniidae]	Yes (Smith <i>et al.</i> 1997)	Assessment not required			
<i>Scutiphora pedicellata</i> Kirby 1826 [Hemiptera: Scutelleridae]	Yes (PHA 2001)	Assessment not required			
<i>Spissistilus festinus</i> Say 1830 [Hemiptera: Membracidae]	Not known to occur	No: This hopper is a pest of soybean (Rice and Drees 1985) and feeds on leaves and stems and lays eggs in the stems of soybean (Hudson and Adams 2008). Dormant grapevines are not preferred sites for egg laying. Therefore, this species is not on the pathway of grapevine propagative material.	Assessment not required		
<i>Targionia vitis</i> Signoret 1876 [Hemiptera: Diaspididae]	Not known to occur	<b>Yes:</b> The black vine scale feeds on stems and branches, especially under bark flakes (Stathas and Kontodimas 2001; Watson 2005). Therefore, dormant cuttings may harbour mated females and provide a pathway for black vine scale.	<b>Yes:</b> The black vine scale has a wide host range (Watson 2005), has established in areas with a wide range of climatic conditions (Ben-Dov <i>et al.</i> 2012) and can spread naturally in infested propagative material. Therefore, this scale has the potential to establish and spread in Australia.	<b>Yes:</b> This species is reported as one of the most important pests of table grapes in Italy (Guario and Laccone 1996). Heavy infestations of black vine scale may encrust the bark with several layers of scale covers (Watson 2005) and may cause defoliation, splitting of bark, twig dieback and an overall decline in host plant health (Beardsley and Gonzalez 1975). Therefore, this	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
				scale has the potential for economic consequences in Australia.	
<i>Tettigades chilensis</i> Amyot & Serville 1843 [Hemiptera: Cicadidae]	Not known to occur	No: This plant hopper feeds on the branches and roots of host plants (Gonzalez 1983). Adults are mobile and are unlikely to remain on grapevine cuttings following harvesting. The young cicada nymphs live underground and feed on the roots of trees. Therefore, dormant cuttings do not provide a pathway for this leafhopper.	Assessment not required		
<i>Trialeurodes vaporariorum</i> 1856 Westwood [Hemiptera: Aleyrodidae]	Yes (PHA 2001)	Assessment not required			
<i>Zygina rhamni</i> Ferrari 1882 [Hemiptera: Cicadellidae]	Not known to occur	No: Adults of Italian grape leafhopper overwinter in the shelter of evergreens. In late spring they migrate to summer host plants, including grapes (Alford 2007). Eggs have been recorded on <i>Vitis vinifera</i> in summer (Mazzoni <i>et al.</i> 2008). Larvae feed on leaves (Bournier 1976). Therefore, foliage free dormant cuttings do not provide a pathway for this leafhopper.	Assessment not required		
<b>HYMENOPTERA (wasps, ants)</b>					
<i>Ametastegia glabrata</i> Fallén 1808	Yes (CSIRO)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Hymenoptera: Tenthredinidae]	2005)				
<i>Ceratina dentipes</i> Friese 1914 [Hymenoptera: Apidae]	Not known to occur	No: These wasps lay eggs on the stem and after hatching larvae bore into the stem and feed on woody parts of the grapevine (Luo <i>et al.</i> 2005). Grape dormant cuttings are not preferred sites to lay eggs. Therefore, this species is not on the pathway.	Assessment not required		
<i>Ceratina viticola</i> Sinich [Hymenoptera: Apidae]	Not known to occur		Assessment not required		
<i>Erythraspdes vitis</i> (Harris) [Hymenoptera: Tenthredinidae]	Not known to occur	No: This species lays eggs on the underside of grape leaves and larvae feed at the edge of the leaf (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Evoxysoma vitis</i> (Saunders 1869) [Hymenoptera: Vespidae]	Not known to occur	<b>Yes:</b> This species lays eggs in grape berries and hatching larvae feed on the seeds and overwinter within grape seed on the ground (Williams <i>et al.</i> 2011). Dormant cuttings do not provide a pathway for this species. However, grape seeds may provide a pathway for this species.	<b>Yes:</b> This chalcid has established in areas with a wide range of climatic conditions (Webb 2003) and distribution of infested seed will facilitate the spread of this species. Therefore, this chalcid has the potential for establishment and spread in Australia.	No: Outbreaks of this species are rare and are generally confined to wild grapes (Williams <i>et al.</i> 2011). This chalcid is not reported to cause significant economic consequences. Therefore, this species is unlikely to be of economic consequence in Australia.	
<i>Iridomyrmex humilis</i> Mayr 1868 [Hymenoptera: Formicidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Solenopsis xyloni</i> McCook 1879 [Hymenoptera: Formicidae]	Yes (PHA 2001)	Assessment not required			
<i>Vespula germanica</i> Fabricius 1793 [Hymenoptera: Vespidae]	Yes (Naumann 1993)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<b>ISOPTERA</b>					
<i>Coptotermes acinaciformis</i> Froggatt 1898 [Isoptera: Rhinotermitidae]	Yes (AFD 2008)	Assessment not required			
<i>Incisitermes minor</i> Hagen 1858 [Isoptera: Kalotermitidae]	Not known to occur	No: Colonies of these termites are often found in dead downed logs, and large, dead branches on the ground (Cabrera and Scheffrahn 2005). Therefore, dormant cuttings do not provide a pathway for western dry wood termites.	Assessment not required		
<i>Neotermes chilensis</i> Blanchard [Isoptera: Kalotermitidae]	Not known to occur	No: Damp wood termites feed on the heartwood (dead tissue) of vines and usually avoid the living sapwood (Rust 1992). Therefore, semi-hardwood dormant cuttings do not provide a pathway for damp wood termite.	Assessment not required		
<i>Paraneotermes simplicicornis</i> Banks & Snyder 1920 [Isoptera: Kalotermitidae]	Not known to occur	No: Desert damp wood termites may girdle young grapevines below the soil line in desert areas (Ebeling 2002). Therefore, root free dormant cuttings do not provide a pathway for desert damp wood termite.	Assessment not required		
<b>LEPIDOPTERA (moths, butterflies)</b>					
<i>Abagrotis barnesi</i> (Benjamin 1921) [Lepidoptera: Noctuidae]	Not known to occur	No: Cutworms conceal themselves underneath loose bark or beneath the grape trellis during the day and	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		crawl up the trunk to feed on swelling buds at night (Williams <i>et al.</i> 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this pest.			
<i>Accuminulia buscki</i> Brown 2000 [Lepidoptera: Tortricidae]	Not known to occur	No: The larvae of these species bore into grape berries (Brown 1999). Therefore, fruit free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Accuminulia longiphallus</i> Brown 2000 [Lepidoptera: Tortricidae]	Not known to occur		Assessment not required		
<i>Acosmeryx castanea</i> Rothschild & Jordan 1903 [Lepidoptera: Sphingidae]	Not known to occur	No: These species have been reported from grapevines (Pittaway and Kitching 2006). The larvae of sphingids generally feed on foliage (Common 1990; USDA 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Acosmeryx naga</i> (Moore 1858) [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		
<i>Acosmeryx sericeus</i> (Walker 1856) [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		
<i>Acosmeryx shervillii</i> Boisduval 1875 [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		
<i>Acronicta rumicis</i> (Linnaeus 1948) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of this noctuid moth feed on the foliage of host plants (Thompson and Nelson 2003). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Actias ningpoana</i> Felder 1862 [Lepidoptera: Saturniidae]	Not known to occur	No: Larvae of this species feed on grapevine foliage (Zhang 2005). Therefore, foliage free dormant cuttings	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		do not provide a pathway for this species.			
<i>Agrotis ipsilon</i> Hufnagel 1766 [Lepidoptera: Noctuidae]	Yes (Common 1990)	Assessment not required			
<i>Agrotis munda</i> Walker 1856 [Lepidoptera: Noctuidae]	Yes (Naumann 1993)	Assessment not required			
<i>Agrotis segetum</i> Denis & Schiffermüller 1775 [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of these noctuid moths feed on grape buds and young stems during spring (AgroAtlas 2009a; Wright <i>et al.</i> 2010). These cutworms occur in the soil and litter during the day and climb grapevines to feed on swelling buds at night (Wright <i>et al.</i> 2010). Therefore, dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Agrotis vetusta</i> (Walker 1856) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Alypia octomaculata</i> (Fabricius 1775) [Lepidoptera: Noctuidae]	Not known to occur	No: Eggs are laid on grape shoots and leaves and larvae feed on foliage (Williams <i>et al.</i> 2011). This species overwinters as pupae in tunnels built in old wood or trash just beneath the soil surface (Arnold 1982; Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Ampelophaga khasiana</i> Rothschild 1895 [Lepidoptera: Sphingidae]	Not known to occur	No: These species have been reported from grapevines (Pittaway and Kitching 2006).	Assessment not required		
<i>Ampelophaga rubiginosa</i> Bremer & Grey	Not known to		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
1853 [Lepidoptera: Sphingidae]	occur	The larvae of sphingids generally feed on foliage (Common 1990; Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these species.			
<i>Amphipyra pyramidoides</i> Guenée 1852 [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of pyramidal fruit worm feed on new foliage (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Amyelois transitella</i> Walker 1863 [Lepidoptera: Pyralidae]	Not known to occur	No: This species is associated with postharvest fruit and dried grape fruits (Johnson 2007). Eggs are laid on dried, fallen fruit (Siegel <i>et al.</i> 2006). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Antispila viticordifoliella</i> Clemens 1860 (Lepidoptera: Heliozelidae).	Not known to occur	No: Leaf miner larvae feed between the upper and lower surfaces of leaves (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Aporia crataegi</i> (Linnaeus 1758) [Lepidoptera: Pieridae]	Not known to occur	No: Larvae of this species feed on foliage of many fruiting plants including grapes (Robinson <i>et al.</i> 2008, Grichanov and Ovsyannikova	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.			
<i>Argyrotaenia citrana</i> Fernald 1889 [Lepidoptera: Tortricidae]	Not known to occur	No: These species lay eggs on the leaves and shoots (Zalom <i>et al.</i> 2008, EPPO 2002b) or newly set grape clusters during spring (Williams <i>et al.</i> 2011). Larvae of these tortricid moths feed during spring on buds and leaves and later feed on berries (Bentley <i>et al.</i> 2008). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Argyrotaenia ljugiana</i> Thunberg 1797 [Lepidoptera: Tortricidae]	Not known to occur		Assessment not required		
<i>Argyrotaenia velutinana</i> (Walker 1863) [Lepidoptera: Tortricidae]	Not known to occur		Assessment not required		
<i>Artena dotata</i> (Fabricius 1794) [Lepidoptera: Noctuidae]	Not known to occur	No: This fruit piercing moth feeds on the grapevine fruit (Li 2004). Therefore, fruit free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Autographa gamma</i> (Linnaeus 1758) [Lepidoptera: Noctuidae]	Not known to occur	No: This species lays eggs on the underside of leaves (AgroAtlas 2009b) and larvae feed externally on young grapevine buds and shoots (Bournier 1976). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Calyptra emarginata</i> Fabricius	Not known to	No: Larvae of these fruit-	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Lepidoptera: Noctuidae]	occur	piercing moths are foliage feeders and adults are associated directly with the fruit and fruit clusters of grapevines (Hanken 2002, Lee <i>et al.</i> 1970). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.			
<i>Calyptra lata</i> (Butler 1881) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Calyptra thalictri</i> (Borkhusen 1790) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Cechenena lineosa</i> (Walker 1856) [Lepidoptera: Sphingidae]	Not known to occur	No: Larvae of sphingid moths are generally foliage feeders (Common 1990). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Cechenena minor</i> (Butler 1875) [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		
<i>Clania variegata</i> (Snellen 1879) [Lepidoptera: Psychidae] (synonym <i>Eumeta variegata</i> (Snellen))	Not known to occur	No: The larvae of this species feed on foliage and also chew the skin of grapes (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Cnephasia longana</i> Haworth 1811 [Lepidoptera: Tortricidae]	Not known to occur	No: This omnivorous leafroller lays eggs on rough barked trunks (Rosenstiel and Ferguson 1944) and larvae feed on grapevine leaves (Norton 1991). Therefore, foliage free, semi-hardwood dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Conogethes punctiferalis</i> Guenée 1854 [Lepidoptera: Crambidae]	Yes (Nielsen <i>et al.</i> 1996)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Cryptophlebia leucotreta</i> Meyrick 1913 [Lepidoptera: Tortricidae]	Not known to occur	No: This tortricid moth lays eggs on the fruit (Grové <i>et al.</i> 1999) and larvae feed within the fruit (Carter 1984). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Ctenopseustis obliquana</i> Walker 1863 [Lepidoptera: Tortricidae]	Not known to occur	No: This tortricid moth lays eggs on buds and larvae feed on the leaves, fruits and buds of hosts (Kay 1979). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Dasychira feminula</i> (Hampson 1891) [Lepidoptera: Lymantriidae]	Not known to occur	No: These species have been recorded on grapevines (Robinson <i>et al.</i> 2008). Larvae of Lymantriid moths are foliage feeders (Common 1990). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Dasychira tenebrosa</i> Walker 1865 [Lepidoptera: Lymantriidae]	Not known to occur		Assessment not required		
<i>Deilephila elpenor</i> Swinhoe 1884 [Lepidoptera: Sphingidae]	Not known to occur	No: This species lays eggs on leaves (CABI 2012a) and larvae feed on the leaves of host plants (Owen 1991). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Desmia funeralis</i> Hübner 1796 [Lepidoptera: Pyralidae]	Not known to occur	No: The grape leaf folder lays eggs on both sides of leaves	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(AliNiazee 1974) and larvae attack leaves (Mead and Webb 2001). Therefore, foliage free dormant cuttings do not provide a pathway for pest.			
<i>Diaphania indica</i> (Saunders 1851) [Lepidoptera: Pyralidae]	Yes (Nielsen <i>et al.</i> 1996)	Assessment not required			
<i>Elibia dolichus</i> (Westwood 1847) [Lepidoptera: Sphingidae]	Not known to occur	No: This moth has been recorded from grapes (Robinson <i>et al.</i> 2008). The larvae of sphingids generally feed only on foliage (Common 1990). Therefore, foliage free dormant cuttings do not provide a pathway for this species	Assessment not required		
<i>Endopiza viteana</i> Clemens 1860 [Lepidoptera: Tortricidae] (synonym: <i>Paralobesia viteana</i> Clemens 1860)	Not known to occur	No: This moth lays eggs on berries, where the larvae then feed (Botero-Garces and Isaacs 2003). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Epiphyas postvittana</i> Walker 1863 [Lepidoptera: Noctuidae]	Yes (Nicholas <i>et al.</i> 1994)	Assessment not required			
<i>Estigmene acraea</i> Drury 1773 [Lepidoptera: Arctiidae]	Not known to occur	No: This species feeds on the leaves of host plants (Stracener 1931). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Eudocima fullonia</i> Clerck 1764 [Lepidoptera: Noctuidae]	Yes (Smith <i>et al.</i> 1997)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Eudocima tyrannus</i> (Guenée 1852) [Lepidoptera: Noctuidae]	Not known to occur	No: This species lays eggs on the underside of leaves of host plants and sometimes on the bark (Kumar and Lal 1983; CABI 2012a). Adults feed on fruit (Hanken 2002). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Eupoecilia ambiguella</i> (Hübner 1796) [Lepidoptera: Tortricidae]	Not known to occur	No: The larvae of this species feed internally on berries and unripe seed of grapes (Frolov 2009). Therefore dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Eulithis diversilineata</i> Hübner 1812 [Lepidoptera: Geometridae]	Not known to occur	No: Larvae of lesser grapevine looper are foliage feeders (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this insect.	Assessment not required		
<i>Eumorpha achemon</i> (Drury 1773) [Lepidoptera: Sphingidae]	Not known to occur	No: This moth lays eggs on the upper surface of the leaves and larvae feed on the foliage (Bournier 1976; Anon 2008). Therefore, foliage free dormant cuttings do not provide a pathway for the Achemon sphinx moth.	Assessment not required		
<i>Eupoecilia ambiguella</i> Hübner 1796 [Lepidoptera: Tortricidae]	Not known to occur	No: This moth lays eggs on buds, bracts and anthophores, less often on	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		young sprouts or on immature berries (AgroAtlas 2011a). Larvae feed on flower buds and flowers (AgroAtlas 2011a). Therefore, dormant cuttings do not provide a pathway for grape moth.			
<i>Euproctis paradoxa</i> (Butler 1886) [Lepidoptera: Lymantriidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Euxoa messoria</i> Harris 1841 [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of these noctuid moths feed on the swelling grape buds (Wright <i>et al.</i> 2010; Williams <i>et al.</i> 2011). Therefore, dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Euxoa scandens</i> Riley 1869 [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Euxoa tessellata</i> (Harris 1841) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Euxoa ochrogaster</i> (Guenée 1852) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Feltia subgothica</i> (Haworth 1809) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of this noctuid moth feed on the swelling grape buds (Williams <i>et al.</i> 2011). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Geina persicidactylus</i> (Fitch) 1855 [Lepidoptera: Pterophoridae]	Not known to occur	No: This moth lays eggs on foliage and hatched larvae feed on the upper surfaces of grape leaves (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Harrisina americana</i> Guérin-Meneville 1844 [Lepidoptera: Zygaenidae]	Not known to occur	No: The grape leaf skeletonizer lays eggs and	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		feeds on the leaves of grapevine (Mead and Webb 2001). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.			
<i>Helicoverpa armigera</i> Hübner 1809 [Lepidoptera: Noctuidae]	Yes (Nielsen <i>et al.</i> 1996)	Assessment not required			
<i>Helicoverpa punctigera</i> Wallengren 1860 [Lepidoptera: Noctuidae]	Yes (Smith <i>et al.</i> 1997)	Assessment not required			
<i>Herpetogramma luctuosalis</i> (Guenée 1854) [Lepidoptera: Pyralidae]	Not known to occur	No: The larvae of this moth feed on grape leaves by rolling the leaves into a cylinder and feeding on them from the inside (Li 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Hippotion celerio</i> (Linnaeus 1758) [Lepidoptera: Sphingidae]	Present (Common 1990)	Assessment not required			
<i>Hyphantria cunea</i> Drury 1770 [Lepidoptera: Arctiidae]	Not known to occur	No: This species lays eggs on the underside of leaves of host plants (Johnson and Lyon 1988). Developing larvae feed on foliage (Warren and Tadic 1970). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Illiberis tenuis</i> (Butler 1877) [Lepidoptera: Zygaenidae]	Not known to occur	No: Larvae feed on young shoots, flowers and leaves of grapevines (Zhang 2005). Therefore, foliage free	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		dormant cuttings do not provide a pathway for this species.			
<i>Ischyja manlia</i> (Cramer 1776) [Lepidoptera: Noctuidae]	Yes (Nielsen <i>et al.</i> 1996)	Assessment not required			
<i>Lacinipolia meditata</i> (Grote 1873) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of these noctuid moths feed on swelling grape buds (Williams <i>et al.</i> 2011). Therefore, dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Lacinipolia renigera</i> (Stephens 1829) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Lithophane antennata</i> (Walker 1858) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of green fruit worm feed on new foliage (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Lobesia botrana</i> Denis & Schiffermüller 1775 [Lepidoptera: Tortricidae]	Not known to occur	No: Larvae of this tortricid moth mainly feed on flowers and berries (Varela <i>et al.</i> 2010). Eggs are laid on flower buds, flower clusters, berries, shoot and leaves during spring and summer (Moreau <i>et al.</i> 2010; Varela <i>et al.</i> 2010). Non-diapausing larvae pupate on the edges of leaves while diapausing larvae pupate in the bark of trunks (Roehrich and Boller 1991). Therefore, dormant, semi-hardwood cuttings do not	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		provide a pathway for this pest.			
<i>Loepa katinka</i> (Westwood 1847) [Lepidoptera: Saturniidae]	Not known to occur	No: Larvae of saturniid moths are foliage feeders (Common 1990). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Mamestra brassicae</i> Linnaeus 1758 [Lepidoptera: Noctuidae]	Not known to occur	No: Larvae of this noctuid moth feed on foliage and lay eggs on the underside of the leaves of host plants (CABI 2012a). Older larvae can also feed on ripening grapes (Voigt 1974). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Marumba gaschkewitschii</i> (Bremer & Grey 1852) [Lepidoptera: Sphingidae]	Not known to occur	No: Larvae of this moth feed on foliage (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Nippoptilia vitis</i> Sasaki 1913 [Lepidoptera: Pterophoridae] (synonym: <i>Stenoptilia vitis</i> Sasaki)	Not known to occur	No: Larvae of this species feed on leaves, stems, and fruit (APHIS–USDA 2002). Larvae may also feed internally on the fruit and seeds of grape, usually causing the young fruit to drop (Li 2004; Zhang 2005). Therefore, foliage free dormant cuttings do not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		provide a pathway for this pest.			
<i>Oraesia emarginata</i> (Fabricius 1794) [Lepidoptera: Noctuidae]	Yes (CSIRO 2013)	Assessment not required			
<i>Oraesia excavata</i> (Butler 1878) [Lepidoptera: Noctuidae]	Not known to occur	No: Adults shelter in foliage and feed on fruits (Li 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Orthodes rufula</i> Grote 1849 [Lepidoptera: Noctuidae]	Not known to occur	No: This moth occurs on grapevines from the time of bud swell to when shoots are several inches long. The larvae feed on grapevine buds and injured buds may fail to develop (Bentley <i>et al.</i> 2008). Dormant cuttings are not preferred feeding sites for this insect and therefore do not provide a pathway for this moth.	Assessment not required		
<i>Orthosia hibisci</i> Guenée 1852 [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of this insect feed on new foliage (Williams <i>et al.</i> 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Paranthrene regalis</i> Butler 1878 [Lepidoptera: Sesiidae]	Not known to occur	<b>Yes:</b> The newly hatched larvae of grape clear-wing moth bore into the tender stems of grapevines; they then develop, overwinter and pupate within the stem of	<b>Yes:</b> Grape clear-wing moth is distributed in China and Korea (Zhou 1991; Seung-Tae <i>et al.</i> 2006). There are similar natural and built environments in parts of Australia that would	<b>Yes:</b> This species damages vines and may cause defoliation and a decline in yield (Li 2004). This species is listed as an insect that can endanger commercial grapevine production in China (Li	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		grapevines (Zhou 1991). Dormant cuttings may harbor overwintering larvae and therefore provide a pathway for grape clear-wing moth.	be suitable for the establishment and spread of this pest.	2004). Therefore, this moth has the potential for economic consequences in Australia.	
<i>Paropta paradoxus</i> Herrich-Schäffer 1851 [Lepidoptera: Cossidae]	Not known to occur	No: This cossid moth lays eggs on the underside of loose bark or on the older wood of grapevines (Plaut 1973). Hatched larvae settle under loose bark and begin feeding. The larvae burrow into the stems and branches of grapevine through dried stubs of pruned canes and excavate galleries along the axes of stems and branches (Plaut 1973). Larvae may also develop under dry bark. This cossid moth overwinters as active immature larvae and diapausing mature prepupal larvae (Plaut 1973). One year old semi-hardwood dormant cuttings are not preferred sites for egg laying and therefore do not provide a pathway for this cossid moth.	Assessment not required		
<i>Pergesa acteus</i> (Cramer 1779) [Lepidoptera: Sphingidae]	Not known to occur	No: This species has been recorded on grapevines (Pittaway and Kitching 2006) and larvae feed on foliage (Common 1990). Therefore, foliage free dormant cuttings	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		do not provide a pathway for this species.			
<i>Peridroma saucia</i> Hübner 1803 [Lepidoptera: Noctuidae]	Not known to occur	No: This moth is associated with <i>Vitis</i> species (Dibble <i>et al.</i> 1979; CABI 2012a) and larvae feed on swelling grape buds (Williams <i>et al.</i> 2011). Therefore, dormant cuttings do not provide a pathway for this noctuid moth.	Assessment not required		
<i>Phalaenoides glycinae</i> Lewin 1805 [Lepidoptera: Noctuidae]	Yes (CSIRO 2005)	Assessment not required			
<i>Platynota stultana</i> Walsingham 1884 [Lepidoptera: Tortricidae]	Not known to occur	No: Omnivorous leafrollers lay eggs on the leaves and newly hatched larvae feed on buds (AliNiazee and Stafford 1972). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Proeulia auraria</i> Clarke 1949 [Lepidoptera: Tortricidae]	Not known to occur	No: These species lay eggs on leaves (Campos <i>et al.</i> 1981) and larvae feed on foliage and fruit (Brown and Passoa 1998; Brown 1999). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Proeulia chrysopteris</i> Butler 1883 [Lepidoptera: Tortricidae]	Not known to occur		Assessment not required		
<i>Proeulia triquetra</i> Obraztsov 1964 [Lepidoptera: Tortricidae]	Not known to occur		Assessment not required		
<i>Psychomorpha epimenis</i> (Drury 1782) [Lepidoptera: Noctuidae]	Not known to occur	No: This species lays eggs on or near new foliage and hatched larvae feed on foliage (Williams <i>et al.</i> 2011). Therefore, foliage free	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		dormant cuttings do not provide a pathway for this pest.			
<i>Rhagastis castor aurifera</i> (Butler 1875) [Lepidoptera: Sphingidae]	Not known to occur	No: These species have been recorded on grapevines (Pittaway and Kitching 2006) and larvae feed on foliage (Common 1990; Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these species.	Assessment not required		
<i>Rhagastis confusa</i> Rothschild and Jordan 1903 [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		
<i>Rhagastis mongoliana</i> (Butler 1876) [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		
<i>Rhynchagrotis cupida</i> (Grote 1864) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of these moths feed on the swelling grape buds (Williams <i>et al.</i> 2011). Therefore, dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Rhynchagrotis placida</i> (Grote 1876) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Sarbanissa subflava</i> (Moore 1877) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of these moths feed on young shoots and leaves of grapevines (Zhang 2005). Therefore, dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Sarbanissa transiens</i> (Walker 1855) [Lepidoptera: Noctuidae]	Not known to occur		Assessment not required		
<i>Spaelotis clandestina</i> (Harris 1862) [Lepidoptera: Noctuidae]	Not known to occur	No: The larvae of this noctuid moth feed on the swelling grape buds (Williams <i>et al.</i> 2011). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Sparganothis pilleriana</i> Denis & Schiffermüller 1776 [Lepidoptera: Tortricidae]	Not known to occur	No: This moth lays eggs on the upper surface of grape leaves (HYPPZ 2008). Hatching larvae shelter under	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		the trunk bark to hibernate. In spring, larval withdrawal from diapause coincides with bud swelling and blossoming and with growth of young leaves (AgroAtlas 2011b). The larvae feed on buds, leaves and young branches and pupate in the folds of leaves (HYPPZ 2008). Therefore, foliage free semi-hardwood dormant cuttings do not provide a pathway for this species.			
<i>Sphecodina caudata</i> (Bremer & Grey 1853) [Lepidoptera: Sphingidae]	Not known to occur	No: The larvae of this moth feed on leaves of grapevines (Zhang 2005). Therefore, dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Spirama retorta</i> (Clerck 1764) [Lepidoptera: Noctuidae]	Not known to occur	No: Larvae of this moth feed on young foliage and new shoots, whereas adults feed on fruits (Kim and Lee 1985; Sambath and Joshi 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Spodoptera exigua</i> (Hübner 1803) [Lepidoptera: Noctuidae]	Yes (Naumann 1993)	Assessment not required			
<i>Spodoptera frugiperda</i> Smith & Abbot 1797 [Lepidoptera: Noctuidae]	Not known to occur	No: <i>Spodoptera</i> species lay eggs on leaves, often near blossoms (Capinera and Fasulo 2006). Larvae and	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		adults feed on leaves, buds and flowers (Balikai <i>et al.</i> 1999; Papademetriou and Dent 2001; Capinera 2008). Therefore, foliage free dormant cuttings do not provide a pathway for this species.			
<i>Spodoptera litura</i> Fabricius 1775 [Lepidoptera: Noctuidae]	Yes (Naumann 1993)	Assessment not required			
<i>Spodoptera praefica</i> Grote 1875 [Lepidoptera: Noctuidae]	Not known to occur	No: <i>Spodoptera</i> species lay eggs on leaves, often near blossoms (Capinera and Fasulo 2006). Larvae and adults feed on leaves, buds and flowers (Balikai <i>et al.</i> 1999; Papademetriou and Dent 2001; Capinera 2008). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Sylepta lunalis</i> Guenee 1854 [Lepidoptera: Pyralidae]	Not known to occur	No: Larvae of this species feed on foliage and destroy the parenchyma tissue of the leaves (Bournier 1976). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Theretra alecto</i> (Linnaeus 1758) [Lepidoptera: Sphingidae]	Not known to occur	No: These species have been recorded on grapevines (Pittaway and Kitching 2006). Sphingid larvae generally feed	Assessment not required		
<i>Theretra boisduvalii</i> (Bugnion 1839) [Lepidoptera: Sphingidae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		only on foliage (Common 1990). Therefore, foliage free dormant cuttings do not provide a pathway for this species.			
<i>Theretra clotho</i> Drury 1773 [Lepidoptera: Sphingidae]	Yes (PHA 2001)	Assessment not required			
<i>Theretra japonica</i> (Boisduval 1869) [Lepidoptera: Sphingidae]	Not known to occur	No: Sphingid larvae feed on grapevine leaves (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Theretra latreillei</i> Macleay 1827 [Lepidoptera: Sphingidae]	Yes (Common 1990)	Assessment not required			
<i>Theretra oldenlandiae</i> Fabricius 1775 [Lepidoptera: Sphingidae]	Yes (Naumann 1993)	Assessment not required			
<i>Theretra pallicosta</i> (Walker 1856) [Lepidoptera: Sphingidae]	Not known to occur	No: This species has been recorded on grapevines (Pittaway and Kitching 2006). Sphingid larvae generally feed only on foliage (Common 1990). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Vitacea polistiformis</i> Harris 1854 [Lepidoptera: Sesiidae]	Not known to occur	No: This species is a root borer and caterpillars damage roots (Bournier 1976). Therefore, root free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Xestia c-nigrum</i> (Linneaus 1958) [Lepidoptera: Noctuidae]	Not known to occur	No: Larvae of this species feed on developing shoots	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		and buds (Dibble <i>et al.</i> 1979). Therefore, dormant cuttings do not provide a pathway for this pest.			
<i>Zeuzera coffeae</i> Nietner 1861 [Lepidoptera: Cossidae]	Not known to occur	<b>Yes:</b> Newly hatched larvae enter young twigs and later move into larger branches or trunks (Cheng 1984). Eggs are laid in strings in cracks of the bark of branches. Therefore, dormant cuttings may provide a pathway for the larvae of this species.	<b>Yes:</b> This species has established in areas with a wide range of climatic conditions (Waller <i>et al.</i> 2007) and can spread naturally in infested propagative material. Therefore, this species has the potential to establish and spread in Australia.	<b>Yes:</b> No information is available on losses caused by this moth on grapevines, but it causes considerable damage in coffee trees due to destruction of branches through boring activity (Waller <i>et al.</i> 2007). Therefore, this moth has the potential for economic consequences in Australia.	<b>Yes</b>
<b>ORTHOPTERA (grasshoppers, crickets)</b>					
<i>Austracris guttulosa</i> Walker 1870 [Orthoptera: Acrididae]	Yes (Coombe and Dry 1992)	Assessment not required			
<i>Austroicetes cruciata</i> Saussure 1888 [Orthoptera: Acrididae]	Yes (PHA 2001)	Assessment not required			
<i>Chortoicetes terminifera</i> Walker 1870 [Orthoptera: Acrididae]	Yes (PHA 2001)	Assessment not required			
<i>Melanoplus devastator</i> Scudder 1778 [Orthoptera: Acrididae]	Not known to occur	No: This species feeds on young foliage (Schell <i>et al.</i> 2007). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Phaulacridium vittatum</i> Sjöstedt 1920 [Orthoptera: Acrididae]	Yes (PHA 2001)	Assessment not required			
<i>Valanga irregularis</i> Walker 1870 [Orthoptera: Acrididae]	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<b>PSOCOPTERA (booklice)</b>					
<i>Ectopsocus briggsi</i> McLachlan 1899 [Psocoptera: Ectopsocidae]	Yes (Ahadiyat and Zangeneh 2007)	Assessment not required			
<i>Graphopsocus cruciatus</i> Linnaeus 1768 [Psocoptera: Ectopsocidae]	Not known to occur	No: This species feeds on the microflora on leaves (Greenwood 1988). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<b>THYSANOPTERA (thrips)</b>					
<i>Aeolothrips fasciatus</i> (Linnaeus 1758) [Thysanoptera: Aeolothripidae]	Yes(PHA 2001)	Assessment not required			
<i>Aeolothrips intermedius</i> Bagnall 1934 [Thysanoptera: Aeolothripidae]	Not known to occur	No: These species are associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for these thrips.	Assessment not required		
<i>Aeolothrips melaleucus</i> Haliday 1852 [Thysanoptera: Aeolothripidae]	Not known to occur		Assessment not required		
<i>Aeolothrips vittatus</i> Haliday 1836 [Thysanoptera: Aeolothripidae]	Not known to occur		Assessment not required		
<i>Caliothrips fasciatus</i> Pergande 1895 [Thysanoptera: Thripidae]	Not known to occur	No: This species feeds on the leaves, stems, buds and flowers (Mound 2008). The eggs are laid in leaf tissue (Harman <i>et al.</i> 2007). Therefore, foliage free dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Chirothrips manicatus</i> Haliday 1836 [Thysanoptera: Thripidae]	Not known to occur	No: These <i>Chirothrips</i> species are associated with foliage	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Chirothrips molestus</i> Priesner 1926 [Thysanoptera: Thripidae]	Not known to occur	and inflorescences of grapevines (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Dendrothrips saltatrix</i> Uzel 1895 [Thysanoptera: Thripidae]	Not known to occur	No: This thrip is associated with the foliage and inflorescences of grapevines (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Dictyothrips betae</i> Uzel 1895 [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Drepanothrips reuteri</i> Uzel 1895 [Thysanoptera: Thripidae]	Not known to occur	No: Grape thrips lay eggs on the young leaves and buds of <i>Vitis vinifera</i> (Marullo 2009) and feed on shoot tips and leaves (Flint 2006). Dormant cuttings are not the preferred egg laying site and therefore do not provide a pathway for grape thrips.	Assessment not required		
<i>Frankliniella australis</i> Morgan 1925 [Thysanoptera: Thripidae]	Not known to occur	No: This species feeds and lays eggs in the flowers of host plants (Borbon <i>et al.</i>	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		2008). Therefore, dormant cuttings do not provide a pathway for this pest.			
<i>Frankliniella intonsa</i> (Trybom 1895) [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with the foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Frankliniella occidentalis</i> Pergande 1895 [Thysanoptera: Thripidae]	Present (PHA 2001)	Assessment not required			
<i>Frankliniella tritici</i> Fitch 1855 [Thysanoptera: Thripidae]	Not known to occur	No: This species feeds on flowers and lays eggs on leaf petioles (Reitz 2002). Therefore, foliage and flower free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Haplothrips acanthoscelis</i> (Karny 1909) [Thysanoptera: Phlaeothripidae]	Not known to occur	No: These <i>Haplothrips</i> species are associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Haplothrips aculeatus</i> (Fabricius 1803) [Thysanoptera: Phlaeothripidae]	Not known to occur		Assessment not required		
<i>Haplothrips froggatti</i> Hood 1918 [Thysanoptera: Phlaeothripidae]	Yes (PHA 2001)	Assessment not required			
<i>Haplothrips kurdjumovi</i> Karny 1913 [Thysanoptera: Phlaeothripidae]	Not known to occur	No: This species is associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009).	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		Therefore, foliage free dormant cuttings do not provide a pathway for this pest.			
<i>Haplothrips leucanthemi</i> (Schrank 1781) [Thysanoptera: Phlaeothripidae]	Yes (PHA 2001)	Assessment not required			
<i>Haplothrips victoriensis</i> Bagnall 1918 [Thysanoptera: Phlaeothripidae]	Yes (PHA 2001)	Assessment not required			
<i>Heliothrips haemorrhoidalis</i> Bouché 1833 [Thysanoptera: Thripidae]	Present (Naumann 1993)	Assessment not required			
<i>Heliothrips sylvanus</i> Faure 1933 [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with foliage (Roditakis and Roditakis 2007). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Neohydatothrips gracilicornis</i> (Williams 1916) [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Retithrips syriacus</i> Mayet 1890 [Thysanoptera: Thripidae]	Not known to occur	No: This species feeds on leaves (Doganlar and Yigit 2002) and lays eggs on leaf surfaces (CPPDR 1994). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Rhipiphorocthrrips cruentatus</i> Hood 1991 [Thysanoptera: Thripidae]	Not known to occur	No: This species feeds on foliage (Bournier 1976;	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		Dahiya and Lakra 2001) and lays eggs on the underside of leaves (Kulkarni <i>et al.</i> 2007). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.			
<i>Rubiothrips vitis</i> (Priesner 1933) [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Scirtothrips citri</i> Moulton 1909 [Thysanoptera: Thripidae]	Not known to occur	No: <i>Scirtothrips</i> species are associated with the foliage of <i>Vitis</i> species (Arpaia and Morse 1991; Roditakis and Roditakis 2007; Nietschke <i>et al.</i> 2008). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Scirtothrips dorsalis</i> Hood 1919 [Thysanoptera: Thripidae]	Yes (PHA 2001)	Assessment not required			
<i>Scirtothrips mangiferae</i> Priesner 1932 [Thysanoptera: Thripidae]	Not known to occur	No: <i>Scirtothrips</i> species are associated with the foliage of <i>Vitis</i> species (Arpaia and Morse 1991; EPPO 2005; Roditakis and Roditakis 2007; Nietschke <i>et al.</i> 2008). Therefore, foliage free dormant cuttings do not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		provide a pathway for this pest.			
<i>Thrips australis</i> Bagnall 1915 [Thysanoptera: Thripidae]	Yes (Naumann 1993)	Assessment not required			
<i>Thrips fulvipes</i> Bagnall 1923 [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		
<i>Thrips hawaiiensis</i> Morgan 1913 [Thysanoptera: Thripidae]	Yes (PHA 2001)	Assessment not required			
<i>Thrips imagines</i> Bagnall 1926 [Thysanoptera: Thripidae]	Present (Naumann 1993)	Assessment not required			
<i>Thrips physapus</i> Linnaeus 1758 [Thysanoptera: Thripidae]	Not known to occur	No: These thrips species are associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for these pests.	Assessment not required		
<i>Thrips pillichii</i> Priesner 1924 [Thysanoptera: Thripidae]	Not known to occur		Assessment not required		
<i>Thrips tabaci</i> Lindeman 1889 [Thysanoptera: Thripidae]	Yes (Naumann 1993)	Assessment not required			
<i>Thrips validus</i> Uzel 1895 [Thysanoptera: Thripidae]	Not known to occur	No: This species is associated with foliage and inflorescences (Vasiliu-Oromulu <i>et al.</i> 2009). Therefore, foliage free dormant cuttings do not provide a pathway for this pest.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<b>PATHOGENS</b>					
<b>BACTERIA</b>					
<i>Pantoea agglomerans</i> (Beijerinck 1888) Gavini <i>et al.</i> 1989 [Enterobacteriales: Enterobacteriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pseudomonas fluorescens</i> Migula 1895 [Pseudomonadales: Pseudomonadaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pseudomonas syringae</i> subsp. <i>syringae</i> van Hall 1902 [Pseudomonadales: Pseudomonadaceae]	Yes (Whitelaw-Weckert <i>et al.</i> 2011)	Assessment not required			
<i>Pseudomonas viridiflava</i> Burkholder 1930) Dowson 1939 [Pseudomonadales: Pseudomonadaceae]	Yes (PHA 2001)	Assessment not required			
<i>Rhizobium radiobacter</i> (Beijerinck and van Delden 1902) Pribram 1933 [Rhizobiales: Rhizobiaceae]	Yes (PHA 2001)	Assessment not required			
<i>Rhizobium vitis</i> (Ophel & Kerr 1990) Young <i>et al.</i> [Rhizobiales: Rhizobiaceae]	Yes (Gillings and Ophel-Keller 1995)	Assessment not required			
<i>Xanthomonas campestris</i> pv. <i>viticola</i> (Nayudu 1972) Dye 1978 [Xanthomonadales: Xanthomonadaceae]	Not known to occur	<b>Yes:</b> <i>Xanthomonas campestris</i> pv. <i>viticola</i> (Xcv) survives in infected plants as an epiphyte on aerial plant parts and may be carried in infected transplants and cuttings (Nascimento and Mariano 2004; Peixoto <i>et al.</i> 2007). Therefore, dormant cuttings may provide a pathway for this bacterium.	<b>Yes:</b> Xcv has established in areas with a wide range of climatic conditions (Trindade <i>et al.</i> 2005) and may spread naturally in infected propagative material (Nascimento and Mariano 2004; Peixoto <i>et al.</i> 2007). Multiplication and marketing of infected propagative material will help spread this bacterium within Australia.	<b>Yes:</b> This bacterium causes leaf blight and cankers on stems and petioles and causes extensive foliage death. It also causes irregular colour and size in berries and may cause necrotic lesions (Nascimento and Mariano 2004), reducing the yield and quality of the grapes. The development of grapevine bacterial canker in Brazil has caused severe crop losses	<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
			Therefore, this bacterium has the potential for establishment and spread in Australia.	(Nascimento <i>et al.</i> 2005). Therefore, this bacterium has the potential for economic consequences in Australia.	
<i>Xanthomonas campestris</i> pv. <i>vitiscarnosae</i> (Moniz & Patel 1958) Dye 1978 [Xanthomonadales: Xanthomonadaceae]	Not known to occur	Assessment not required <sup>8</sup>			
<i>Xanthomonas campestris</i> pv. <i>vitistrifoliae</i> (Padhya <i>et al.</i> 1965) Dye 1978) Dye [Xanthomonadales: Xanthomonadaceae]	Not known to occur	Assessment not required <sup>9</sup>			
<i>Xanthomonas campestris</i> pv. <i>vitiswoodrowii</i> (Patel & Kulkarni 1951) Dye 1978 [Xanthomonadales: Xanthomonadaceae]	Not known to occur	Assessment not required <sup>10</sup>			
<i>Xylella fastidiosa</i> (Wells <i>et al.</i> 1987) <sup>11</sup> – grapevine strain [Xanthomonadales: Xanthomonadaceae]	Not known to occur	<b>Yes:</b> <i>Xylella fastidiosa</i> multiplies and spreads exclusively within the xylem (Purcell 2001). Diseased stems often mature irregularly and show patches of brown and green tissue. The grapevine strain of <i>X.</i>	<b>Yes:</b> Grapevine strain of <i>X. fastidiosa</i> has established in areas with a wide range of climatic conditions (Mizell <i>et al.</i> 2008) and may spread naturally in infected propagative material (Frison and Ikin 1991). Multiplication	<b>Yes:</b> Grapevine strain of <i>X. fastidiosa</i> causing Pierce's disease is a major constraint on grapevine production in the USA and tropical America (CABI/EPPO 1990). Grapevines affected by this pathogen usually die within 1–5 years of infection	<b>Yes</b>

<sup>8</sup> *V. campestris* pv. *vitiscarnosae* attacks *V. carnosae*. *Vitis carnosae* is not an important species of *Vitis* for commercial viticulture, scion cultivars, rootstocks or in breeding programs and therefore will not be imported into Australia. Additionally, *Vitis carnosae* is currently not permitted entry into Australia. Consequently, *V. campestris* pv. *vitiscarnosae* is not on the pathway.

<sup>9</sup> *V. campestris* pv. *vitistrifoliae* attacks *V. trifolia*. *Vitis trifolia* is not an important species of *Vitis* for commercial viticulture, scion cultivars, rootstocks or in breeding programs and therefore will not be imported into Australia. Additionally, *Vitis trifolia* is currently not permitted entry into Australia. Consequently, *V. campestris* pv. *vitistrifoliae* is not on the pathway.

<sup>10</sup> *Xanthomonas campestris* pv. *vitiswoodrowii* attacks *V. woodrowii*. *Vitis woodrowii* is not an important species of *Vitis* for commercial viticulture, scion cultivars, rootstocks or in breeding programs and therefore will not be imported into Australia. Additionally, *Vitis woodrowii* is currently not permitted entry into Australia. Consequently, *V. campestris* pv. *vitiswoodrowii* is not on the pathway.

<sup>11</sup> Strains of this bacterium are the causal agent of phony peach disease (PPD), plum leaf scald, Pierce's disease (PD) of grapes, citrus variegated chlorosis (CVC) and leaf scorch of almond, coffee, elm, oak, oleander pear, and sycamore (Mizell *et al.* 2008). Only information on Pierce's disease (PD) grape strain has been used in this section.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		<i>fastidiosa</i> proliferates only in xylem vessels, in roots, stems and leaves. Therefore, propagative material provides a pathway for the grapevine strain of <i>X. fastidiosa</i> .	and marketing of infected propagative material will help spread this bacterium within Australia. CLIMEX predictions indicate that grape growing regions across southern Australia would be highly suitable for this bacterium (Hoddle 2004). Therefore, this bacterium has the potential for establishment and spread in Australia.	(Pearson and Goheen 1988). <i>X. fastidiosa</i> is an EPPO A1 quarantine pest and is also of quarantine significance for COSAVE. Presence of this bacterium in Australia would impact upon Australia's ability to access overseas markets. Therefore, this bacterium has the potential for significant economic consequences in Australia.	
<i>Xylophilus ampelinus</i> (Panagopoulos 1969) Willems <i>et al.</i> 1987 [Xanthomonadales: Xanthomonadaceae]	Not known to occur	<b>Yes:</b> <i>Xylophilus ampelinus</i> is a systemic pathogen infecting xylem (Grall and Manceau 2003) and overwinters in plant tissue. Primary infection occurs on one or two year old shoots (Rid� <i>et al.</i> 1977). This bacterium often presents as a latent infection (Rid� <i>et al.</i> 1983; Panagopoulos 1987). This may lead to the propagation and distribution of infected propagative material, suggesting that this bacterium could be introduced into Australia.	<b>Yes:</b> <i>Xylophilus ampelinus</i> has established in areas with a wide range of climatic conditions (Botha <i>et al.</i> 2001; Manceau <i>et al.</i> 2005; CABI/EPPO 1999; Dre� <i>et al.</i> 2005) and has spread naturally in infected propagative material (Frison and Ikin 1991). Multiplication and marketing of latently infected propagative material will help spread this bacterium within Australia. Therefore, this bacterium has the potential for establishment and spread in Australia.	<b>Yes.</b> <i>Xylophilus ampelinus</i> is a destructive pathogen of multiple grapevine cultivars (Serfontein <i>et al.</i> 1997). <i>Xylophilus ampelinus</i> is an EPPO A2 quarantine organism (OEPP/EPPO 1984) and is also of quarantine significance for NAPPO and the IAPSC. Presence of this bacterium in Australia would impact upon Australia's ability to access overseas markets. Therefore, this bacterium has the potential for economic consequences in Australia.	<b>Yes</b>
<b>FUNGI</b>					
<i>Acanthonitschkea tristis</i> (Pers.) Nannf. [Coronophorales: Nitschkiaceae]	Not known to occur	No: This species is found on the decaying wood and bark of host plants (Miller and Huhndorf 2009). Therefore,	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		dormant cuttings do not provide a pathway for this fungus.			
<i>Acremonium alternatum</i> Link [Hypocreales: Unassigned]	Not known to occur	No: This fungus is a mycoparasite and consequently feeds on other pathogens occurring on the plant (Romero <i>et al.</i> 2003). For instance, it is known to control powdery mildews (Romero <i>et al.</i> 2003). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Acremonium charticola</i> (Lindau) W. Gams [Hypocereales: Incertae sedis]	Not known to occur	<b>Yes:</b> This fungus is isolated from vascular tissues of grapevines (Halleen <i>et al.</i> 2005). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has a wide distribution (Farr and Rossman 2011) and parts of Australia would be suitable for its establishment and spread. Distribution of infected propagative material would help spread this fungus in Australia. Therefore, this species has the potential for establishment and spread in Australia.	No: This fungus is considered to be non-pathogenic (Halleen <i>et al.</i> 2005). There is no evidence that it has the potential for economic consequences in Australia.	
<i>Acremonium strictum</i> W. Gams [Hypocereales: Incertae sedis]	Yes (McGee 1989; PHA 2001)	Assessment not required			
<i>Acrogenotheca ornata</i> Deighton & Pirozynski [Unassigned]	Not known to occur	No: Other members of this genus are associated with sooty molds (Reynolds 1971). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Acrospermum viticola</i> Ikata & Hitomi [Acrospermales: Acrospermaceae]	Not known to occur	No: This fungus is associated with grapevine foliage (Li 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Acrostalagmus luteoalbus</i> (Link) Zare et al. [Hypocreales: Hypocreaceae]	Not known to occur	No: This species is commonly found in soil (Thormann and Rice 2007). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Actinomucor elegans</i> (Eidam) CR Benj. & Hesselt [Mucorales: Mucoraceae]	Yes (PHA 2001)	Assessment not required			
<i>Aecidium cissigenum</i> Welw. [Pucciniales: Unassigned]	Not known to occur	<b>Yes:</b> <i>Aecidium</i> species produce small yellowing pustules, either scattered or densely distributed on the lower leaf surface and occasionally on petioles, young shoots and rachises of host plants (Pearson and Goheen 1988). Therefore, dormant cuttings may provide a pathway for these rust fungi.	<b>Yes:</b> Grape rusts have established in areas with a wide range of climatic conditions (Pearson and Goheen 1988), and could spread naturally in infested propagative material. Therefore, these fungi have the potential for the establishment and spread in Australia.	No: There is limited information on these species. Although these species are reported to occur on <i>Vitis</i> species (Pearson and Goheen 1988), a literature search failed to provide information on the economic importance of these <i>Aecidium</i> species. Therefore, these species are not considered to be economically significant.	
<i>Aecidium guttatum</i> Kunze [Pucciniales: Unassigned]	Not known to occur				
<i>Aecidium vitis</i> Smith [Pucciniales: Unassigned]	Not known to occur				
<i>Aleurodiscus botryosus</i> Burt [Russulales: Stereaceae]	Not known to occur	No: This species occurs on dead stems (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Alternaria alternata</i> (Fr.) Keissl. [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Alternaria tenuis</i> Link [Pleosporales:	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Pleosporaceae]					
<i>Alternaria tenuissima</i> (Kunze) Wiltshire [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Alternaria viticola</i> Brunaud [Pleosporales: Pleosporaceae]	Not known to occur	<b>Yes:</b> <i>Alternaria viticola</i> mainly attacks young, tender stalks (Ma <i>et al.</i> 2004). This fungus overwinters on tendrils, branches and in bud scales (Ma <i>et al.</i> 2004). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> <i>Alternaria viticola</i> has established in areas with a wide range of climatic conditions (Ma <i>et al.</i> 2004) and may spread naturally in infected propagative material. Therefore, this fungus has the potential for establishment and spread in Australia.	<b>Yes:</b> <i>Alternaria viticola</i> can cause serious drop off of flowers and young fruit. Grape production has been seriously damaged in some areas of China. Yield losses of 30–40% have been reported from Xinjiang province (Ma <i>et al.</i> 2004) and 30–50% in southeast Shandong (Zhu <i>et al.</i> 2006). Therefore, this fungus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Alternaria vitis</i> Cavara [Pleosporales: Pleosporaceae]	Not known to occur	No: <i>Alternaria vitis</i> is associated with the foliage of grapevines (Suhag <i>et al.</i> 1982). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Amerosporium concinnum</i> Petr. [Helotiales: Sclerotiniaceae]	Not known to occur	No: This species occurs on the dead stems of host plants (Hayova and Minter 2009). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Ampelomyces quisqualis</i> Ces [Anamorphic Phaeosphaeriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Amphisphaeria sylvana</i> Saccardo & Spegazzini [Xylariales: Amphisphaeriaceae]	Not known to occur	No: Members of this genus are associated with the dried stems of host plants (Rao	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		1965). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Apiospora montagnei</i> Saccardo [Incertae sedis: Apiosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Aplosporella beaumontiana</i> S. Ahmad [Botryosphaerales: Botryosphaeriaceae]	Not known to occur	No: Fungi in this genus are associated with thin dead twigs (Damm <i>et al.</i> 2007). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Aplosporella fabaeformis</i> (Pass. & Thüm.) Petr. & Syd. [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required		
<i>Armillaria limonea</i> (G. Stev.) Boesew. [Agaricales: Physalacriaceae]	Not known to occur	No: Members of the genus <i>Armillaria</i> occur in the roots of the host plant and cause root rot (Farr <i>et al.</i> 1989; van der Kamp and Hood 2002; CABI 2012a). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Armillaria luteobubalina</i> Watling & Kile [Agaricales: Physalacriaceae]	Yes (Cook and Dubé 1989)	Assessment not required	.		
<i>Armillaria mellea</i> (Vahl) P. Kumm. [Agaricales: Physalacriaceae]	Not known to occur	No: This soil-borne fungus survives in infected wood and roots below ground (Flaherty <i>et al.</i> 1992). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Armillaria novae-zelandiae</i> (G. Stev.) Boesew. [Agaricales: Physalacriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Armillaria tabescens</i> (Scop.) Emel [Agaricales: Physalacriaceae]	Not known to occur	No: Members of the genus <i>Armillaria</i> occur in the roots of the host plant and cause root	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		rot (Farr <i>et al.</i> 1989; van der Kamp and Hood 2002; CABI 2012a). Therefore, root free dormant cuttings do not provide a pathway for this fungus.			
<i>Arxiomyces vitis</i> (Fuckel) P.F. Cannon & D. Hawksw. [Melanosporales: Ceratostomataceae]	Not known to occur	No: This fungus occurs on the bark of woody shrubs and trees (Ferreira <i>et al.</i> 2005). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Ascochyta ampelina</i> Sacc. [Pleosporales: Incertae sedis]	Not known to occur	No: <i>Ascochyta ampelina</i> has been recorded on grapes, causing leaf spot (Kiewnick 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Ascospora viticola</i> Nasyrov [Incertae sedis]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species, causing leaf spot (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Aspergillus aculeatus</i> Iizuka [Eurotiales: Trichocomaceae]	Not known to occur	No: Members of this genus occur on the fruits and seeds of the host plant, causing storage rot (Farr <i>et al.</i> 1989; Leong <i>et al.</i> 2006; CABI 2012a). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Aspergillus carbonarius</i> (Bainier) Thom [Eurotiales: Trichocomaceae]	Yes (Leong <i>et al.</i> 2006)	Assessment not required			
<i>Aspergillus flavus</i> Link. [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Aspergillus glaucus</i> (L.) Link [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Aspergillus niger</i> var. <i>niger</i> Tiegh. [Eurotiales: Trichocomaceae]	Yes (Cook and Dubé 1989)	Assessment not required			
<i>Aspergillus wentii</i> Wehmer [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Asperisporium minutulum</i> (Sacc.) Deighton [Unassigned]	Not known to occur	No: These species occur on the leaves of the plant, causing leaf spot (Farr <i>et al.</i> 1989; Farr and Rossman 2011). <i>Asperisporium vitiphyllum</i> also occurs on fruit (USDA 2005). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Asperisporium vitiphyllum</i> (Speschnew) Deighton [Unassigned]	Not known to occur		Assessment not required		
<i>Asterina viticola</i> AK Kar & SN Ghosh [Capnodiales: Asterinaceae]	Not known to occur	No: <i>Asterina viticola</i> has been recorded on leaves of <i>Vitis</i> species (Hosagoudar 2003). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Athelia arachnoidea</i> (Berk.) Jülich [Atheliales: Atheliaceae]	Not known to occur	No: This fungus occurs on the wood and roots of plants (Farr <i>et al.</i> 1989; Farr and Rossman 2011). Therefore, root free dormant cuttings do not provide a pathway for this	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		fungus.			
<i>Aureobasidium pullulans</i> var. <i>pullulans</i> (de Bary) G. Arnaud [Dothideales: Dothioraceae]	Yes (Simmonds 1966)	Assessment not required			
<i>Bactrodesmium pallidum</i> MB Ellis [Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus occurs on wood (Tsui <i>et al.</i> 2003). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Bartalinia robillardoides</i> Tassi [Xylariales: Amphispheariaceae]	Yes (PHA 2001)	Assessment not required			
<i>Berkleasium corticola</i> (P. Karst.) R.T. Moore [Pelopsporales Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus occurs on dead wood (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Bertia vitis</i> R. Schulzer [Coronophorales: Bertiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, <i>Bertia</i> species are mainly associated with wood and dead limbs of	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		forest trees (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Bipolaris papendorffii</i> (Aa) Alcorn [Pleosporales: Pleosporaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Biscogniauxia capnodes</i> var. <i>capnodes</i> (Berk.) YM Ju & JD Rogers [Xylariales: Xylariaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts these fungi occur on wood (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Biscogniauxia mediterranea</i> var. <i>mediterranea</i> (De Not.) Kuntze [Xylariales: Xylariaceae]	Not known to occur		Assessment not required		
<i>Bispora antennata</i> (Pers.) EW Mason [Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this dematiaceous fungus is saprobic on wood (Ellis and Ellis 1997). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Botryodiplodia palmarum</i> (Cooke) Petr. & Syd. [Diaporthales: Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts fungus causes sett rot	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Sharma and Sharma 2006). Therefore, root free dormant cuttings do not provide a pathway for this fungus.			
<i>Botryodiplodia vitis</i> Sousa da Câmara [Diaporthales: Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since being reported on <i>Vitis</i> species in 1969 in Pakistan (Farr and Rossman 2011), it has not been reported from any other country. This indicates that dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Botryosphaeria australis</i> Slippers <i>et al.</i> [Botryosphaeriales: Botryosphaeriaceae] (synonym: <i>Fusicoccum australe</i> Slippers <i>et al.</i> ; <i>Neofusicoccum australe</i> (Slippers <i>et al.</i> ) Crous <i>et al.</i> )	Yes (Taylor <i>et al.</i> 2005)	Assessment not required			
<i>Botryosphaeria corticola</i> A.J.L. Phillips, A. Alves & J. Luque [Botryosphaeriales: Botryosphaeriaceae] (synonym: <i>Diplodia corticola</i> Phillips <i>et al.</i> )	Not known to occur	<b>Yes:</b> This fungus has been recorded on <i>Vitis</i> species (Carlucci and Frisullo 2009). This fungus is reported to cause cankers in the vascular tissue of one year old canes, spurs and cordons in Texas (Úrbez-Torres <i>et al.</i> 2010b). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Carlucci and Frisullo 2009, Úrbez-Torres <i>et al.</i> 2010b) and may spread naturally in infected propagative material. Therefore, this fungus has the potential for establishment and spread in Australia.	No: This fungus causes dieback of young shoots, defoliation, sub cortical brown streaks on the canes, and wedge-shaped necrotic areas within trunks and branches (Carlucci and Frisullo 2009). This fungus causes cankers, vascular necrosis and dieback in oak ( <i>Quercus</i> ) species (Dreaden <i>et al.</i> 2011). While this species can have strong pathogenic effects on	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
				cork oak, the fungus only colonises decorticated trunks after cork extraction (Luque <i>et al.</i> 2008). It is considered to be one of the less virulent of the <i>Botryosphaeriaceae</i> species (Gubler <i>et al.</i> 2010)	
<i>Botryosphaeria dothidea</i> (Moug.) Ces. & De Not. [Botryosphaeriales: Botryosphaeriaceae]	Yes (Pitt <i>et al.</i> 2009)	Assessment not required			
<i>Botryosphaeria lutea</i> AJL Phillips [Botryosphaeriales: Botryosphaeriaceae] (synonyms: <i>Fusicoccum luteum</i> Pennycook & Samuels; <i>Neofusicoccum luteum</i> (Pennycook & Samuels) Crous <i>et al.</i> )	Yes (Qui <i>et al.</i> 2011)	Assessment not required			
<i>Botryosphaeria obtusa</i> (Schwein.) Shoemaker [Botryosphaeriales: Botryosphaeriaceae]	Yes (Castillo-Pando <i>et al.</i> 2001)	Assessment not required			
<i>Botryosphaeria parva</i> Pennycook & Samuels [Botryosphaeriales: Botryosphaeriaceae] (synonym: <i>Fusicoccum parvum</i> Pennycook & Samuels; <i>Neofusicoccum parvum</i> (Pennycook & Samuels) Crous <i>et al.</i> )	Yes (Pitt <i>et al.</i> 2009)	Assessment not required			
<i>Botryosphaeria rhodina</i> (Berk. & M.A. Curtis) Arx [Botryosphaeriales: Botryosphaeriaceae] (synonym: <i>Lasiodiplodia theobromae</i> (Pat.) Griffon & Maubl.)	Yes (Taylor <i>et al.</i> 2005)	Assessment not required			
<i>Botryosphaeria ribis</i> Grossenb. & Duggar [Botryosphaeriales: Botryosphaeriaceae] (synonym: <i>Fusicoccum tingens</i> Goid.)	Yes (Constable and Drew 2004)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Botryosphaeria stevensii</i> Shoemaker [Botryosphaerales: Botryosphaeriaceae] (synonym: <i>Diplodia mutila</i> (Fr.) Mont.)	Yes (Taylor <i>et al.</i> 2005)	Assessment not required			
<i>Botryosphaeria viticola</i> AJL Phillips & J Luque [Botryosphaerales: Botryosphaeriaceae] ( <i>Dothiorella viticola</i> AJL Phillips & J Luque)	Yes (Wunderlich <i>et al.</i> 2008)	Assessment not required			
<i>Botrytis ampelophila</i> Speg[Helotiales: Sclerotiniaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis riparia</i> (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis riparia</i> in 1973 in Argentina (Farr and Rossman 2011), it has not been reported from any other country, indicating that dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Botrytis cinerea</i> Pers. [Helotiales: Sclerotiniaceae]	Yes (Rogiers <i>et al.</i> 2005)	Assessment not required			
<i>Briosia ampelophaga</i> Cavara [Unassigned]	Not known to occur	No: This species is associated with foliage and causes leaf spot in <i>Vitis</i> (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Cadophora luteo-olivacea</i> (JFH Beyma) TC Harr. & McNew [Helotiales: Leotiomyetidae] (synonym: <i>Phialophora luteo-olivacea</i> JFH Beyma)	Not known to occur	<b>Yes:</b> This endophytic fungus attacks young grapevines (Gramaje <i>et al.</i> 2010; Navarrete <i>et al.</i> 2010) and has been isolated from the	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Prodi <i>et al.</i> 2008; Gramaje and Armengol 2011)	<b>Yes:</b> This species is involved in the decline of young grapevines in vineyards and nurseries (Gramaje <i>et al.</i> 2010) and is common on grapevines affected	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		vascular tissue of grapevines (Halleen <i>et al.</i> 2007). Infection of this fungus can be symptomatic (Navarrete <i>et al.</i> 2010) or asymptomatic (Halleen <i>et al.</i> 2007). Therefore, this fungus has the potential to be on the pathway of grapevine dormant cuttings.	and may spread naturally in infected propagative material. Therefore, this fungus has the potential for establishment and spread in Australia.	by esca and Petri disease in parts of its current range (Gramaje and Armengol 2011). This species has also been reported as the causal agent of kiwifruit leader dieback (Riccioni <i>et al.</i> 2007; Prodi <i>et al.</i> 2008). Therefore, this fungus has the potential for economic consequences in Australia.	
<i>Cadophora melinii</i> Nannf. [Helotiales: Leotiomycetidae] (synonym: <i>Phialophora melinii</i> (Nannf.) Conant)	Not known to occur	<b>Yes:</b> This species is associated with trunk diseases of young grapevines (Gramaje <i>et al.</i> 2010). Therefore, this species has the potential to be on the pathway of dormant grapevine cuttings.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Prodi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2010; Navarrete <i>et al.</i> 2010) and may spread naturally in infected propagative material. Therefore, this fungus has the potential for establishment and spread in Australia.	<b>Yes:</b> This fungus has been isolated from young grapevines affected by esca and Petri disease (Gramaje <i>et al.</i> 2010; Gramaje <i>et al.</i> 2011). There is no evidence that this species is an economically important pathogen of grapevines, however it is associated with trunk hypertrophy and elephantiasis in kiwifruit (Prodi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2011; Spadaro <i>et al.</i> 2011) resulting in reduced foliage and small, unsalable fruits (Prodi <i>et al.</i> 2008). Therefore, this species has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Calonectria kyotensis</i> Terash. 1968 [Hypocreales: Nectriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Camarosporium viniferum</i> S. Ahmad [Botryosphaerales: Unassigned]	Not known to occur	<b>Yes:</b> This species occurs on <i>Vitis</i> branches (Farr and Rossman 2011). Therefore, dormant cuttings may provide	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Farr and Rossman	No: There is no evidence that this species has the potential for economic consequences.	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		a pathway for this fungus.	2011) and may spread naturally in infected propagative material. Therefore, this fungus has the potential for establishment and spread in Australia.		
<i>Campylocarpon fasciculare</i> Schroers <i>et al.</i> [Hypocreales: Nectriaceae]	Not known to occur	No: These species are associated with grapevine roots causing sunken necrotic root lesions (Halleen <i>et al.</i> 2006a). Therefore, root free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Campylocarpon pseudofasciculare</i> Halleen <i>et al.</i> [Hypocreales: Nectriaceae]	Not known to occur		Assessment not required		
<i>Capnodinula tonduzii</i> Speg. [Incertae sedis: Pseudoperisporiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species from Costa Rica in 1973 (Farr and Rossman 2011), it has not been reported from any other country, indicating that dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Capnodium elongatum</i> Berk. & Desm. [Capnodiales: Capnodiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus occurs on leaves (Farr and Rossman	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Capnodium salicinum</i> Mont [Capnodiales: Capnodiaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Capronia mansonii</i> (Schol-Schwarz) Müller <i>et al.</i> [Chaetothyriales: Herpotrichiellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus occurs on leaves (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Cephalotrichum microsporum</i> (Sacc.) PM Kirk [Microascales: Microascaceae] (synonym: <i>Doratomyces microsporus</i> (Sacc.) F.J. Morton & G. Sm.)	Yes (Eicker 1973; PHA 2001)	Assessment not required			
<i>Cephalotrichum stemonitis</i> (Pers.) Nees [Microascales: Microascaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Cercospora coryneoides</i> Sävul. & Rayss [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: These <i>Cercospora</i> species have been recorded on <i>Vitis</i> species (Farr and Rossman 2011). Generally, <i>Cercospora</i> species occur on the leaves of host plants and cause leaf spot (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for these	Assessment not required		
<i>Cercospora daspurensis</i> AK Kar & M Mandal [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora fuckelii</i> (Thüm.) Jacz. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora judaica</i> Rayss [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora sessilis</i> Sorokīn	Not known to		Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Capnodiales: Mycosphaerellaceae]	occur	fungi.			
<i>Cercospora truncata</i> Ellis & Everh. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora truncatella</i> G.F. Atk. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora vitiphylla</i> (Speschnew) Barbarin. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora vitis-heterophyllae</i> Hennings. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercospora vulpinae</i> Ellis & Kellerm [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Cercosporidium vitis</i> MS Patil & Sawant [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, <i>Cercosporidium</i> species are associated with foliage and cause late leaf spot in host plants (Meena 2010). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Chaetospermum chaetosporum</i> (Pat.) AL Smith & Ramsb. [Unassigned]	Yes (PHA 2001)	Assessment not required			
<i>Chalara ampullula</i> (Sacc.) Sacc. [Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, <i>Chalara</i>	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		species are associated with wood and dead leaves in other host plants (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Cladosporium asperulatum</i> Bensch <i>et al.</i> [Capnodiales: Davidiellaceae]	Not known to occur	<b>Yes:</b> <i>Cladosporium</i> species are saprobic on dead plant material (Farr <i>et al.</i> 1989), are associated with foliage and cause leaf spot (Pearson and Goheen 1988; Farr and Rossman 2011) or are associated with canes (Pearson and Goheen 1988). Therefore, dormant cuttings may provide a pathway for these fungi.	<b>Yes.</b> These fungi have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and may spread naturally in infected propagative material. Therefore, these fungi have the potential for establishment and spread in Australia.	No: <i>Cladosporium</i> species usually cause minor foliage diseases or fruit rot in storage (Pearson and Goheen 1988). The two <i>Cladosporium</i> species ( <i>C. cladosporioides</i> and <i>C. herbarum</i> ) that are present in Australia are associated with berry rot causing yield losses and reducing wine quality (Briceño and Latorre 2008). <i>Cladosporium asperulatum</i> , <i>C. autumnale</i> and <i>C. baccae</i> are not associated with berry rot (Briceño and Latorre 2008) and are therefore not economically important.	
<i>Cladosporium autumnale</i> Kübler [Capnodiales: Davidiellaceae]	Not known to occur				
<i>Cladosporium baccae</i> Verwoerd & Dippen. [Capnodiales: Davidiellaceae]	Not known to occur				
<i>Cladosporium cladosporioides</i> (Fresen.) G.A. de Vries [Capnodiales: Davidiellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Cladosporium fasciculatum</i> Corda [Capnodiales: Davidiellaceae]	Not known to occur	<b>Yes:</b> <i>Cladosporium</i> species are saprobic on dead plant material (Farr <i>et al.</i> 1989), are associated with foliage and cause leaf spot (Pearson and Goheen 1988; Farr and Rossman 2011) or are associated with canes	<b>Yes.</b> <i>Cladosporium</i> species have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and may spread naturally in infected propagative material. Therefore, these fungi have	No: <i>Cladosporium</i> species cause minor foliage diseases or fruit rot in storage (Pearson and Goheen 1988). The two <i>Cladosporium</i> species ( <i>C. cladosporioides</i> and <i>C. herbarum</i> ) associated with berry rot, causing yield losses and reducing wine quality	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Pearson and Goheen 1988). Therefore, dormant cuttings may provide a pathway for this fungus.	the potential for establishment and spread in Australia.	(Briceño and Latorre 2008), are present in Australia. <i>C. fasciculatum</i> is not associated with berry rot (Briceño and Latorre 2008) and is therefore not economically important.	
<i>Cladosporium herbarum</i> (Pers.) Link [Capnodiales: Davidiellaceae]	Yes (Cook and Dubé 1989)	Assessment not required			
<i>Cladosporium longipes</i> Sorokín [Capnodiales: Davidiellaceae]	No records found	<b>Yes:</b> <i>Cladosporium</i> species are saprobic on dead plant material (Farr <i>et al.</i> 1989), are associated with foliage and cause leaf spot (Pearson and Goheen 1988; Farr and Rossman 2011) or are associated with canes (Pearson and Goheen 1988). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes.</b> <i>Cladosporium</i> species have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and may spread naturally in infected propagative material. Therefore, these fungi have the potential for establishment and spread in Australia.	No: <i>Cladosporium</i> species cause minor foliage diseases or fruit rot in storage (Pearson and Goheen 1988). The two <i>Cladosporium</i> species ( <i>C. cladosporioides</i> and <i>C. herbarum</i> ) associated with berry rot, causing yield losses and reducing wine quality (Briceño and Latorre 2008), are present in Australia. <i>C. longipes</i> is not associated with berry rot (Briceño and Latorre 2008) and is therefore not economically important.	
<i>Cladosporium macrocarpum</i> Preuss [Capnodiales: Davidiellaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Cladosporium oxysporum</i> Berk. & M.A. Curtis [Capnodiales: Davidiellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Cladosporium roesleri</i> Catt. [Capnodiales: Davidiellaceae]	No records found	<b>Yes:</b> <i>Cladosporium</i> species are saprobic on dead plant material (Farr <i>et al.</i> 1989), are associated with foliage and cause leaf spot (Pearson and Goheen 1988; Farr and Rossman 2011) or are associated with canes	<b>Yes.</b> <i>Cladosporium</i> species have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and may spread naturally in infected propagative material. Therefore, these fungi have	No: <i>Cladosporium</i> species cause minor foliage diseases or fruit rot in storage (Pearson and Goheen 1988). The two <i>Cladosporium</i> species ( <i>C. cladosporioides</i> and <i>C. herbarum</i> ) associated with berry rot, causing yield losses and reducing wine quality	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Pearson and Goheen 1988). Therefore, dormant cuttings may provide a pathway for this fungus.	the potential for establishment and spread in Australia.	(Briceño and Latorre 2008), are present in Australia. <i>C. roesleri</i> is not associated with berry rot (Briceño and Latorre 2008) and is therefore not economically important.	
<i>Cladosporium tenuissimum</i> Cooke [Capnodiales: Davidiellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Cladosporium uvarum</i> McAlpine [Capnodiales: Davidiellaceae]	Yes (Dugan <i>et al.</i> 2004)	Assessment not required			
<i>Cladosporium viticola</i> Ces. [Capnodiales: Davidiellaceae]	No records found	<b>Yes:</b> <i>Cladosporium</i> species are saprobic on dead plant material (Farr <i>et al.</i> 1989), are associated with foliage and cause leaf spot (Pearson and Goheen 1988; Farr and Rossman 2011) or are associated with canes (Pearson and Goheen 1988). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes.</b> <i>Cladosporium</i> species have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and may spread naturally in infected propagative material. Therefore, these fungi have the potential for establishment and spread in Australia.	No: <i>Cladosporium</i> species cause minor foliage diseases or fruit rot in storage (Pearson and Goheen 1988). The two <i>Cladosporium</i> species ( <i>C. cladosporioides</i> and <i>C. herbarum</i> ) associated with berry rot, causing yield losses and reducing wine quality (Briceño and Latorre 2008), are present in Australia. <i>C. viticola</i> is not associated with berry rot (Briceño and Latorre 2008) and is therefore not economically important.	
<i>Clathrospora turkestanica</i> Domashova [Pleosporales: Pleosporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011) but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species from Central Asia in 1973 (Farr and Rossman 2011), it has not been reported from any other country, indicating that	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		grapevine propagative material does not provide a pathway for this fungus.			
<i>Cochliobolus bicolor</i> AR Paul & Parbery [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Cochliobolus geniculatus</i> RR Nelson [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Colletotrichum acutatum</i> J.H. Simmonds [Incertae sedis: Glomerellaceae]	Yes (Whitelaw-Weckert <i>et al.</i> 2007a)	Assessment not required			
<i>Colletotrichum ampelinum</i> Cavara [Incertae sedis: Glomerellaceae]	Not known to occur	No: <i>Colletotrichum</i> species are foliar pathogens (Mohan 2005). This fungus is associated with grapevines, causing anthracnose in China (Anon 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Colletotrichum capsici</i> (Syd. & P. Syd.) E.J. Butler & Bisby [Incertae sedis: Glomerellaceae]	Yes (Shivas 1989)	Assessment not required			
<i>Colletotrichum crassipes</i> (Speg.) Arx [Incertae sedis: Glomerellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Colletotrichum gloeosporioides</i> (Penz.) Penz. & Sacc. [Incertae sedis: Glomerellaceae]	Yes (Shivas 1989)	Assessment not required			
<i>Colletotrichum simmondsii</i> RG Shivas & YP Tan [Incertae sedis: Glomerellaceae]	Yes (Shivas and Tan 2009)	Assessment not required			
<i>Coniella castaneicola</i> (Ellis & Everh.) B. Sutton [Diaporthales: Schizoparmaceae]	Yes (PHA 2001)	Assessment not required			
<i>Coniella diplodiella</i> (Speg.) Petr. & Syd. [Diaporthales: Schizoparmaceae]	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Coniella fragariae</i> (Oudem.) B. Sutton [Diaporthales: Schizoparmaceae]	Yes (PHA 2001)	Assessment not required			
<i>Coniella granati</i> (Sacc.) Petr. & Syd. [Diaporthales: Schizoparmaceae]	Yes (PHA 2001)	Assessment not required			
<i>Coniella petrakii</i> B. Sutton [Diaporthales: Schizoparmaceae]	Not known to occur	No: This soil-borne fungus has been recorded on <i>Vitis</i> species, causing white root rot (König <i>et al.</i> 2009). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Corticium cremeoalbidum</i> (MJ Larsen & Nakasone) MJ Larsen [Corticiales: Corticiaceae] (synonym: <i>Laeticorticium cremeoalbidum</i> MJ Larsen & Nakasone)	Not known to occur	No: This fungus has been recorded on the wood of <i>Vitis</i> species (CABI 2012b). There is no evidence that this species occurs on the young stems of grapevines. Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Coryneopsis microsticta</i> Grove [Xylariales: Amphisphaeriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Corynespora calicioidea</i> (Berk. & Broome) M.B. Ellis [Pleosporales: Corynesporascaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, other <i>Corynespora</i> species are associated with wood, bark and dead wood (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for these	Assessment not required		
<i>Corynespora karnati</i> (VG Rao) MB Ellis. [Pleosporales: Corynesporascaceae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		fungi.			
<i>Coryneum discolor</i> Fautrey [Diaporthales: Pseudovalsaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, <i>Coryneum</i> species occur on twigs and foliage of other hosts (Schloemann 2003-2004; Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Coryneum microstictum</i> Berk. & Broome [Diaporthales: Pseudovalsaceae]	Not known to occur		Assessment not required		
<i>Coryneum vitiphyllum</i> Speschnew [Diaporthales: Pseudovalsaceae]	Not known to occur		Assessment not required		
<i>Crepidotus amarus</i> Murrill [Agaricales: Inocybaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, <i>Crepidotus</i> species occur on bark and wood of hardwoods on other hosts (Farr and Rossman 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Cryptophaeella trematosphaeriicola</i> Frolov [Pleosporales: Montagnulaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species from Central Asia in 1973 (Farr and Rossman 2011), it has not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		been reported from any other country. This indicates that dormant cuttings do not provide a pathway for this fungus.			
<i>Cryptosphaeria pullmanensis</i> Glawe [Xylariales: Diatrypaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species and has been isolated from cankered wood (Trouillas and Gubler 2010). Generally, <i>Cryptosphaeria</i> species occur on bark of host plants (Romero and Carmaran 2003). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Cryptostictis hysterioides</i> Fuckel [Xylariales: Amphisphaeriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Cryptostictis inaequalis</i> Tehon & Stout [Xylariales: Amphisphaeriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus occurs on the leaves (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Cryptovalsa ampelina</i> Fuckel [Incertae sedis: Incertae sedis]	Yes (Sosnowski <i>et al.</i> 2007)	Assessment not required			
<i>Cryptovalsa protracta</i> (Pers.) De Not. [Unassigned]	Yes (Yuan 1996)	Assessment not required			



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Cryptovalsa rabenhorstii</i> (Nitschke) Sacc. [Unassigned]	Yes (Trouillas <i>et al.</i> 2011)	Assessment not required			
<i>Cylindrocarpon destructans</i> var. <i>destructans</i> (Zinssm.) Scholten [Hypocreales: Nectriaceae]	Yes (Sweetingham 1983)	Assessment not required			
<i>Cylindrocarpon lichenicola</i> (C. Massal.) D. Hawksw. [Hypocreales: Nectriaceae]	Yes (Brayford 1987)	Assessment not required			
<i>Cylindrocarpon liriodendri</i> JD MacDonald & EE Butler [Hypocreales: Nectriaceae]	Yes (Whitelaw-Weckert <i>et al.</i> 2007b)	Assessment not required			
<i>Cylindrocarpon macrodidymum</i> Schroers <i>et al.</i> [Hypocreales: Nectriaceae]	Yes (Whitelaw-Weckert 2008)	Assessment not required			
<i>Cylindrocarpon obtusisporum</i> (Cooke & Harkn.) Wollenw [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Cylindrocarpon pauciseptatum</i> Schroers & Crous [Hypocreales: Nectriaceae]	Not known to occur	No: This species is associated with black foot disease in grapevines (Schroers <i>et al.</i> 2008; Martin <i>et al.</i> 2011a). This fungus has reported to occur in the roots (Alaniz <i>et al.</i> 2007), stem vascular tissue and brown wood of young grapevines (Martin <i>et al.</i> 2011a). Semi-hardwood, root free dormant cuttings therefore do not provide a pathway for this fungus.	Assessment not required		
<i>Cylindrocladiella lageniformis</i> Crous <i>et al.</i> [Hypocreales: Nectriaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but	Assessment not required		
<i>Cylindrocladiella parva</i> (P.J. Anderson)	Not known to		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Boesew [Hypocreales: Nectriaceae]	occur	affected plant parts are not mentioned. However, on other hosts these fungi occur on the roots (Van-Coller <i>et al.</i> 2005). Therefore, root free dormant cuttings do not provide a pathway for these fungi.			
<i>Cylindrocladiella peruviana</i> (Bat. <i>et al.</i> ) Boesew [Hypocreales: Nectriaceae]	Not known to occur		Assessment not required		
<i>Daldinia concentrica</i> sensu auct. [Xylariales: Xylariaceae]	Yes (PHA 2001)	Assessment not required			
<i>Daldinia vernicosa</i> (Schwein.) Ces. & De Not. [Xylariales: Xylariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts, this fungus occurs on burnt wood (Rhoads 1918; Whalley and Watling 1980; Farr and Rossman 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Dendrophoma pleurospora</i> Sacc [Xylariales: Xylariaceae] (synonym: <i>Dinemasporium pluerospora</i> (Sacc.) Shkarupa)	Not known to occur	No: This fungus is associated with <i>Vitis</i> species and has been isolated from the necrotic and healthy stem tissue of older grapevines (Serra <i>et al.</i> 2000). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Dendryphiella infuscans</i> (Thüm.) M.B. Ellis [Pleosporales: Pleosporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		affected plant parts are not mentioned. However, on other hosts this fungus occurs on dead stems (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Dendryphion acinorum</i> Ellis & Everh. [Pleosporales: Pleosporaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since their report on <i>Vitis</i> species in the United States in 1952 (Farr and Rossman 2011), they have not been reported from any other country indicating that dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Dendryphion harknessii</i> var. <i>leptaleum</i> Ellis [Pleosporales: Pleosporaceae]	Not known to occur		Assessment not required		
<i>Dermatella viticola</i> Ellis & Everh. [Helotiales: Dermateaceae]	Not known to occur	No: This species is associated with dead shoots (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Diaporthe australafricana</i> Crous & Niekirk [Diaporthales: Diaporthaceae]	Yes (Van Niekirk <i>et al.</i> 2005)	Assessment not required			
<i>Diaporthe eres</i> Nitschke [Diaporthales: Diaporthaceae] (synonym <i>Diaporthe ambigua</i> Nitschke)	Yes (PHA 2001)	Assessment not required			
<i>Diaporthe kyushuensis</i> Kajitani & Kanem [Diaporthales: Diaporthaceae]	Not known to occur	<b>Yes:</b> This fungus is associated with <i>Vitis</i> species,	<b>Yes:</b> This fungus has established in areas with a	No: This species has been reported on grapes, causing	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		causing leaf and cane spot (Kajitani and Kanematsu 2000). Small black spots appear at the base of the green shoot which later coalesces to form a blackened zone (Kajitani and Kanematsu 2000). Infection may also be latent (Kajitani and Kanematsu 2000). Therefore, propagative material may provide a pathway for this fungus.	wide range of climatic conditions and it can spread naturally in infected propagative material (Kajitani and Kanematsu 2000). Propagation and distribution of infected material will help spread this fungus within Australia. Therefore, this fungus has the potential to establish and spread in Australia.	canker in the 1960s in Japan (Kajitani and Kanematsu 2000). Since then, no economic losses have been reported. Therefore, this fungus is not of economic concern for host plants.	
<i>Diaporthe medusaea</i> Nitschke [Diaporthales: Diaporthaceae]	Yes (PHA 2001)	Assessment not required			
<i>Diaporthe perijuncta</i> Niessl. [Diaporthales: Diaporthaceae]	Yes (Van Niekerk <i>et al.</i> 2005)	Assessment not required			
<i>Diaporthe viticola</i> Nitschke [Diaporthales: Diaporthaceae]	Yes (Scheper <i>et al.</i> 2000)	Assessment not required			
<i>Diatrype nigerrima</i> Ellis & Everh. [Xylariales: Diatrypaceae]	Not known to occur	No: Most species of diatrypaceous fungi have been regarded as saprobes (Glawe and Rogers 1984). Species in the Diatrypaceae family have been isolated from the cankered wood of grapevines (Trouillas and Gubler 2010). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Diatrype oregonensis</i> (Wehm.) Rappaz [Xylariales: Diatrypaceae]	Not known to occur		Assessment not required		
<i>Diatrype stigma</i> (Hoffm.) Fr. [Xylariales: Diatrypaceae]	Not known to occur		Assessment not required		
<i>Diatrype vitis</i> Ellis & Everh. [Xylariales: Diatrypaceae]	Not known to occur		Assessment not required		
<i>Diatrype whitmanensis</i> J.D. Rogers & Glawe [Xylariales: Diatrypaceae]	Not known to occur		Assessment not required		
<i>Diatrypella verruciformis</i> (Ehrh.) Fr. [Xylariales: Diatrypaceae]	Not known to occur	No: Species in the Diatrypaceae family have been isolated from the	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		cankered wood of grapevines (Trouillas and Gubler 2010). Pathogenicity studies indicate that this species is saprophytic rather than pathogenic on grapes (Trouillas and Gubler 2010). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Diatrypella vulgaris</i> Trouillas <i>et al.</i> [Xylariales: Diatrypaceae]	Yes (Trouillas <i>et al.</i> 2011)	Assessment not required			
<i>Dichomera viticola</i> Cooke & Harkn. [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur	No: This species occurs on the dead stems of the plant (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Dictyosporium elegans</i> Corda [Pleosporales: Unassigned]	Yes (PHA 2001)	Assessment not required			
<i>Didymosphaeria sarmenti</i> (Cooke & Harkness) Berl. & Voglino [Pleosporales: Didymosphaeriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, the majority of <i>Didymosphaeria</i> species are saprobes that grow mostly on dead plant material (Aptroot 1995). Therefore, dormant cuttings are unlikely to provide a pathway for this fungus.	Assessment not required		
<i>Diplodia ampelina</i> (Cooke ) [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur	<b>Yes:</b> These species have been isolated from cankered	<b>Yes:</b> These species occur in a wide range of climates (Farr	No: There is no evidence that these species cause significant	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Diplodia porosum</i> Van Niekerk & Crous [Botryosphaerales: Botryosphaeriaceae] (synonym: <i>Phaeobotryosphaeria porosa</i> (Van Niekerk & Crous) Crous & A.J.L. Phillips)	Not known to occur	grapevines (Phillips <i>et al.</i> 2008; Úrbez-Torres and Gubler 2009). These species are endophytic (Paoletti <i>et al.</i> 2007) and both saprophytic and pathogenic (Úrbez-Torres 2011) and have been isolated from the shoots of grapevines (Aroca <i>et al.</i> 2010). Therefore, dormant cuttings may provide a pathway for these fungi.	and Rossman 2011). Therefore, parts of Australia will be suitable for the establishment and spread of these species. Distribution of infected propagative material will assist the establishment and spread of these fungi in Australia.	economic consequences. Therefore, these species do not have the potential for economic consequences in Australia.	
<i>Diplodia seriata</i> De Not [Botryosphaerales: Botryosphaeriaceae]	Yes (Pitt <i>et al.</i> 2009)	Assessment not required			
<i>Diplodina vitis</i> Brunaud [Diaporthales: Gnomoniaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species in Central Asia in 1973 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Discohainesia oenotherae</i> (Cooke & Ellis) Nannf [Unassigned]	Yes (PHA 2001)	Assessment not required			
<i>Discosia artocreas</i> (Tode) Fr. [Xylariales: Amphispheeriaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts <i>Discosia</i> species occur on leaves (Farr and Rossman	Assessment not required		
<i>Discosia vitis</i> Schulzer [Xylariales: Amphispheeriaceae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		2011). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.			
<i>Discostroma corticola</i> (Fuckel) Brockmann [Xylariales: Amphisphaeriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Dothidella confluens</i> (Welw. & Curr.) Sacc. [Incertae sedis: Polystomellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus is associated with foliage and causes leaf spot (Chee 1976). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Dothiorella americana</i> Urbez-Torres <i>et al.</i> sp. nov. [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur	<b>Yes:</b> This fungus is associated with die-back of <i>Vitis</i> species and has been isolated from grapevine vascular tissue (Urbez-Torres <i>et al.</i> 2012). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Urbez-Torres <i>et al.</i> 2012). Propagation and distribution of infected material will help spread this fungus within Australia. Therefore, this fungus has the potential to establish and spread in Australia.	No: Although this fungus is associated with die-back, it is considered a weak pathogen of grapevines (Urbez-Torres <i>et al.</i> 2012). Other <i>Dothiorella</i> species are also generally considered weak pathogens of grapevines (Urbez-Torres <i>et al.</i> 2006; Urbez-Torres and Gubler 2009). This <i>Dothiorella</i> species has not been recorded to have economic consequences. Therefore, this fungus is not of economic concern to Australia.	
<i>Dothiorella iberica</i> Phillips <i>et al.</i>	Yes (Wunderlich)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Botryosphaerales: Botryosphaeriaceae]	<i>et al.</i> 2008, Pitt <i>et al.</i> 2009)				
<i>Dothiorella sarmentorum</i> (Fr.) Phillips <i>et al.</i> [Botryosphaerales: Botryosphaeriaceae] (synonym <i>Diplodia sarmentorum</i> (Fr.) Fr.)	Not known to occur	<b>Yes:</b> This species has been isolated from the trunks of grapevines (Gramaje <i>et al.</i> 2009b). <i>Dothiorella</i> species have been isolated from the vascular tissue of grapevines (Urbez-Torres <i>et al.</i> 2012). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This species is distributed across a wide range of climates (Gramaje <i>et al.</i> 2009b; Farr and Rossman 2011). Parts of Australia have suitable climatic conditions for the establishment and spread of this species. Propagation and distribution of infected material will help spread this fungus within Australia.	No: This species occurs on a range of hosts, including elms, grapevines, <i>Malus</i> species and <i>Prunus</i> species (Phillips <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009b; Gramaje <i>et al.</i> 2012). <i>Dothiorella</i> species are generally considered weak pathogens of grapevines (Urbez-Torres <i>et al.</i> 2006; Urbez-Torres and Gubler 2009). This <i>Dothiorella</i> species has not been recorded to have significant economic consequences. Therefore, this fungus is not of economic concern to Australia.	
<i>Drechslera tetramera</i> (McKinney) Subram. & B.L. Jain [Pleosporales: Pleosporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus is associated with roots (Nan 1995). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Ellisembia brachypus</i> (Ellis & Everh.) Subram. [Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		hosts this species occurs on dead wood (Sivichai <i>et al.</i> 2000). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Elsinoë ampelina</i> Shear [Myriangiales: Elsinoaceae]	Yes (Magarey <i>et al.</i> 1993)	Assessment not required			
<i>Endothia radicalis</i> (Schwein.) De Not. [Diaporthales: Cryphonectriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, this species is saprophytic and occurs in dead stems on other hosts (Hoegger <i>et al.</i> 2002). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Epicoccum nigrum</i> Link [Pleosporales: Pleosporaceae] (synonym: <i>Epicoccum granulatum</i> Penz.)	Yes (PHA 2001)	Assessment not required			
<i>Eriosphaeria oenotria</i> Sacc. & Speg. [Trichosphaeriales: Trichosphaeriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species in Italy in 1973 (Farr and Rossman 2011), it has not been reported from any other country, indicating that dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Erysiphe necator</i> Schwein. [Erysiphales:	Yes (Magarey <i>et</i>	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Erysiphaceae]	<i>al.</i> 1997)				
<i>Eutypa lata</i> (Pers.) Tul. & C. Tul [Xylariales: Diatrypaceae]	Yes (Constable and Drew 2004)	Assessment not required			
<i>Eutypa leptoplaca</i> (Mont.) Rappaz [Xylariales: Diatrypaceae]	Yes (Trouillas <i>et al.</i> 2010)	Assessment not required			
<i>Eutypa ludibunda</i> Sacc. [Xylariales: Diatrypaceae]	Not known to occur	No: This fungus occurs on the dead wood of host plants (Rolshausen 2004). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Eutypella citricola</i> Speg [Xylariales: Diatrypaceae]	Yes (Trouillas <i>et al.</i> 2011)	Assessment not required			
<i>Eutypella fraxinicola</i> (Cooke & Peck) Sacc. [Xylariales: Diatrypaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this fungus occurs on dead branches of <i>Fraxinus</i> and <i>Ulmus</i> species (Vasilyeva and Stephenson 2006; Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Eutypella leprosa</i> (Pers.) Berl. [Xylariales: Diatrypaceae]	Not known to occur	<b>Yes:</b> This fungus has been isolated from grapevines showing canker symptoms and vascular necrosis of trunks, arms and spurs (Diaz <i>et al.</i> 2011). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Diaz <i>et al.</i> 2011; Farr and Rossman 2011) and it may spread naturally in infected propagative material. Therefore, this fungus has the	<b>Yes:</b> This fungus causes cankers and vascular necrosis of trunks, arms and spurs, along with general decline and dieback of grapevines (Diaz <i>et al.</i> 2011). Therefore, this fungus has potential for economic consequences in Australia.	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
			potential to establish and spread in Australia.		
<i>Eutypella microtheca</i> Trouillas <i>et al.</i> [Xylariales: Diatrypaceae]	Yes (Trouillas <i>et al.</i> 2011)	Assessment not required			
<i>Eutypella vitis</i> (Schwein.Fr.) Ellis & Everhart [Xylariales: Diatrypaceae] (synonym: <i>Eutypella aequilinearis</i> (Schwein. Fr.) Starb.)	Not known to occur	<b>Yes:</b> This fungus is associated with Eutypa dieback and has been isolated from the trunks and branches of <i>Vitis</i> species (Catal <i>et al.</i> 2007). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Catal <i>et al.</i> 2007; Farr and Rossman 2011) and it may spread naturally in infected propagative material. Therefore, this fungus has the potential to establish and spread in Australia.	<b>Yes:</b> <i>Eutypella vitis</i> has been identified as an additional causal agent of Eutypa dieback, an important disease of grapevine (Navarrete <i>et al.</i> 2010). Therefore, this fungus has potential for economic consequences in Australia.	<b>Yes</b>
<i>Exosporium sultanae</i> Du Plessis [Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, <i>Exosporium</i> species occur on the leaves of other hosts (Pitta 1994). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Favolus tenuiculus</i> P. Beauv. [Polyporales: Polyporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this saprobic mushroom species occurs on decaying hardwood (Ruan-Soto <i>et al.</i> 2006). Therefore, semi-hardwood	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		dormant cuttings do not provide a pathway for this fungus.			
<i>Fomes fomentarius</i> (L.) J. Kickx [Polyporales: Polyporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this mushroom species occurs on decaying hardwood (Monthey and Cross 2000). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Fomitiporia australiensis</i> Fischer <i>et al.</i> [Hymenochaetales: Hymenochaetaceae]	Yes (Pascoe <i>et al.</i> 2005)	Assessment not required			
<i>Fomitiporia mediterranea</i> M. Fischer <sup>12</sup> [Hymenochaetales: Hymenochaetaceae]	Not known to occur	<b>Yes:</b> <i>Fomitiporia</i> species are associated with wood decay of grapevines showing esca symptoms (Cortesi <i>et al.</i> 2000; Sparapano <i>et al.</i> 2000; Ciccarone <i>et al.</i> 2004; Fischer 2006; Amalfi <i>et al.</i> 2010). <i>Fomitiporia</i> species cause spongy wood decay in the trunks of growing <i>Vitis</i> plants (Sparapano <i>et al.</i> 2000; Amalfi <i>et al.</i> 2010). Therefore, dormant cuttings may provide a pathway for these fungi.	<b>Yes:</b> These fungi have been established in areas with a wide range of climatic conditions (Cortesi <i>et al.</i> 2000; Sparapano <i>et al.</i> 2000; Ciccarone <i>et al.</i> 2004; Fischer 2006; Amalfi <i>et al.</i> 2010) and may spread naturally in infected propagative material. Propagation and distribution of infected material will help spread these fungi within Australia. Therefore, these fungi have the potential to	<b>Yes:</b> <i>Fomitiporia</i> species constitute the complex of pathogens associated with the diseases forming the esca complex (Abou-Mansour <i>et al.</i> 2009). Esca is a complex trunk disease including a vascular disease and an internal white rot of the trunk, which gradually changes the hard wood to a soft, friable, spongy mass (Graniti <i>et al.</i> 1994; Mugnai <i>et al.</i> 1999). Grapevine trunk diseases cause a slow decline and yield loss in	<b>Yes</b>
<i>Fomitiporia polymorpha</i> M. Fisch. [Hymenochaetales: Hymenochaetaceae]	Not known to occur				<b>Yes</b>

<sup>12</sup> *Fomitiporia punctata* has been mentioned in literature as being associated with grapevines however, these records of *Fomitiporia punctata* on grapevine have more recently been attributed to *Fomitiporia mediterranea* (Fischer 2002); grapevine is no longer considered a host of this species.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
			establish and spread in Australia.	grapevines at all stages of growth, the death of spurs, arms, and cordons, and the eventual death of the vines due to a progressive wood necrosis and decay of plant tissue (Andolfi <i>et al.</i> 2011). Therefore, these fungi have potential for economic consequences in parts of Australia.	
<i>Fusarium acuminatum</i> Ellis & Everh [Hypocreales: Nectriaceae]	Yes (Wong <i>et al.</i> 1985)	Assessment not required			
<i>Fusarium anthophilum</i> (A. Braun) Wollenw. [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Fusarium avenaceum</i> (Fr.) Sacc. [Hypocreales: Nectriaceae]	Yes (Summerell <i>et al.</i> 2011)	Assessment not required			
<i>Fusarium culmorum</i> (W.G. Sm.) Sacc. [Hypocreales: Nectriaceae]	Yes (Summerell <i>et al.</i> 2011)	Assessment not required			
<i>Fusarium equiseti</i> (Corda) Sacc. [Hypocreales: Nectriaceae]	Yes (Wong <i>et al.</i> 1985)	Assessment not required			
<i>Fusarium moniliforme</i> J. Sheld. [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Fusarium oxysporum</i> f. sp. <i>herbemontis</i> (Tochetto) W.L. Gordon [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus is associated with grapevines causing Fusarium wilt (de Andrade <i>et al.</i> 1995). It occurs in the root vascular system of the plant, causing vascular root discolouration (Gallotti 1991). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Fusarium oxysporum</i> Schtdl.	Yes (Summerell	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Hypocreales: Nectriaceae]	<i>et al.</i> 2011)				
<i>Fusarium poae</i> (Peck) Wollenw. [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Fusarium proliferatum</i> (Matsush.) Nirenberg ex Gerlach & Nirenberg [Hypocreales: Nectriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Fusarium schweinitzii</i> Ell. & Hark. [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on the dead wood of <i>Vitis</i> species (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Fusarium solani</i> (Mart.) Sacc. [Hypocreales: Nectriaceae]	Yes (Summerell <i>et al.</i> 2011)	Assessment not required			
<i>Fusarium sporotrichioides</i> Sherb. [Hypocreales: Nectriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Fusarium subglutinans</i> (Wollenw. & Reinking) Nelson <i>et al.</i> [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Fusarium volutella</i> Ellis & Everh. [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011). <i>Fusarium</i> species are soil-borne, causing root rot (Lew <i>et al.</i> 1996; Farr and Rossman 2011). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Fusicladium viticis</i> M.B. Ellis [Pleosporales: Venturiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		affected plant parts are not mentioned. Generally, <i>Fusicladium</i> species occur on foliage (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Fusicoccum macroclavatum</i> Burgess <i>et al.</i> [Botryosphaerales: Botryosphaeriaceae] (synonym: <i>Neofusicoccum macroclavatum</i> (Burgess <i>et al.</i> ) Burgess <i>et al.</i> )	Yes (Burgess <i>et al.</i> 2005)	Assessment not required			
<i>Fusicoccum viticlavatum</i> Niekirk & Crous [Botryosphaerales: Botryosphaeriaceae] (synonym: <i>Neofusicoccum viticlavatum</i> (Van Niekirk & Crous) Crous <i>et al.</i> )	Not known to occur	<b>Yes:</b> These fungi have been recorded on <i>Vitis</i> species causing brown wood streaking and internal necrotic lesions (Van Niekirk <i>et al.</i> 2004). Therefore, dormant cuttings may provide a pathway for these fungi.	<b>Yes.</b> These fungi have established in areas with a wide range of climatic conditions (Van Niekirk <i>et al.</i> 2004) and may spread naturally in infected propagative material. Therefore, these fungi have the potential for establishment and spread in Australia.	No: These species have been recorded on grapevines causing canker in association with other species (Van Niekirk <i>et al.</i> 2004). However, no information is available on the losses caused by these pathogens. Therefore, these fungi are not of economic concern for host plants.	
<i>Fusicoccum vitifusiforme</i> Niekirk & Crous [Botryosphaerales: Botryosphaeriaceae] (synonym: <i>Neofusicoccum vitifusiforme</i> (Van Niekirk & Crous) Crous <i>et al.</i> )	Not known to occur				
<i>Gliocladium roseum</i> Bainier [Hypocreales: Hypocreaceae] (synonym <i>Clonostachys rosea</i> (Link) Schroers <i>et al.</i> )	Yes (PHA 2001)	Assessment not required			
<i>Gloeosporium sarmenticola</i> Speg. [Helotiales: Dermateaceae]	No records found	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species in Argentina in 1973 (Farr and Rossman	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.			
<i>Glonium clavisporum</i> Seaver [Hysteriales: Hysteriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts <i>Glonium</i> species occur on bark and dead wood (Farr <i>et al.</i> 1989; Farr and Rossman 2011). Therefore, dormant grapevine cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Greeneria uvicola</i> (Berkley & M.A. Curtis) Punithalingam [Diaporthales: Unassigned]	Yes (Castillo-Pando <i>et al.</i> 1999; Sergeeva <i>et al.</i> 2001)	Assessment not required			
<i>Grovesinia pyramidalis</i> M.N. Cline, J.L. Crane & S.D. Cline [Helotiales: Sclerotiniaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus causes leaf spot (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz [Botryosphaerales: Botryosphaeriaceae]	Not known to occur	<b>Yes:</b> These fungi are associated with the foliage,	<b>Yes:</b> These fungi have established in areas with a	<b>Yes.</b> These fungi cause black rot, an important disease of	<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
(synonym: <i>Greenaria uvicola</i> (Berk. & M.A. Curtis) Punith.)		shoots, tendrils, cluster stems and fruit of <i>Vitis</i> species (University of Illinois 2001; Ellis 2008; Ullrich <i>et al.</i> 2009).	wide range of climatic conditions (Farr and Rossman 2011) and can spread naturally in infected propagative material.	grapevine that affects the foliage, petioles, shoots, tendrils, cluster stems and fruit (University of Illinois 2001; Ellis 2008; Ullrich <i>et al.</i> 2009). These fungi can cause substantial economic losses (Ramsdell and Milholland 1988; Wilcox 2003). For instance, crop loss due to black rot can range from 5–100% (Kummuang <i>et al.</i> 1996; Eyres <i>et al.</i> 2006). Therefore, these fungi have the potential for significant economic consequences in parts of Australia.	
<i>Guignardia bidwellii</i> f. <i>euvitis</i> Luttrell [Botryosphaerales: Botryosphaeriaceae]	Not known to occur	These fungi overwinter in infected canes, tendrils, fallen leaves and in mummified fruit on vines or on the ground (Kummuang <i>et al.</i> 1996; Ellis 2008). Therefore, dormant grapevine cuttings may provide a pathway for these fungi.	Multiplication and marketing of infected propagative material will help spread these fungi within Australia. Ascospores and conidia are the primary inoculum and are spread by air and rain (Pearson and Goheen 1988). Therefore, these fungi have the potential to establish and spread in Australia.		Yes
<i>Guignardia bidwellii</i> f. <i>muscadinii</i> Luttrell [Botryosphaerales: Botryosphaeriaceae]	Not known to occur				Yes
<i>Hapalopilus nidulans</i> (Fr.) P. Karst. [Polyporales: Polyporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this species grows on decaying logs, sticks or hardwood debris and causes white rot (Kuo 2003). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Helicobasidium mompa</i> Nobuj. Tanaka [Helicobasidiales: Helicobasidiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, this species infects the below	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		ground parts of a variety of other host plants (Matsubara <i>et al.</i> 2000). Therefore, root free dormant cuttings do not provide a pathway for this fungus.			
<i>Helminthosporium decacuminatum</i> Thüm. & Pass. [Pleosporales: Massarinaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts these fungi occur on dead and dying plant material (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Helminthosporium siliquosum</i> Berk. & MA Curtis [Pleosporales: Massarinaceae]	Not known to occur		Assessment not required		
<i>Helminthosporium velutinum</i> Link [Pleosporales: Massarinaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Helotium sarmentorum</i> De Not [Helotiales: Helotiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its report on <i>Vitis</i> species in Portugal in 1941 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Hendersonia cookeana</i> Speg. [Pleosporales: Phaeosphaeriaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but	Assessment not required		
<i>Hendersonia corticalis</i> Ellis & Everhart	Not known to	(Farr and Rossman 2011), but	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Pleosporales: Phaeosphaeriaceae]	occur	affected plant parts are not mentioned. Species of the genus are foliar pathogens (Sinclair <i>et al.</i> 1987; Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.			
<i>Hendersonia sarmentorum</i> Westend. [Pleosporales: Phaeosphaeriaceae]	Not known to occur		Assessment not required		
<i>Hendersonia tenuipes</i> McAlpine [Pleosporales: Phaeosphaeriaceae]	Not known to occur		Assessment not required		
<i>Hendersonia viticola</i> S. Ahmad [Pleosporales: Phaeosphaeriaceae]	Not known to occur		Assessment not required		
<i>Hinomyces moricola</i> (I. Hino) Narumi & Y. Harada [Helotiales: Sclerotiniaceae]	Not known to occur	No. This species has been recorded on <i>Vitis</i> species causing leaf spot (Li 2004). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Hyphodontia pruni</i> (Lasch) Svrček [Hymenochaetales: Schizoporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts this fungus causes wood rot (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Hypocrea gelatinosa</i> (Tode) Fr. [Hypocreales: Hypocreaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, this species occurs on dead wood and other decaying matter on other hosts (Farr <i>et al.</i> 1989). Therefore, dormant cuttings	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		do not provide a pathway for this fungus.			
<i>Hypoderma commune</i> (Fr.) Duby [Rhytismatales: Rhytismataceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, this species occurs on dead stems of many herbaceous plants (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Hypoxylon rubiginosum</i> var. <i>rubiginosum</i> (Pers.) Fr. [Xylariales: Xylariaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, these species occur on hardwoods and cause heart rot (Farr and Rossman 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Hypoxylon tinctor</i> (Berk.) Cooke [Xylariales: Xylariaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, these species occur on hardwoods and cause heart rot (Farr and Rossman 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Hysteroglyphium flexuosum</i> (Schwein.) Sacc. [Hysteriales: Hysteriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, congeneric species are saprobic or hemibiotrophic (Barr 1990) on wood and bark or on fallen branches (Lorenzo and Messuti 2009).	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.			
<i>Hysterographium mori</i> (Schwein.) Rehm [Hysteriales: Hysteriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Hysterographium viticola</i> (Cooke & Peck) Rehm [Hysteriales: Hysteriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, species of this fungus are saprobic or hemibiotrophic (Barr 1990) on wood and bark or on fallen branches (Lorenzo and Messuti 2009). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Hysterographium vulvatum</i> (Schwein.) Rehm [Hysteriales: Hysteriaceae]	Not known to occur		Assessment not required		
<i>Inocutis jamaicensis</i> (Murrill) Gottlieb <i>et al.</i> [Hymenochaetales: Hymenochaetaceae]	Not known to occur	<b>Yes:</b> This species is associated with grapevines, causing white rot in the trunk and main branches (Pérez <i>et al.</i> 2008) and has also been isolated from esca-affected grapevine stems (Fischer 2006). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Fischer 2006; Pérez <i>et al.</i> 2008) and may spread naturally in infected propagative material. Multiplication and marketing of infected propagative material would help spread this fungus into new areas. Therefore, this fungus has the potential for establishment and spread in Australia.	<b>Yes:</b> This fungus is associated with Esca disease of grapevine, which is one of the most important diseases of grapevine worldwide (Romanazzi <i>et al.</i> 2009). This fungus is able to colonise wide variety of hosts, including grapevine and <i>Eucalyptus</i> , in diverse conditions (Pérez <i>et al.</i> 2008). The wine industry and native <i>Eucalyptus</i> plantations in Australia could be severely affected by this fungus. Therefore, this fungus has potential for economic	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
				consequences in parts of Australia.	
<i>Irpex lacteus</i> (Fr.) Fr. [Polyporales: Meruliaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, this species occurs on trunks, dead stems and wood of host plants (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Isariopsis clavispora</i> (Berk. & MA Curtis) Sacc. [Capnodiales: Mycosphaerellaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Isariopsis fuckelii</i> (Thüm.) du Plessis [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, this species is associated with the foliage of host plants (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Kuehneola vitis</i> (E.J. Butler) Syd. & P. Syd. [Pucciniales: Phragmidaceae]	Not known to occur	No: This fungus infects fully grown leaves or older leaves and may cause leaf rust (Papademetriou and Dent 2001). Therefore, foliage free dormant cuttings do not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		provide a pathway for this fungus.			
<i>Lachnella alboviolascens</i> (Alb. & Schwein.) Fr. [Agaricales: Niaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Lachnella macrochaeta</i> Speg. [Agaricales: Niaceae] (synonym <i>Trichopezizella macrochaeta</i> (Speg.) Gamundi)	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. There is little information on the biology of these species. However, generally <i>Lachnella</i> species occur on dead twigs, dead shoots, dead stems and bark (Ellis and Everhart 1897; Seaver 1911). Some <i>Lachnella</i> species have also been reported to occur on the young shoots of herbaceous species such as senecio (McKenzie and Foggo 1989), but are not reported to occur on the living stems or shoots of grapevines. Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Lachnella myceliosa</i> WB Cooke [Agaricales: Niaceae]	Not known to occur		Assessment not required		
<i>Lasiodiplodia crassispora</i> Tl Burgess & Barber [Botryosphaerales: Botryosphaeriaceae]	Yes (Burgess <i>et al.</i> 2006)	Assessment not required			
<i>Lasiodiplodia missouriana</i> Úrbez-Torres <i>et al.</i> [Botryosphaerales: Botryosphaeriaceae]	Not known to occur	<b>Yes:</b> These species cause cankers in the vascular tissue of grapevines (Úrbez-Torres <i>et al.</i> 2012). Therefore,	<b>Yes.</b> These fungi have established in areas with a wide range of climatic conditions (Úrbez-Torres <i>et al.</i>	No: These species have been recorded on grapevines causing canker in association with other species (Úrbez-Torres <i>et al.</i>	
<i>Lasiodiplodia viticola</i> Úrbez-Torres <i>et al.</i>	Not known to				

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Botryosphaerales: Botryosphaeriaceae]	occur	dormant cuttings may provide a pathway for these fungi.	2012) and may spread naturally in infected propagative material. Therefore, these fungi have the potential for establishment and spread in Australia.	2012). However, no information is available on the losses caused by these pathogens. Therefore, these fungi are not of economic concern for Australia.	
<i>Lepteutypa cupressi</i> (Nattrass <i>et al.</i> ) HJ Swart [Xylariales: Amphisphaeriaceae] (synonym: <i>Monochaetia unicornis</i> (Cooke & Ellis) Sacc. & D. Sacc.)	Yes (PHA 2001)	Assessment not required			
<i>Leptosphaeria ampelina</i> Curzi & Barbaini [Pleosporales: Leptosphaeriaceae]	Not known to occur	No: These <i>Leptosphaeria</i> species occur on dead stems and dry runners of grapevine and on wood and dead plant material (Grand and Vernia 2004; Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Leptosphaeria cerlettii</i> Speg. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria chaetostoma</i> Sacc. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria cirricola</i> Pass. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria cookei</i> Pirota [Pleosporales: Leptosphaeriaceae] <sup>13</sup>	Yes (Shivas 1989)				
<i>Leptosphaeria gibelliana</i> Pirota [Pleosporales: Leptosphaeriaceae]	Not known to occur	No: These <i>Leptosphaeria</i> species occur on dead stems and dry runners of grapevine and on wood and dead plant material (Grand and Vernia 2004; Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Leptosphaeria ogilviensis</i> (Berk. & Broome) Ces. & De Not. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria pampini</i> (Thüm.) Sacc. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria vinealis</i> Pass. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		

<sup>13</sup> Listed as *Phoma vitis* (Shivas 1989)



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Leptosphaeria viticola</i> Fautrey & Roum. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria vitigena</i> (Schulzer) Sacc [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptosphaeria vitis</i> Schulzer ex Sacc. [Pleosporales: Leptosphaeriaceae]	Not known to occur		Assessment not required		
<i>Leptothyrium passerinii</i> Thüm. [Incertae sedis]	Not known to occur	No: This species has been recorded on grape clusters (Pearson and Goheen 1988). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Leptoxylum fumago</i> (Woron.) RC Srivastava [Capnodiales: Capnodiaceae] (synonym: <i>Fumago vagans</i> Pers.)	Yes (Phillips 1994)	Assessment not required			
<i>Lewia scrophulariae</i> (Desm.) ME Barr & EG Simmons [Pleosporales: Pleosporaceae]	Not known to occur	No: This species is a saprophyte (Bahcecioglu <i>et al.</i> 2006). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Lopharia crassa</i> (Lév.) Boidin [Polyporales: Polyporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Lophiostoma elegans</i> (Fabre) Sacc. [Pleosporales: Lophiostomataceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Lophiostoma</i> species occur on bark, dead wood and dead stems of various herbaceous plants (Farr <i>et al.</i> 1989; Farr	Assessment not required		
<i>Lophiostoma macrostomum</i> (Tode) Ces. & De Not. [Pleosporales: Lophiostomataceae]	Not known to occur		Assessment not required		
<i>Lophiostoma rhopalosporum</i> Ellis & Everh. [Pleosporales: Lophiostomataceae]	Not known to occur		Assessment not required		
<i>Lophiostoma stenostomum</i> Ellis &	Not known to		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Everh. [Pleosporales: Lophiostomataceae]	occur	and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for these fungi.			
<i>Lycoperdon radicatum</i> Durieu & Mont. [Agaricales: Agaricaceae]	Not known to occur	No: <i>Lycoperdon</i> species are saprobic and occur on soil or decayed wood in deciduous woodland (Pegler <i>et al.</i> 1995). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Macrophoma farlowiana</i> (Viala & Sauv.) Tassi [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur	<b>Yes:</b> These <i>Macrophoma</i> species have been recorded on <i>Vitis</i> species occurring on foliage, twigs, stems and fruits (Pearson and Goheen 1988; Farr and Rossman 2011). Therefore, dormant cuttings may provide a pathway for these fungi.	<b>Yes:</b> These fungi have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and may spread naturally in infected propagative material. Multiplication and marketing of infected propagative material would help spread these fungi into new areas. Therefore, these fungi have the potential for establishment and spread in Australia.	No: There is no information on the economic impact of these fungi in grape production areas. These <i>Macrophoma</i> species have not been recorded to have economic consequences. Therefore, these fungi are not of economic concern for host plants.	
<i>Macrophoma flaccida</i> (Viala & Ravaz) Cavara [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur				
<i>Macrophoma longispora</i> (Thüm. & Pass.) Berl. & Voglino [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur				
<i>Macrophoma peckiana</i> (Thüm.) Berl. & Voglino [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur				
<i>Macrophoma reniformis</i> (Viala & Ravaz) Cavara [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur				
<i>Macrophoma rimiseda</i> (Sacc.) Berl. & Voglino [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur				
<i>Macrophoma sicula</i> Scalia [Botryosphaeriales: Botryosphaeriaceae]	Not known to occur				
<i>Macrophomina phaseolina</i> (Tassi) Goid. [Botryosphaeriales: Botryosphaeriaceae]	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Marssonina viticola</i> (I. Miyake) F.L. Tai [Helotiales: Dermateaceae]	No records found	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Marssonina</i> species generally occur on leaves and cause leaf diseases on host species (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Massarina microcarpa</i> (Fuckel) Sacc. [Pleosporales: Massarinaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr <i>et al.</i> 1989), but affected plant parts are not mentioned. <i>Massarina</i> species generally have been detected on dead stems (Kirk and Cooper 2009). Therefore, dormant cuttings do not provide a pathway for this fungus	Assessment not required		
<i>Meliola vitis</i> Hansford [Meliolales: Meliolaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Hosagoudar and Archana 2009), but affected plant parts are not mentioned. <i>Meliola</i> species are associated with foliage, causing black mildew (Hosagoudar <i>et al.</i> 2010). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Micropera ampelina</i> Saccardo & Fairman [Unassigned]	Not known to occur	No: There is one record of this fungus occurring on the living limbs of <i>Vitis vinifera</i> (Farr and Rossman 2011). However, since being reported on <i>Vitis</i> species from New York in 1906 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Mollisia cinerea</i> f. <i>cinerea</i> (Batsch) P. Karst. [Helotiales: Dermateaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Mollisia</i> species are generally associated with leaves, dead wood, and old stems (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Mollisia melaleuca</i> (Fr.) Sacc. [Helotiales: Dermateaceae]	Not known to occur		Assessment not required		
<i>Mollisia pullata</i> (WR Gerard) Dennis [Helotiales: Dermateaceae]	Not known to occur		Assessment not required		
<i>Monilinia fruticola</i> (G. Winter) Honey [Helotiales: Sclerotiniaceae]	Yes (PHA 2001)	Assessment not required			
<i>Monilinia fructigena</i> Honey [Helotiales: Sclerotiniaceae]	Not known to occur	Yes: This species is associated with <i>Vitis</i> species (CABI 2012a). Cankers may develop on infected twigs and branches (Mackie 2005). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This pathogen is established in areas with a wide range of climatic conditions (Machowicz-Stefaniak and Zalewska 2002; Mackie 2005) and may spread naturally in infected propagative material. Therefore, it has the potential	<b>Yes:</b> This pathogen is less damaging on grapes (CABI 2012a); however it is of significant economic importance for apples, pears, peaches and apricots (Mackie 2005). This pathogen can cause fruit losses of 5%–35% (Mackie 2005). If introduced to Australia, it is likely	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
			for establishment and spread in Australia.	to cause serious losses to apple and pear industries in particular (Mackie 2005). Therefore, this fungus has potential for economic consequences in parts of Australia.	
<i>Monilinia laxa</i> (Aderh. & Ruhland) Honey [Helotiales: Sclerotiniaceae]	Yes (PHA 2001)	Assessment not required			
<i>Monochaetia ellisiana</i> var. <i>affinis</i> Sacc. & Briard [Xylariales: Amphisphaeriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Monochaetia sarmentii</i> (Pass.) Sacc. [Xylariales: Amphisphaeriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Monochaetia uniseta</i> (Tracy & Ellis) Sacc. [Xylariales: Amphisphaeriaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Monochaetia</i> species are generally associated with foliage or dead leaves (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Monochaetia viticola</i> (Cavara) Sacc. & D. Sacc. [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Monochaetinula ampelophila</i> (Speg.) Nag Raj [Xylariales: Amphisphaeriaceae] (synonym: <i>Monochaetia ampelophila</i> Speg)	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. These fungi are generally associated with foliage or dead leaves (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not	Assessment not required		
<i>Monochaetinula terminaliae</i> (Bat. & J.L. Bezerra) Muthumary <i>et al.</i> [Xylariales: Amphisphaeriaceae] (synonym: <i>Monochaetia terminaliae</i> Bat. & J.L. Bezerra)	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		provide a pathway for these fungi.			
<i>Mucor circinelloides</i> Tiegh. [Mucorales: Mucoraceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this fungus occurs in soils and on a variety of organic substrates (Farr <i>et al.</i> 1989). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Mucor racemosus</i> Fresen. [Mucorales: Mucoraceae]	Yes (PHA 2001)	Assessment not required			
<i>Mycosphaerella angulata</i> W.A. Jenkins [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: These <i>Mycosphaerella</i> species are associated with <i>Vitis</i> species (Farr and Rossman 2011) and cause leaf spot (Farr <i>et al.</i> 1989), resulting in premature defoliation (Pearson and Goheen 1988). These fungi overwinter in dead leaves (Pearson and Goheen 1988). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Mycosphaerella cuboniana</i> (D. Sacc.) Tomilin [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Mycosphaerella manganottiana</i> (C. Massal.) Tomilin [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Mycosphaerella personata</i> B.B. Higgins [Capnodiales: Mycosphaerellaceae]	Yes (Simmonds 1966)	Assessment not required			
<i>Mycosphaerella vitis</i> Koshk. [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		affected plant parts are not mentioned. Generally, <i>Mycosphaerella</i> species are associated with foliage, causing leaf spot (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Mycovellosiella vitis</i> Y.L. Guo & X.J. Liu [Capnodiales: Mycosphaerellaceae] (synonym: <i>Passalora vitis-piadezkii</i> U. Braun & Crous)	Not known to occur	No: This fungus has been recorded on leaves of <i>Vitis</i> species (Kirk 2012). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Myxosporium viticola</i> Dearn. & House [Unassigned]	Not known to occur	No: There is one record of this fungus occurring on the stems of <i>Vitis</i> species in Alabama in 1960 (Farr and Rossman 2011). However, this fungus has not been recorded from any other location, indicating propagative material does not provide a pathway for this fungus.	Assessment not required		
<i>Natrassia mangiferae</i> (Syd. & P. Syd.) B. Sutton & Dyko [Botryosphaerales: Botryosphaeriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Nectria cinnabarina</i> (Tode) Fr. [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Nectria coccinea</i> (Pers.) Fr. [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		affected plant parts are not mentioned. Generally, <i>Nectria</i> species are associated with hardwood trees and soil (Farr <i>et al.</i> 1989). Therefore, semi-hardwood, root free dormant cuttings do not provide a pathway for this fungus.			
<i>Nectria haematococca</i> Berk. & Broome [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Nectria radicola</i> Gerlach & L. Nilsson [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Nectria ramulariae</i> (Wollenw.) E. Müller [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Nectria</i> species are associated with hardwood trees (Farr <i>et al.</i> 1989). Therefore, semi-hardwood, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Nectria viticola</i> Berk. & Curt. [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on the limbs of <i>Vitis</i> species in Alabama in 1960 (Farr and Rossman 2011). However, this fungus has not been recorded since from any other location, indicating that propagative material does not provide a pathway for this fungus.	Assessment not required		
<i>Neofusicoccum mediterraneum</i> Crous <i>et</i>	Not known to	<b>Yes:</b> This fungus has been	<b>Yes:</b> This fungus has	No: This fungus causes	



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>al.</i> [Botryosphaeriales: Botryosphaeriaceae]	occur	recorded on grapevines (Úrbez-Torres <i>et al.</i> 2010a) and has been isolated from the vascular tissue and brown wood of young, declining grapevines (Úrbez-Torres <i>et al.</i> 2010a; Martin <i>et al.</i> 2011b). Therefore, dormant cuttings may provide a pathway for this fungus.	established in areas with a wide range of climatic conditions (Úrbez-Torres <i>et al.</i> 2010a; Martin <i>et al.</i> 2011b) and may spread naturally in infected propagative material. Therefore, this fungus has the potential for establishment and spread in Australia.	Botryosphaeria canker in association with other species (Úrbez-Torres <i>et al.</i> 2010a). However, no information is available on the losses caused by this pathogen. Therefore, this fungus is not of economic concern.	
<i>Neonectria fuckeliana</i> (C. Booth) Castl. & Rossman [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species causing basal rot (Halleen <i>et al.</i> 2006a, b). Therefore, root-free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Neonectria macrodidyma</i> Halleen <i>et al.</i> [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Neonectria mammoidea</i> W. Phillips & Plowr. [Hypocreales: Nectriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species causing basal rot (Halleen <i>et al.</i> 2006a, b). Therefore, root-free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Neonectria radicola</i> (Gerlach & L. Nilsson) Mantiri & Samuels [Hypocreales: Nectriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pareutypella sulcata</i> YM Ju & JD Rogers [Xylariales: Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		hosts it is recorded on fallen twigs (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Passalora dissiliens</i> (Duby) U. Braun & Crous [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts these fungi are associated with foliage, causing leaf spots (Farr <i>et al.</i> 1989; Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Passalora vitis</i> (MS Patil & Sawant) Poonam Srivastava [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Passalora vitis-ripariae</i> (U. Braun) U. Braun & Crous [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Patellaria atrata</i> (Hedw.) Fr. [Patellariales: Patellariaceae]	No records found	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since they were reported on <i>Vitis</i> species in 1973 in Central Asia ( <i>P. atrata</i> ) and Spain ( <i>P. viticola</i> ) (Farr and Rossman 2011), there have been no reports from any other country, indicating dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Patellaria viticola</i> Pers. [Patellariales: Patellariaceae]	Not known to occur		Assessment not required		
<i>Penicillium adametzioides</i> S. Abe ex G. Sm. [Eurotiales: Trichocomaceae]	Not known to occur	No: <i>Penicillium</i> species occur in soil, on decaying plant	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Penicillium ardesiacum</i> Novobr. [Eurotiales: Trichocomaceae]	Not known to occur	debris, decomposing fruits and stored products (Jones and Aldwinkle 1991; Shim <i>et al.</i> 2002; Schmidt <i>et al.</i> 2006; Okafor <i>et al.</i> 2007). Dormant cuttings therefore do not provide a pathway for these species.	Assessment not required		
<i>Penicillium aurantiogriseum</i> Dierckx [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Penicillium brevicompactum</i> Dierckx [Eurotiales: Trichocomaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Penicillium canescens</i> Sopp [Eurotiales: Trichocomaceae]	Not known to occur	No: <i>Penicillium</i> species occur in soil, on decaying plant debris, decomposing fruits and stored products (Jones and Aldwinkle 1991; Shim <i>et al.</i> 2002; Schmidt <i>et al.</i> 2006; Okafor <i>et al.</i> 2007). Dormant cuttings therefore do not provide a pathway for these species.	Assessment not required		
<i>Penicillium chrysogenum</i> var. <i>chrysogenum</i> Thom [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium citrinum</i> Thom [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium decumbens</i> Thom [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Penicillium digitatum</i> (Pers.) Sacc. [Eurotiales: Trichocomaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Penicillium elongatum</i> Dierckx [Eurotiales: Trichocomaceae]	Not known to occur	No: <i>Penicillium</i> species occur in soil, on decaying plant debris, decomposing fruits and stored products (Jones and Aldwinkle 1991; Shim <i>et al.</i> 2002; Schmidt <i>et al.</i> 2006; Okafor <i>et al.</i> 2007). Dormant cuttings therefore do not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		provide a pathway for this species.			
<i>Penicillium expansum</i> Link [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Penicillium funiculosum</i> Thom [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Penicillium glabrum</i> (Wehmer) Westling [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Penicillium griseoroseum</i> Dierckx [Eurotiales: Trichocomaceae]	Not known to occur	No: <i>Penicillium</i> species occur in soil, on decaying plant debris, decomposing fruits and stored products (Jones and Aldwinkle 1991; Shim <i>et al.</i> 2002; Schmidt <i>et al.</i> 2006; Okafor <i>et al.</i> 2007). Therefore, dormant cuttings do not provide a pathway for this species.	Assessment not required		
<i>Penicillium italicum</i> Wehmer [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			
<i>Penicillium janthenellum</i> Biourge [Eurotiales: Trichocomaceae]	Not known to occur	No: <i>Penicillium</i> species occur in soil, on decaying plant debris, decomposing fruits and stored products (Jones and Aldwinkle 1991; Shim <i>et al.</i> 2002; Schmidt <i>et al.</i> 2006; Okafor <i>et al.</i> 2007). Dormant cuttings therefore do not provide a pathway for these species.	Assessment not required		
<i>Penicillium kloeckeri</i> Pitt [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium purpurascens</i> (Sopp) Biourge [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium rolfsii</i> Thom [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium solitum</i> var. <i>crustosum</i> (Thom) Bridge <i>et al.</i> [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium thomii</i> Maire [Eurotiales: Trichocomaceae]	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Penicillium variabile</i> Sopp [Eurotiales: Trichocomaceae]	Not known to occur	No: <i>Penicillium</i> species occur in soil, on decaying plant debris, decomposing fruits and stored products (Jones and Aldwinkle 1991; Shim <i>et al.</i> 2002; Schmidt <i>et al.</i> 2006; Okafor <i>et al.</i> 2007). Dormant cuttings therefore do not provide a pathway for these species.	Assessment not required		
<i>Penicillium viridicatum</i> Westling [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium vitis</i> Novobr. [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Penicillium vulpinum</i> (Cooke & Massee) Seifert & Samson [Eurotiales: Trichocomaceae]	Not known to occur		Assessment not required		
<i>Peniophora albobadia</i> (Schwein.) Boidin [Russulales: Peniophoraceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Peniophora</i> species are saprobic and found on dead, bark-covered branches (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Perenniporia medulla-panis</i> (Jacq.) Donk [Polyporales: Polyporaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Perenniporia tenuis</i> var. <i>tenuis</i> (Schwein.) Ryvarden [Polyporales: Polyporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. This species occurs on bark or wood causing white rot (Gilbertson and Bigelow 1998; Farr and Rossman 2011). Therefore, semi-hardwood dormant	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		cuttings do not provide a pathway for this fungus.			
<i>Periconia byssoides</i> Pers. [Pleosporales: Unassigned]	Yes (PHA 2001)	Assessment not required			
<i>Pestalotia briardii</i> Lendn. [Xylariales: Amphisphaeriaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, members of this genus are secondary pathogens; they are saprophytic on dead and dying plant tissues (CAES 2008). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Pestalotia europaea</i> Grove [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia malicola</i> Hori. [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia menezesiana</i> Bres. & Torrend [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia monochaetoidea</i> var. <i>affinis</i> Sacc. & Briard [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia pezizoides</i> De Not. [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia pitospora</i> MEA Costa & Sousa da Câmara [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia quadriciliata</i> Bubak & Dearness [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia thuemenii</i> Speg. [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia uniseta</i> Tracy & Earle [Xylariales: Amphisphaeriaceae]	Not known to occur		Assessment not required		
<i>Pestalotia viticola</i> Cavara [Xylariales: Amphisphaeriaceae]	Yes (Sergeeva <i>et al.</i> 2005)	Assessment not required			
<i>Pestalotiopsis funerea</i> (Desm.) Steyaert [Xylariales: Amphisphaeriaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pestalotiopsis guepinii</i> (Desm.) Steyaert [Xylariales: Amphisphaeriaceae]	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Pestalotiopsis menezesiana</i> (Bres. & Torrend) Bissett [Xylariales: Amphisphaeriaceae]	Yes (Sergeeva <i>et al.</i> 2005)	Assessment not required			
<i>Pestalotiopsis uvicola</i> (Speg.) Biss. [Xylariales: Amphisphaeriaceae]	Yes (Sergeeva <i>et al.</i> 2005)	Assessment not required			
<i>Phaeoacremonium aleophilum</i> Gams <i>et al.</i> [Diaporthales: Togniniaceae]	Yes (Edwards and Pascoe 2004)	Assessment not required			
<i>Phaeoacremonium alvesii</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur	<b>Yes:</b> <i>Phaeoacremonium</i> species <sup>14</sup> colonise the vascular system of plants (Chicau <i>et al.</i> 2000; Marco <i>et al.</i> 2004; Eskalen <i>et al.</i> 2005; Mostert <i>et al.</i> 2006b; Gramaje <i>et al.</i> 2007; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a). These fungi have been found in apparently healthy asymptomatic grapevines (Ridgway <i>et al.</i> 2003; Aroca and Raposo 2009). Therefore, propagative material may provide a pathway for these	<b>Yes:</b> These fungi have established in areas with a wide range of climatic conditions (Chicau <i>et al.</i> 2000; Marco <i>et al.</i> 2004; Eskalen <i>et al.</i> 2005; Mostert <i>et al.</i> 2006b; Gramaje <i>et al.</i> 2007; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a) and can spread naturally in infected propagative material (Mugnai <i>et al.</i> 1999; Ridgway <i>et al.</i> 2003; Giménez-Jaime <i>et al.</i> 2006; Aroca and Raposo 2009). Multiplication and	<b>Yes.</b> <i>Phaeoacremonium</i> species are involved in Petri disease in young vines and esca in adult vines (Mostert <i>et al.</i> 2006a,b; Aroca and Raposo 2009, Gramaje <i>et al.</i> 2009a). Petri disease pathogens act as pioneer organisms that facilitate the invasion of the wood decay fungi that cause the typical symptoms of Esca disease inside the trunk and branches (Larignon and Dubos 1997). Petri disease and Esca disease limit both vineyard longevity and	Yes
<i>Phaeoacremonium angustius</i> Gams <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium argentinense</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium armeniacum</i> Graham <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes

<sup>14</sup> Taxonomy of this genus has been repeatedly reviewed, with new species described in recent years. Several species of *Phaeoacremonium* have been isolated from grapevines, although their pathogenicity has not been demonstrated for all of them (Aroca and Raposo 2009). Four species (*P. aleophilum*, *P. angustius*, *P. inflatipes*, and *P. parasiticum*) were described based on morphological and cultural characteristics (Crous *et al.* 1996). Two additional species (*P. viticola* and *P. mertoniae*) were described based on phenotypic characters, the internal transcribed spacer (ITS) regions 1 and 2, the 5.8S rDNA (Dupont *et al.* 2000) and the b-tubulin gene (Groenewald *et al.* 2001). Subsequent studies based on actin and calmodulin gene regions identified seven additional species (*P. australiense*, *P. austroafricanum*, *P. iraniamum*, *P. krajdenii*, *P. scolyti*, *P. subulatum*, *P. venezuelense*) from grapevines (Mostert *et al.* 2005, 2006b). Recently, four more species of *Phaeoacremonium* (*P. croatiense*, *P. hungaricum*, *P. sicilianum*, *P. tuscanum*) from grapevine has been described (Essakhi *et al.* 2008). Additionally, three more species of *Phaeoacremonium* (*P. alvesii*, *P. griseorubrum*, *P. rubrigenum*) previously known from humans have been reported on grapevines (Essakhi *et al.* 2008). More recently two species (*P. cinereum*, *P. hispanicum*) have been identified based on combined DNA sequences of the actin and b-tubulin genes (Gramaje *et al.* 2009a). *Phaeoacremonium* species occur as part of a disease complex with *Phaeomoniella chlamydospora* causing Petri disease in younger vines and with several basidiomycete species causing esca in older vines (Mugnai *et al.* 1999; Edwards and Pascoe 2004; Fischer 2006).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		pathogens.	marketing of infected propagative material will help spread these pathogens within Australia. Additionally, these fungi are also known to be wind-borne (Rooney-Latham <i>et al.</i> 2005) or spread by grafting (Halleen <i>et al.</i> 2003) and pruning tools (Mugnai <i>et al.</i> 1999). Therefore, they have the potential to establish and spread in Australia.	productivity as woody parts of the vine are killed (Urbez- Torres <i>et al.</i> 2012) and affect yield, wine quality and berry quality (White 2010). Consequently, <i>Phaeoacremonium</i> species have great impact on the wine, table grape and raisin industries (White 2010). Therefore, <i>Phaeoacremonium</i> strains from grapevines have the potential for economic consequences in Australia.	
<i>Phaeoacremonium australiense</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Yes (PHA 2001)	Assessment not required			
<i>Phaeoacremonium austroafricanum</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur	<b>Yes:</b> <i>Phaeoacromonium</i> species colonise the vascular system of plants (Chicau <i>et al.</i> 2000; Marco <i>et al.</i> 2004; Eskalen <i>et al.</i> 2005; Mostert <i>et al.</i> 2006b; Gramaje <i>et al.</i> 2007; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a). <i>Phaeoacremonium</i> species have been found in apparently healthy asymptomatic grapevines (Ridgway <i>et al.</i> 2003). Therefore, propagative material may provide a pathway for these pathogens.	<b>Yes:</b> These fungi have established in areas with a wide range of climatic conditions (Chicau <i>et al.</i> 2000; Marco <i>et al.</i> 2004; Eskalen <i>et al.</i> 2005; Mostert <i>et al.</i> 2006b; Gramaje <i>et al.</i> 2007; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a) and can spread naturally in infected propagative material (Mugnai <i>et al.</i> 1999; Ridgway <i>et al.</i> 2003; Giménez-Jaime <i>et al.</i> 2006; Aroca and Raposo 2009). Multiplication and marketing of infected propagative material will help spread these pathogens	<b>Yes.</b> <i>Phaeoacremonium</i> species are involved in Petri disease in young vines and esca in adult vines (Mostert <i>et al.</i> 2006a,b; Aroca and Raposo 2009, Gramaje <i>et al.</i> 2009a). Petri disease pathogens act as pioneer organisms that facilitate the invasion of the wood decay fungi that cause the typical symptoms of Esca disease inside the trunk and branches (Larignon and Dubos 1997). Petri disease and Esca disease limit both vineyard longevity and productivity as woody parts of the vine are killed (Urbez- Torres <i>et al.</i> 2012) and affect yield, wine	<b>Yes</b>
<i>Phaeoacremonium cinereum</i> Gramaje <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>
<i>Phaeoacremonium croatiense</i> Essakhi <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>
<i>Phaeoacremonium globosum</i> Graham <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>
<i>Phaeoacremonium griseorubrum</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>
<i>Phaeoacremonium hispanicum</i> Gramaje <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>
<i>Phaeoacremonium hungaricum</i> Essakhi <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>
<i>Phaeoacremonium inflatipes</i> Gams <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Phaeoacremonium iraniana</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur		within Australia. Additionally, these fungi are also known to be wind-borne or spread by grafting and pruning tools (Mugnai <i>et al.</i> 1999). Therefore, they have the potential to establish and spread in Australia.	quality and berry quality (White 2010). Consequently, <i>Phaeoacremonium</i> species have great impact on the wine, table grape and raisin industries (White 2010). Therefore, <i>Phaeoacremonium</i> strains from grapevines have the potential for economic consequences in Australia.	Yes
<i>Phaeoacremonium kraidenii</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium mortoniae</i> Crous & W. Gams [Diaporthales: Togniniaceae] (synonym <i>Togninia fraxinopennsylvanica</i> (T.E. Hinds) Hausner <i>et al.</i> )	Not known to occur				Yes
<i>Phaeoacremonium occidentale</i> Graham <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium parasiticum</i> (Ajello <i>et al.</i> ) Gams [Diaporthales: Togniniaceae]	Yes (Mostert <i>et al.</i> 2006b)	Assessment not required			
<i>Phaeoacremonium rubrigenum</i> Gams <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur	<b>Yes:</b> <i>Phaeoacromonium</i> species colonise the vascular system of plants (Chicau <i>et al.</i> 2000; Marco <i>et al.</i> 2004; Eskalen <i>et al.</i> 2005; Mostert <i>et al.</i> 2006b; Gramaje <i>et al.</i> 2007; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a). <i>Phaeoacremonium</i> species have been found in apparently healthy asymptomatic grapevines (Ridgway <i>et al.</i> 2003). Therefore, propagative material may provide a pathway for these pathogens.	<b>Yes:</b> These fungi have established in areas with a wide range of climatic conditions (Chicau <i>et al.</i> 2000; Marco <i>et al.</i> 2004; Eskalen <i>et al.</i> 2005; Mostert <i>et al.</i> 2006b; Gramaje <i>et al.</i> 2007; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a) and can spread naturally in infected propagative material (Mugnai <i>et al.</i> 1999; Ridgway <i>et al.</i> 2003; Giménez-Jaime <i>et al.</i> 2006; Aroca and Raposo 2009). Multiplication and marketing of infected propagative material will help spread these pathogens within Australia. Additionally, these fungi are also known to be wind-borne or spread by grafting and pruning tools	<b>Yes.</b> <i>Phaeoacremonium</i> species are involved in Petri disease in young vines and esca in adult vines (Mostert <i>et al.</i> 2006a,b; Aroca and Raposo 2009, Gramaje <i>et al.</i> 2009a). Petri disease pathogens act as pioneer organisms that facilitate the invasion of the wood decay fungi that cause the typical symptoms of Esca disease inside the trunk and branches (Larignon and Dubos 1997). Petri disease and Esca disease limit both vineyard longevity and productivity as woody parts of the vine are killed (Urbez- Torres <i>et al.</i> 2012) and affect yield, wine quality and berry quality (White 2010). Consequently, <i>Phaeoacremonium</i> species have great impact on the wine, table	Yes
<i>Phaeoacremonium scolymii</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium sicilianum</i> Essakhi <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium subulatum</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium tuscanicum</i> Essakhi <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium venezuelense</i> Mostert <i>et al.</i> [Diaporthales: Togniniaceae]	Not known to occur				Yes
<i>Phaeoacremonium viticola</i> J. Dupont [Diaporthales: Togniniaceae].	Not known to occur				Yes

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
			(Mugnai <i>et al.</i> 1999). Therefore, they have the potential to establish and spread in Australia.	grape and raisin industries (White 2010). Therefore, <i>Phaeoacremonium</i> strains from grapevines have the potential for economic consequences in Australia.	
<i>Phaeomoniella chlamydospora</i> (Gams <i>et al.</i> ) Crous & W. Gams [Chaetothyriales: Herpotrichiellaceae]	Yes (Edwards and Pascoe 2004)	Assessment not required			
<i>Phakopsora ampelopsidis</i> Dietel & P. Syd. [Pucciniales: Phakopsoraceae]	Not known to occur	No <sup>15</sup> : This fungus is host specific and does not occur on grapevines (Ono 2000) and therefore is not on the pathway.	Assessment not required		
<i>Phakopsora cronartiiformis</i> Dietel [Pucciniales: Phakopsoraceae]	Not known to occur	No <sup>16</sup> : This fungus is host specific and does not occur on grapevines (Ono <i>et al.</i> 1990) and therefore is not on the pathway.	Assessment not required		
<i>Phakopsora euvitidis</i> Y. Ono [Pucciniales: Phakopsoraceae]	Not known to occur <sup>17</sup>	Yes <sup>18</sup> : These fungi are associated with grapevine causing leaf rust (Chatasiri	Yes: These rust fungi have established in areas with a wide range of climatic	Yes. These rust fungi are serious pathogens of grapevines (Leu 1988; EPPO 2002a;	Yes
<i>Phakopsora muscadinae</i> Buritica	Not known to				Yes

<sup>15</sup> Recent taxonomic studies partly clarified the situation of *Phakopsora* species causing grapevine rust. *Phakopsora ampelopsidis* was previously identified as the pathogen causing grape leaf rust of *Vitis* spp. (Hiratsuka 1935 cited in Hennessy *et al.* 2007). However, recent studies based on differences in host specificity, lifecycle and morphology of *Phakopsora ampelopsidis* isolated from these hosts indicated that this fungus consists of three taxonomically distinct species (Ono 2000). *Phakopsora ampelopsidis* and *Phakopsora vitis* are host specific and occur on *Ampelopsis brevipedunculata* and *Parthenocissus tricuspidata* respectively (Hennessy *et al.* 2007). Therefore *Phakopsora ampelopsidis* is not considered in this assessment. Based on the work by Ono (2000), the records of *P. ampelopsidis* on *Vitis* species are assumed to be *P. euvitidis*.

<sup>16</sup> *Phakopsora cronartiiformis* has previously been recorded on grapevine, however, further studies indicated that it is host specific and occurs on *Parthenocissus semicordata* (Ono *et al.* 1990). Therefore *Phakopsora cronartiiformis* is not considered in this assessment.

<sup>17</sup> *Phakopsora euvitidis* was detected in Darwin in 2001 (Weinert *et al.* 2003) and declared eradicated in 2006 (Liberato *et al.* 2007).

<sup>18</sup> Three rust fungi namely *Phakopsora euvitidis* (Asian grapevine leaf rust), *Phakopsora muscadinae* and *Phakopsora uva* (American grapevine leaf rust) are associated with grapevines in Asian and Americas (Chatasiri and Ono 2008).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Pucciniales: Phakopsoraceae] <i>Phakopsora uva</i> Buriticá & J.F. Hennen [Pucciniales: Phakopsoraceae]	occur  Not known to occur	and Ono 2008). These rust species generally infect leaves (Ono 2000; Weinert <i>et al.</i> 2003; Hennessy <i>et al.</i> 2007; Chatasiri and Ono 2008), however they can overwinter as mycelium in grapevine shoots (EPPO 2002a) or dormant buds (Weinert <i>et al.</i> 2003; Hennessy <i>et al.</i> 2007). Therefore, dormant cuttings may provide a pathway for these rust fungi.	conditions (EPPO 2002a; Chatasiri and Ono 2008) and can spread naturally in infected propagative material (EPPO 2002a). Distribution of propagative material carrying mycelium in dormant buds will help spread these rust fungi within Australia. Additionally, spores are dispersed by wind and rain splash (EPPO 2002a). These dispersal mechanisms would facilitate spread within Australia. Therefore, these rust fungi have the potential to establish and spread in Australia.	Angelotti <i>et al.</i> 2008) and have potential to be destructive under favourable conditions (Tessmann <i>et al.</i> 2004; Angelotti <i>et al.</i> 2008). Heavy infection causes necrosis of leaves and in severe cases can lead to defoliation of the host plant. The disease can cause poor shoot growth, reduction of fruit quality and yield loss in commercial grapevine production (Leu 1988; EPPO 2002a; Angelotti <i>et al.</i> 2008). Therefore, <i>Phakopsora</i> species have the potential for economic consequences in Australia.	Yes
<i>Phakopsora vitis</i> P. Syd. [Pucciniales: Phakopsoraceae]	Not known to occur	No <sup>19</sup> : This fungus is host specific and does not occur on grapevines (Hennessy <i>et al.</i> 2007) and therefore is not on the pathway.	Assessment not required		
<i>Phanerochaete flavidoalba</i> (Cooke) S.S. Rattan [Polyporales: Phanerochaetaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. On other hosts, these species are associated with dead branches of fallen trees and cause white rot of hardwood, conifer and other	Assessment not required		
<i>Phanerochaete viticola</i> (Schwein.) Parmasto [Polyporales: Phanerochaetaceae]	Not known to occur		Assessment not required		

<sup>19</sup> *Phakopsora vitis* has previously been recorded on grapevine, however, further studies indicated that this fungus is host specific and occurs on *Parthenocissus tricuspidata* (Hennessy *et al.* 2007). Therefore *Phakopsora vitis* is not considered in this assessment.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		woody debris (Burdall 1985). Therefore, dormant cuttings do not provide a pathway for these fungi.			
<i>Phellinus gilvus</i> (Schwein.) Pat. [Hymenochaetales: Hymenochaetaceae]	Yes (PHA 2001)	Assessment not required			
<i>Phellinus igniarius</i> (L.) Quél. [Hymenochaetales: Hymenochaetaceae]	Not known to occur	No: This fungus was considered the causal agent of esca disease in grapevines (Reisenzein <i>et al.</i> 2000). However, further studies indicate that the isolates from esca affected vines, identified as <i>P. igniarius</i> , were misidentifications of <i>Fomitiporia punctata</i> (Mugnai <i>et al.</i> 1999; Cortesi <i>et al.</i> 2000). Therefore, this species is not assessed.	Assessment not required		
<i>Phellinus noxius</i> (Corner) G. Cunn. [Hymenochaetales: Hymenochaetaceae]	Yes (PHA 2001)	Assessment not required			
<i>Phellinus viticola</i> (Schwein.) Donk [Hymenochaetales: Hymenochaetaceae]	Not known to occur	No: Members of this genus occur on living or dead wood and cause wood rot (Farr <i>et al.</i> 1989; Brooks 2002; Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Phlyctibasidium polyporoideum</i> (Berkeley & MA Curtis) Jülich [Unassigned]	Not known to occur	No: This species occurs on rotting wood (Gilbertson and Bigelow 1998). Therefore, dormant cuttings do not provide a pathway for this	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		fungus.			
<i>Phoma ampelina</i> Berk. & M.A. Curtis [Pleosporales: Incertae sedis]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Phoma</i> species are soil-borne, weakly parasitic or saprophytic species and are associated with roots, dead stems and foliage of host plants (Boerema 1976; Farr <i>et al.</i> 1989; Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Phoma ampelocarpa</i> Pass. [Pleosporales: Incertae sedis]	Not known to occur		Assessment not required		
<i>Phoma exigua</i> Sacc. [Pleosporales: Incertae sedis]	Yes (PHA 2001)	Assessment not required			
<i>Phoma glomerata</i> (Corda) Wollenw. & Hochapfel [Pleosporales: Incertae sedis]	Yes (PHA 2001)	Assessment not required			
<i>Phoma lenticularis</i> Cavara [Pleosporales: Incertae sedis]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Phoma</i> species are soil-borne, weakly parasitic or saprophytic species and are associated with roots, dead stems and foliage of host plants (Boerema 1976; Farr <i>et al.</i> 1989; Farr and Rossman 2011). Therefore, dormant cuttings do not provide a			
<i>Phoma negriana</i> Thüm. [Pleosporales: Incertae sedis]	Not known to occur		Assessment not required		
<i>Phoma plurivora</i> PR Johnston [Pleosporales: Incertae sedis]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		pathway for these fungi.			
<i>Phoma pomorum</i> Thüm. [Pleosporales: Incertae sedis]	Yes (Cook and Dubé 1989)	Assessment not required			
<i>Phomopsis longiparaphysata</i> Uecker & KC Kuo [Diaporthales: Diaporthaceae]	Not known to occur	No: This fungus is known to occur on fruit (Uecker and Kuo 1992). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Phomopsis viticola</i> (Sacc.) Sacc [Diaporthales: Diaporthaceae]	Yes (Savocchia <i>et al.</i> 2007)	Assessment not required			
<i>Phyllachora picea</i> (Berk. & M.A. Curtis) Sacc. [Phyllachorales: Phyllachoraceae]	Not known to occur	<b>Yes:</b> This species has been recorded on <i>Vitis</i> species and is associated with the stem (Farr and Rossman 2011). Therefore, dormant cuttings may provide a pathway for this fungus.	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and it may spread in infected propagative material. Therefore, these fungi have the potential to establish and spread in Australia	No: This species has been reported on grapes but no economic losses have been reported. Therefore, this fungus is not of economic concern to Australia.	
<i>Phyllachora pomigena</i> (Schwein.) Sacc. [Phyllachorales: Phyllachoraceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Phyllachora vitis</i> MS Patil & AB Pawar [Phyllachorales: Phyllachoraceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Phyllachora</i> species are generally associated with foliage (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Phyllactinia ampelopsidis</i> YX Yu & YQ	Not known to	No: This fungus has been	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Lai [Erysiphales: Erysiphaceae]	occur	recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Phyllactinia</i> species occur on foliage and cause powdery mildew (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Phyllactinia guttata</i> (Wallr.) Lév. [Erysiphales: Erysiphaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Phyllosticta ampelophila</i> Politis [Botryosphaerales: Botryosphaeriaceae]	Not known to occur	No: These <i>Phyllosticta</i> species have been recorded on the foliage of <i>Vitis</i> species (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for these species.	Assessment not required.		
<i>Phyllosticta badhami</i> Cooke [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required.		
<i>Phyllosticta dzumajensis</i> Bubák [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required.		
<i>Phyllosticta labruscae</i> Thüm. [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required		
<i>Phyllosticta microspila</i> Pass. [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required.		
<i>Phyllosticta pilispora</i> Speschnew [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required.		
<i>Phyllosticta spermoides</i> Peck. [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required.		
<i>Phyllosticta vitis</i> Sacc. [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required.		
<i>Phymatotrichopsis omnivora</i> (Duggar) Hennebert [Pezizales: Rhiziniaceae]	Not known to occur	No: This species is a soil-borne pathogen associated with the roots of host plants (Farr <i>et al.</i> 1989). Therefore,	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		root free dormant cuttings do not provide a pathway for this fungus.			
<i>Physalospora baccae</i> Cavara [Xylariales: Hyponectriaceae]	Not known to occur	No: This species infects grape berries, leaves, pedicels and peduncles (Zhang 2005). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Pilidiella diplodiopsis</i> Crous & Van Niekerc [Diaporthales: Schizoparmaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this species affects fruit (Lauber and Schuepp 1968). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Pleospora betae</i> (Berl.) Nevod. [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pleospora herbarum</i> (Pers.) Rabenh. [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pleospora penicillus</i> var. <i>penicillus</i> Fuckel [Pleosporales: Pleosporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Therefore, there is no evidence that propagative material provides a pathway for this fungus.	Assessment not required		
<i>Pleospora phaeocomoides</i> (Berk. & Broome) G. Winter [Pleosporales:	Yes (Farr and Rossman 2011)	Assessment not required			



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Pleosporaceae]					
<i>Pleospora vitis</i> Catt. [Pleosporales: Pleosporaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Therefore, there is no evidence that propagative material provides a pathway for this fungus.	Assessment not required		
<i>Pleospora vitis-viniferae</i> Frolov [Pleosporales: Pleosporaceae]	Not known to occur		Assessment not required		
<i>Pleurostomophora richardsiae</i> (Nannf.) Mostert <i>et al.</i> [Calosphaeriales: Pleurostomataceae]	Not known to occur	No: This fungus has been isolated from cankered grapevines (Varela <i>et al.</i> 2011) and cankers generally develop in the woody parts of the vine (Urbez-Torres <i>et al.</i> 2012). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Pleurotus ostreatus</i> (Jacq.) P. Kumm. [Agaricales: Pleurotaceae]	Yes (PHA 2001)	Assessment not required			
<i>Poria papyracea</i> (Schwein.) Cooke [Polyporales: Polyporaceae]	Yes (May <i>et al.</i> 2003)	Assessment not required			
<i>Pseudocercospora riachueli</i> (Speg.) Deighton. [Capnodiales: Mycosphaerellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pseudocercospora vitis</i> (Lév.) Speg. [Capnodiales: Mycosphaerellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pseudopezicula tetraspora</i> Korf <i>et al.</i> [Helotiales: Helotiaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011) and occur on the leaves (Pearson and Goheen 1988). The pathogen overwinters in fallen	Assessment not required		
<i>Pseudopezicula tracheiphila</i> (Müll.-Thurg.) Korf & WY Zhuang [Helotiales: Helotiaceae]	Not known to occur		Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		leaves (Pearson and Goheen 1988). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.			
<i>Pseudovalsa viticola</i> Ellis & Everh. [Diaporthales: Pseudovalsaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species and occurs on dead stems (Farr and Rossman 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Pyrenochaeta vitis</i> Viala & Sauv. [Pleosporales: Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species causing leaf spot (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Pyrenophora phaeocomes</i> (Rebent.) Fr. [Pleosporales: Pleosporaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Pyrenophora</i> species are associated with foliage and cause leaf spot (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Ramularia khandalensis</i> Patw. & A.K. Pande [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		affected plant parts are not mentioned. <i>Ramularia</i> species generally occur on leaves and cause leaf spot (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Resupinatus poriaeformis</i> (Pers.) Thorn <i>et al.</i> [Agaricales: Tricholomataceae]	Not known to occur	No: Species of this genus occur on rotting logs and other herbaceous and woody debris (Farr <i>et al.</i> 1989; Thorn <i>et al.</i> 2005). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Rhabdospora ampelina</i> (Thüm.) Sacc. [Capnodiales: Mycosphaerellaceae]	Not known to occur	<b>Yes:</b> These fungi have been recorded on <i>Vitis</i> species, occurring on stems (Farr and Rossman 2011). Therefore, dormant cuttings may provide a pathway for these fungi.	<b>Yes:</b> These fungi have established in areas with a wide range of climatic conditions (Farr and Rossman 2011) and it may spread in infected propagative material. Therefore, these fungi have the potential to establish and spread in Australia.	No: These fungi have been reported on grapes but no economic consequences have been reported. Therefore, these fungi are not of economic concern to Australia.	
<i>Rhabdospora labruscae</i> Gonz. Frag. [Capnodiales: Mycosphaerellaceae]	Not known to occur				
<i>Rhabdospora mueggenburgii</i> (Pirota) Sacc. [Capnodiales: Mycosphaerellaceae]	Not known to occur				
<i>Rhabdospora vitis</i> Koshk. & Frolov [Capnodiales: [Mycosphaerellaceae]	Not known to occur				
<i>Rhacodiella vitis</i> Sterenberg [Helotiales: Sclerotiniaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011) and causes spotted necrosis on woody vines (Winkler <i>et al.</i> 1974; Cline and Farr 2006). This fungus only affects grapevines that have been subject to the poor	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		management practice of covering vines with soil over winter (Winkler <i>et al.</i> 1974). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Rhizoctonia solani</i> JG Kuhn [Ceratobasidiales: Ceratobasidiaceae]	Yes (Neate <i>et al.</i> 1988)	Assessment not required			
<i>Rhizopus arrhizus</i> var. <i>arrhizus</i> A. Fisch [Mucorales: Mucoraceae]	Yes (PHA 2001)	Assessment not required			
<i>Rhizopus stolonifer</i> var. <i>stolonifer</i> (Ehrenb.) Vuill. [Mucorales: Mucoraceae]	Yes (PHA 2001)	Assessment not required			
<i>Rhytisma vitis</i> Schwein. [Rhytismatales: Rhytismataceae]	Not known to occur	No: This species occurs on leaves and causes the formation of black spots (Pearson and Goheen 1988). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Robillarda sessilis</i> (Sacc.) Sacc. [Unassigned]	Yes (PHA 2001)	Assessment not required			
<i>Robillarda vitis</i> Prillieux & Delacroix [Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Other species in the genus <i>Robillarda</i> occur on foliage and cause leaf spots (Giri <i>et al.</i> 1996). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Roesleria subterranea</i> (Weinm.) Redhead [Incertae sedis: Roesleriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species and causes root rot (Farr and Rossman 2011). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Rosellinia amblystoma</i> Berl. & F. Sacc. [Xylariales: Xylariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Members of the genus <i>Rosellinia</i> occur on roots and cause root rot (Petrini and Petrini 2005). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Rosellinia aquila</i> (Fr.) Ces. & De Not. [Xylariales: Xylariaceae]	Yes (PHA 2001)	Assessment not required			
<i>Rosellinia langloisii</i> Ellis & Everh. [Xylariales: Xylariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Members of the genus <i>Rosellinia</i> occur on roots and cause root rot (Petrini and Petrini 2005). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Rosellinia necatrix</i> Berl. ex Prill. [Xylariales: Xylariaceae]	Yes (PHA 2001)	Assessment not required			
<i>Rosellinia pulveracea</i> (Ehrh.) Fuckel	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Xylariales: Xylariaceae]					
<i>Rosellinia rosarum</i> Niessl [Xylariales: Xylariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Members of the genus <i>Rosellinia</i> occur on roots and cause root rot (Petrini and Petrini 2005). Therefore, root free dormant cuttings do not provide a pathway for this fungus.			
<i>Sacidium viticola</i> Cooke [Mucorales: Pilobolaceae]	Not known to occur	No: These fungi have been recorded on the leaves of <i>Vitis</i> species (Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Sacidium vitis</i> Ellis & Everh. [Mucorales: Pilobolaceae]	Not known to occur		Assessment not required		
<i>Schizophyllum commune</i> Fr. [Agaricales: Schizophyllaceae]	Yes (PHA 2001)	Assessment not required			
<i>Schizopora paradoxa</i> (Schr.) Donk [Hymenochaetales: Schizoporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Schizoxylon insigne</i> (De Not.) Rehm [Ostropales: Stictidaceae]	Not known to occur	No: This species has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. There is no evidence of this species occurring on grapevine stems. Therefore, this species is not on the pathway of dormant grapevine cuttings.	Assessment not required		
<i>Sclerotinia sclerotiorum</i> (Lib.) de Bary	Yes (Shivas	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Helotiales: Sclerotiniaceae]	1989)				
<i>Sclerotium rolfsii</i> Sac [Helotiales: Sclerotiniaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Scytinostroma alutum</i> Lanq. [Russulales: Lachnocladiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this species occurs on dead wood (BCCM 2012). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Sebacina incrustans</i> (Pers.) Tul. & C. Tul. [Sebacinales: Sebacinaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. This fungus occurs on woody stems, leaves and plant debris (Farr <i>et al.</i> 1989). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Seimatosporium hysterioides</i> (Fuckel) Brockmann [Xylariales: Amphisphaeriaceae]	Yes (Sergeeva <i>et al.</i> 2005)	Assessment not required			
<i>Seimatosporium lonicerae</i> (Cooke) Shoemaker [Xylariales: Amphisphaeriaceae]	Yes (Shivas 1989)	Assessment not required			
<i>Seimatosporium macrospermum</i> (Berk. & Broome) B. Sutton [Xylariales: Amphisphaeriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		mentioned. There is no evidence that this species occurs on the stems of grapevines. Therefore, this species is not on the pathway of dormant grapevine cuttings.			
<i>Seimatosporium parasiticum</i> (Dearn. & House) Shoemaker [Xylariales: Amphisphaeriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Septoria ampelina</i> Berk. & M.A. Curtis [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: These fungi have been recorded on the foliage of <i>Vitis</i> species (Pearson and Goheen 1988; Farr <i>et al.</i> 1989; Farr and Rossman 2011). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Septoria badhami</i> Berk. & Broome [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Septoria kellermaniana</i> Thüm. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Septoria melanopsis</i> Pat. [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Septoria tassiana</i> Syd [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Septoria vineae</i> Pass [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Septoria viticola</i> Berk. & M.A. Curtis [Capnodiales: Mycosphaerellaceae]	Not known to occur		Assessment not required		
<i>Septosporium heterosporum</i> Ellis & Galloway [Unassigned] (synonym: <i>Passalora heterosporella</i> U. Braun & Crous, <i>Phaeoramularia heterospora</i> (Ellis & Galloway) Deighton)	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. On other hosts, this species occurs on leaves (Deighton 1976). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Setosphaeria rostrata</i> K.J. Leonard [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Sorosphaera viticola</i> Kirchmair <i>et al.</i> [Plasmodiophorida: Plasmodiophoraceae]	Not known to occur	No: This species has been recorded on <i>Vitis</i> species and is associated with roots (Kirchmair <i>et al.</i> 2005). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Sphaceloma viticola</i> Sawada ex Jenkins & Bitanc [Myriangiales: Elsinoaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its first report on <i>Vitis</i> species in Taiwan in 1944 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Sphaeria antiqua</i> Ellis & Everh [Xylariales: Xylariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since its first report on <i>Vitis</i> species in New Jersey in 1954 (Farr and Rossman 2011), it has not been reported from any other country, indicating propagative material does not provide a pathway for this fungus.	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Sphaeropsis ampelos</i> (Schwein.) Cooke [Botryosphaerales: Botryosphaeriaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011); but affected plant parts are not mentioned. <i>Sphaeropsis</i> species are generally associated with the foliage, cones, bark and wood of host plants (Farr <i>et al.</i> 1989). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Sphaeropsis vitigena</i> Ellis & Everh [Botryosphaerales: Botryosphaeriaceae]	Not known to occur		Assessment not required		
<i>Sporidesmium raulii</i> Ellis & Harkn. [Pleosporales: Unassigned]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since being reported on <i>Vitis</i> species from Pennsylvania in 1954 and 1959 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Sporocadus rhododendri</i> (Schwein.) M. Morelet [Xylariales: Amphispheeriaceae]	Yes (Sergeeva <i>et al.</i> 2005)	Assessment not required			
<i>Stachybotrys chartarum</i> (Ehrenb.) SJ Hughes [Hypocreales: Unassigned]	Yes (PHA 2001)	Assessment not required			
<i>Stagonospora bulgarica</i> Vanev [Pleosporales: Phaeosphaeriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		mentioned. <i>Stagonospora</i> species are generally associated with foliage (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.			
<i>Stemphylium botryosum</i> Sacc. [Pleosporales: Pleosporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Stereum albobadium</i> (Schwein.) Fr. [Russulales: Stereaceae]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Stereum</i> species are associated with hardwood (Farr <i>et al.</i> 1989). Therefore, semi-hardwood dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Stereum crassum</i> (Lév.) Fr. [Russulales: Stereaceae]	Not known to occur		Assessment not required		
<i>Stereum hirsutum</i> (Willd.) Pers. [Russulales: Stereaceae]	Yes (Tovar <i>et al.</i> 2008)	Assessment not required			
<i>Stereum purpureum</i> Pers. [Russulales: Stereaceae]	Yes (Cook and Dubé 1989)	Assessment not required			
<i>Stigmina esfandiarii</i> Petr. [Capnodiales: Mycosphaerellaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Members of this genus occur on foliage, bark and dead twigs (Farr <i>et al.</i> 1989). Therefore, foliage free, semi-hardwood dormant cuttings do not provide a	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		pathway for this fungus.			
<i>Strickeria sylvana</i> (Sacc. & Speg.) Cooke [Unassigned]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since they were reported on <i>Vitis</i> species in Poland ( <i>S. sylvana</i> ) and Central Asia ( <i>S. trubicola</i> ) in 1973 (Farr and Rossman 2011), they have not been reported from any other country, indicating dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Strickeria trubicola</i> (Fuckel) G. Winter [Unassigned]	Not known to occur		Assessment not required		
<i>Synchytrium parthenocissi</i> M.T. Cook [Chytridiales: Synchytriaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since it was reported on <i>Vitis</i> species in Louisiana in 1964 (Farr and Rossman 2011), it has not been reported elsewhere, indicating dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Teichospora winteriana</i> Berl. [Pleosporales: Dacampiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Other species of this genus have been recorded on dead branches	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		and stems of host plants (Rao 1966). Therefore, dormant cuttings do not provide a pathway for this fungus.			
<i>Thelephora atra</i> Weinm. [Thelephorales: Thelephoraceae]	Yes (May <i>et al.</i> 2003)	Assessment not required			
<i>Thielaviopsis basicola</i> (Berk. & Broome) Ferraris [Microascales: Ceratocystidaceae]	Yes (PHA 2001)	Assessment not required			
<i>Thyridium vitis</i> Ellis & Everh. [Incertae sedis: Thyridiaceae]	Not known to occur	No: This species is recorded on the dead shoots of <i>Vitis</i> species (Anon 2011). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Tilletiopsis minor</i> Nyland [Unassigned]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species	Assessment not required		
<i>Tilletiopsis washingtonensis</i> Nyland [Unassigned]	Not known to occur	(Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts they occur on leaves (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Tomentella bryophila</i> (Pers.) MJ Larsen [Thelephorales: Thelephoraceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. However, on other hosts it occurs on wood (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for this	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		fungus.			
<i>Trametes hirsuta</i> (Wulfen) Lloyd [Polyporales: Polyporaceae]	Yes (PHA 2001)	Assessment not required			
<i>Trametes ochracea</i> (Pers.) Gilb. & Ryvarden [Polyporales: Polyporaceae]	Yes (GBIF 2012)	Assessment not required			
<i>Trametes versicolor</i> (L.) Lloyd [Polyporales: Polyporaceae]	Yes (Tovar <i>et al.</i> 2008)	Assessment not required			
<i>Trematosphaeria vitigena</i> Ellis & Everhart [Pleosporales: Pleomassariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since it was reported on <i>Vitis</i> species in West Virginia in 1954 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Trichocladium asperum</i> Harz [Sordariales: Chaetomiaceae]	Yes (PHA 2001)	Assessment not required			
<i>Trichoderma koningii</i> Oudem. [Hypocreales: Hypocreaceae]	Yes (PHA 2001)	Assessment not required			
<i>Trichoderma viride</i> Pers. [Hypocreales: Hypocreaceae]	Yes (PHA 2001)	Assessment not required			
<i>Trichothecium roseum</i> (Pers.) Link [Hypocreales: Incertae sedis]	Yes (PHA 2001)	Assessment not required			
<i>Trullula melanochlora</i> (Desm.) Höhn. [Unassigned]	Not known to occur	<b>Yes:</b> This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011) and is associated with cane bleaching (Phillips 2000).	<b>Yes:</b> This fungus has established in areas with a wide range of climatic conditions (Phillips 2000) and it may spread in infected	No: This fungus has been reported on grapes, but no economic losses have been reported (Phillips 2000). Therefore, this fungus is not of	

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		Therefore, dormant cuttings may provide a pathway for this fungus.	propagative material. Therefore, this fungus has the potential to establish and spread in Australia.	economic concern to Australia.	
<i>Truncatella angustata</i> (Pers.) S. Hughes [Xylariales: Amphispheariaceae]	Yes (Sergeeva <i>et al.</i> 2005)	Assessment not required			
<i>Tryblidaria indica</i> Tilak [Patellariales Patellariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Since it was reported on <i>Vitis</i> species in India in 1966 (Farr and Rossman 2011), it has not been reported from any other country, indicating dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Tubercularia acinorum</i> Cavara [Hypocreales: Nectriaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Tubeufia pezizula</i> (Berk. & M.A. Curtis) M.E. Barr [Pleosporales: Tubeufiaceae]	Yes (Farr and Rossman 2011)	Assessment not required			
<i>Typhula viticola</i> (Peck) Berthier [Agaricales: Typhulaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Typhula</i> species generally occur on fallen, rotting leaves (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Ulocladium atrum</i> Preuss [Pleosporales:	Yes (PHA 2001)	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Pleosporaceae]					
<i>Uredo cissicola</i> Cummins [Unassigned]	Not known to occur	No: These fungi have been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Uredo</i> species generally occur on leaves and cause leaf rust (Farr <i>et al.</i> 1989). Therefore, foliage free dormant cuttings do not provide a pathway for these fungi.	Assessment not required		
<i>Uredo cissi-pterocladae</i> Hirats. [Unassigned]	Not known to occur		Assessment not required		
<i>Uromyces cladomanes</i> Traverso [Pucciniales: Pucciniaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. <i>Uromyces</i> species are generally associated with leaf and stem rust (Farr <i>et al.</i> 1989). However, since it was reported on <i>Vitis</i> in 1937, it has not been reported from any other country, indicating that dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Valsa ceratosperma</i> (Tode) Maire [Diaporthales: Valsaceae]	Yes (PHA 2001)	Assessment not required			
<i>Valsa vitigera</i> Cooke [Diaporthales: Valsaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Most <i>Valsa</i> species affect the dead twigs	Assessment not required		



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		and bark of mature trees (Jones and Aldwinkle 1991, Farr and Rossman 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.			
<i>Valsaria insitiva</i> (Tode) Ces. & De Not. [Diaporthales: Valsaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this species is saprobic on dead wood (Farr <i>et al.</i> 1989; Ellis and Ellis 1997). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Vararia pectinata</i> (Burt) DP Rogers & HS Jacks. [Russulales: Lachnocladiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Vararia</i> species occur on wood and dead branches (Farr <i>et al.</i> 1989). Therefore, dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Vermicularia compacta</i> Cooke & Ellis [Incertae sedis: Glomerellaceae]	Yes (PHA 2001)	Assessment not required			
<i>Verticillium albo-atrum</i> Reinke & Berthold [Incertae sedis: Plectosphaerellaceae]	Yes (Walker 1990)	Assessment not required			
<i>Verticillium dahliae</i> Kleb. [Incertae sedis: Plectosphaerellaceae]	Yes (Harding and	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Plectosphaerellaceae]	Wicks 2007)				
<i>Xenosporium berkeleyi</i> (M.A. Curtis) Piroz. [Pleosporales: Tubeufiaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, this species occurs on decaying, woody substrates (Farr and Rossman 2011). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Xylaria arbuscula</i> Sacc. [Xylariales: Xylariaceae]	Not known to occur	No: This fungus has been recorded on <i>Vitis</i> species (Farr and Rossman 2011), but affected plant parts are not mentioned. Generally, <i>Xylaria</i> species cause decay of dead stumps and hardwood timber (Sivanesan and Holliday 1972). Therefore, semi-hardwood dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Xylaria hypoxylon</i> (L.) Grev. [Xylariales: Xylariaceae]	Yes (PHA 2001)	Assessment not required			
<i>Xylaria polymorpha</i> (Pers.) Grev. [Xylariales: Xylariaceae]	Yes (PHA 2001)	Assessment not required			
<b>STRAMINOPIILA</b>					
<i>Phytophthora cactorum</i> (Lebert & Cohn) J. Schröt. [Peronosporales: Peronosporaceae]	Yes (Golzar <i>et al.</i> 2007)	Assessment not required			
<i>Phytophthora cambivora</i> (Petri) Buisman	Yes (Wicks and	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Peronosporales: Peronosporaceae]	Hall 1990)				
<i>Phytophthora cinnamomi</i> Rands [Peronosporales: Peronosporaceae]	Yes (Cahill <i>et al.</i> 2008)	Assessment not required			
<i>Phytophthora citricola</i> Sawada [Peronosporales: Peronosporaceae]	Yes (Stukely <i>et al.</i> 2007)	Assessment not required			
<i>Phytophthora cryptogea</i> Pethybr. & Laff. [Peronosporales: Peronosporaceae]	Yes (Stukely <i>et al.</i> 2007)	Assessment not required			
<i>Phytophthora drechsleri</i> Tucker [Peronosporales: Peronosporaceae]	Yes (Stukely <i>et al.</i> 2007)	Assessment not required			
<i>Phytophthora megasperma</i> Drechsler [Peronosporales: Peronosporaceae]	Yes (Stukely <i>et al.</i> 2007)	Assessment not required			
<i>Phytophthora nicotianae</i> Breda de Haan [Peronosporales: Peronosporaceae]	Yes (Stukely <i>et al.</i> 2007)	Assessment not required			
<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni [Peronosporales: Peronosporaceae]	Yes (Constable and Drew 2004)	Assessment not required			
<i>Pythium acanthicum</i> Drechsler [Pythiales: Pythiaceae]	Yes (Vaartaja 1965)	Assessment not required			
<i>Pythium aphanidermatum</i> (Edson) Fitzp. [Pythiales: Pythiaceae]	Yes (Male and Vawdrey 2010)	Assessment not required			
<i>Pythium debaryanum</i> R. Hesse [Pythiales: Pythiaceae]	Yes (Wong <i>et al.</i> 1985)	Assessment not required			
<i>Pythium irregulare</i> Buisman [Pythiales: Pythiaceae]	Yes (Vaartaja 1965)	Assessment not required			
<i>Pythium mamillatum</i> Meurs [Pythiales: Pythiaceae]	Yes (Vaartaja 1965)	Assessment not required			
<i>Pythium middletonii</i> Sparrow [Pythiales: Pythiaceae]	Yes (Irwin and Jones 1977)	Assessment not required			
<i>Pythium parasiticum</i> S. Rajagop. & K. Ramakr. [Pythiales: Pythiaceae]	Not known to occur	No: <i>Pythium</i> species are soil-borne and infect the roots of host plants, causing various rots, lesions, damping-off, discoloration, abnormal	Assessment not required		

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		growth, dieback and death (Farr and Rossman 2011). Therefore, root free dormant cuttings do not provide a pathway for this fungus.			
<i>Pythium rostratum</i> E.J. Butler [Pythiales: Pythiaceae]	Yes (Vaartaja 1965)	Assessment not required			
<i>Pythium spinosum</i> Sawada [Pythiales: Pythiaceae]	Yes (Wong <i>et al.</i> 1985)	Assessment not required			
<i>Pythium splendens</i> Hans Braun [Pythiales: Pythiaceae]	Yes (PHA 2001)	Assessment not required			
<i>Pythium sylvaticum</i> WA Campbell & FF Hendrix [Pythiales: Pythiaceae]	Not known to occur	No: <i>Pythium</i> species are soil-borne and infect the roots of host plants, causing various rots, lesions, damping-off, discoloration, abnormal growth, dieback and death (Farr and Rossman 2011). Therefore, root free dormant cuttings do not provide a pathway for this fungus.	Assessment not required		
<i>Pythium ultimum</i> Trow [Pythiales: Pythiaceae]	Yes (Vaartaja 1965)	Assessment not required			
<i>Pythium vexans</i> de Bary [Pythiales: Pythiaceae]	Yes (Irwin and Jones 1977)	Assessment not required			
<b>PHYTOPLASMA<sup>20</sup></b>					
Buckland Valley grapevine yellows	Yes (Constable	Assessment not required			

<sup>20</sup> Phytoplasmas are classified on the basis of molecular data obtained from 16S rDNA and other conserved genes into distinct groups, subgroups and species belonging to the newly established 'Candidatus Phytoplasma' taxon (IRPCM 2004). Initially, differentiation of the phytoplasma was based on the geographical origins of the diseases, the specific hosts and insect vectors and the symptoms exhibited by the host plant. However, given that the same phytoplasma strain may induce different symptoms in different hosts and different strains may share common vectors or cause diseases showing similar symptoms, this approach did not provide an accurate means of phytoplasma classification (Weintraub and Jones 2010). Therefore, the designation of a new/distinct 'Candidatus Phytoplasma' species is based on the nucleotide sequence of the 16S rRNA gene.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
(BVGY) Phytoplasma [16Srl-related]	2010)				
<i>Candidatus</i> Phytoplasma asteris [16Srl – Aster yellows group] <sup>21</sup> (Virginia grapevine yellows I (VGYI), Aster yellow phytoplasma)	Not known to occur	<b>Yes:</b> Phytoplasmas are obligate parasitic, phloem-restricted pathogens that cause grapevine yellows <sup>22</sup> (Weintraub and Jones 2010). Several molecularly distinct phytoplasma groups which cause grapevine yellows have been identified (Hren <i>et al.</i> 2009). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994). Propagative material therefore provides a pathway for this phytoplasma.	<b>Yes:</b> <i>Candidatus</i> phytoplasma asteris has established in areas with a wide range of climatic conditions of different grapevine regions of the world (Constable 2010) and can spread naturally in infected propagative material (Caudwell <i>et al.</i> 1994; Matus <i>et al.</i> 2008; Constable 2010). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, grape infecting phytoplasmas have the potential to establish and spread in Australia.	<b>Yes:</b> The aster yellows group of phytoplasmas are associated with over 100 economically important diseases worldwide (Lee <i>et al.</i> 2004a). Typical symptoms of grapevine yellows include leaf chlorosis and rolling, flower abortion or berry withering, uneven or total lack of lignification of canes and stunting (Olivier <i>et al.</i> 2009b). Therefore, this phytoplasma group has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Candidatus</i> Phytoplasma australiense [16SrXII-B] (strains: Australian)	Yes (Constable 2010)	Assessment not required			

<sup>21</sup> Phytoplasmas classified in subgroups 16Srl-A, 16Srl-B and 16Srl-C ('*Candidatus* Phytoplasma asteris'-related strains) are associated with grapevine yellows in several countries (Bianco *et al.* 1994; Alma *et al.* 1996; Davis *et al.* 1998). 16Srl-B and 16Srl-C have sporadically been found in grapevine (the strains related to 'Ca. Phytoplasma asteris' comprises of a large number of related phytoplasma worldwide, representing the most diverse and widespread phytoplasma group [Lee *et al.* 2004a]). Although there is relatively high similarity in the 16S rDNA sequence, the strains in this group occupy diverse ecological niches and show substantial genetic variation (Firrao *et al.* 2005). Earlier studies placed Tomato big bud mycoplasma like organism and Tomato big bud phytoplasma in the 'Ca. Phytoplasma asteris group' (Firrao 2004). However, recent studies have placed Tomato big bud phytoplasma in the SrlI-D ribosomal group (Constable 2010).

<sup>22</sup> Grapevine yellows (GY) is a term that is used to refer to any of several grapevine diseases that are currently attributed to infection of grapevine plants by phytoplasmas. Grapevine yellows diseases include flavescence dorée, Palatinate grapevine yellows, and Bois noir (black wood, legno nero), reported in southern Europe and the Mediterranean region; North American grapevine yellows (Virginia grapevine yellows I, Virginia grapevine yellows III, New York grapevine yellows, and grapevine yellows in Canada); Australian grapevine yellows in Australia and New Zealand and Buckland Valley grapevine yellows in Australia; and grapevine yellows diseases that have been reported in other regions including South Africa and Chile. While the symptoms caused by different GY are similar, they show considerable differences in epidemiology due to the different life history of their respective vectors (Boudon-Padieu 2005). All vectors of GY identified so far are leafhoppers and planthoppers (Boudon-Padieu 2005).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
grapevine yellows (AGY) Phytoplasma) <i>Candidatus</i> Phytoplasma fraxini [16SrVII-A] (Ash yellows group) – Chile grapevine yellows strain	Not known to occur	<b>Yes:</b> Phytoplasmas are phloem restricted and symptoms include downward leaf rolling, yellowing or reddening of the leaves and incomplete shoot lignification (Gajardo <i>et al.</i> 2009). Mixed phytoplasma infections and infections of phytoplasmas together with one or more viruses also occur (Gajardo <i>et al.</i> 2009). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994); therefore dormant cuttings provide a pathway for this phytoplasma.	<b>Yes:</b> Chile grapevine yellows has established in areas with a wide range of climatic conditions in different regions of the world (Gajardo <i>et al.</i> 2009). Phytoplasmas generally spread naturally in infected propagative material (Constable 2010). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, grape infecting phytoplasmas have the potential to establish and spread in Australia.	<b>Yes:</b> This phytoplasma occurs on a range of hosts, including economically important crops such as peach (Zunnoo-Khan <i>et al.</i> 2010). It also causes devastating effects in ornamentals (Zunnoo-Khan <i>et al.</i> 2010). This phytoplasma causes leaf-reddening, yellowing, shortening of internodes, shoot proliferation, reduced fruit size and plant decline (Griffiths <i>et al.</i> 1999; Zunnoo-Khan <i>et al.</i> 2010). Therefore, this phytoplasma group has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Candidatus</i> Phytoplasma phoenicium [16SrIX]	Not known to occur	<b>Yes:</b> Phytoplasmas associated with grape yellows are obligate parasites and phloem restricted. Infected grapevines show redness and inward curling of leaves (Canik <i>et al.</i> 2011). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994); therefore dormant cuttings provide a pathway for these phytoplasmas.	<b>Yes:</b> This phytoplasma group has established in areas with a wide range of climatic conditions in different regions of the world (Canik <i>et al.</i> 2011). Phytoplasmas generally spread naturally in infected propagative material (Constable 2010). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, grape infecting phytoplasmas have the potential to establish and spread in Australia.	<b>Yes:</b> The 16SrIX group has been identified in grapevines in Turkey, and can cause severe diseases in host plants (Canik <i>et al.</i> 2011). Therefore, this phytoplasma group has the potential for economic consequences in Australia.	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Candidatus</i> Phytoplasma pruni [16SrIII – peach X-disease phytoplasmas group]	Not known to occur	<b>Yes:</b> Phytoplasmas are phloem restricted and symptoms include yellowing of the leaves and die-back of young shoot tips (Martelli and Boudon-Padieu 2006). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994); therefore dormant cuttings provide a pathway for this phytoplasma.	<b>Yes:</b> North American grapevine yellows has established in areas with a wide range of climatic conditions in different regions of the world (Martelli and Boudon-Padieu 2006). Phytoplasmas generally spread naturally in infected propagative material (Martelli and Boudon-Padieu 2006). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, this phytoplasma group has the potential to establish and spread in Australia.	<b>Yes:</b> <i>Candidatus</i> Phytoplasma pruni is the causal agent for several diseases, previously known as peach leaf roll, peach rosette, little peach, red suture and cherry buckskin (Olivier <i>et al.</i> 2009a). Disease incidences of up to 10% have been reported in peach orchards in the United States (Olivier <i>et al.</i> 2009a). This phytoplasma group causes economic losses associated with reduced fruit quality and yield (Olivier <i>et al.</i> 2009a). Therefore, this phytoplasma group has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Candidatus</i> Phytoplasma solani [16SrXII–A] (Stolbur group) (strains: Vergilbungskrankheit (VK) phytoplasma, Bois noir (BN) phytoplasma) <sup>23</sup>	Not known to occur	<b>Yes:</b> Phytoplasmas are phloem restricted and symptoms include downward leaf rolling, yellowing or reddening of the leaves and incomplete shoot lignification (Gajardo <i>et al.</i> 2009). Mixed phytoplasma infections and infections of phytoplasmas	<b>Yes:</b> <i>Candidatus</i> Phytoplasma solani has established in areas with a wide range of climatic conditions of different regions of the world (Constable 2010) and can spread naturally in infected propagative material (Constable 2010; Zorloni <i>et al.</i>	<b>Yes:</b> Bois noir Phytoplasma causes severe damage in European vineyards (Mori <i>et al.</i> 2007). Existence of different strains and mixed infections of different strains (Pacífico <i>et al.</i> 2009) may increase the severity of damage in vineyards and sometimes infected vines die-off	<b>Yes</b>

<sup>23</sup> Bois Noir (BN) was considered a form of Flavescence doree (FD) phytoplasma with a possible common aetiology (Caudwell 1961). Further studies indicated that BN phytoplasma is different from FD phytoplasma as both phytoplasma have different vectors (Caudwell 1961, Sforza *et al.* 1998). BN phytoplasma is associated with the stolbur group and the name *Candidatus* Phytoplasma solani has been recommended as it infects various solanaceous plants (Firrao *et al.* 2005). '*Candidatus* Phytoplasma solani'-related strains; have been classified in group 16SrXII (the stolbur phytoplasmas group (STOL)) subgroup A (formerly called subgroup 16SrI-G). Three STOL types I, II and III have been identified and was shown to be associated with distinctive host plants (Langer and Maixner 2004, Berger *et al.* 2009). Type I and II are more common in grapevine but both have different alternative hosts (Pacífico *et al.* 2009).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		together with one or more viruses also occur (Gajardo <i>et al.</i> 2009). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994); therefore dormant cuttings provide a pathway for this phytoplasma.	2011). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, grape infecting phytoplasmas have the potential to establish and spread in Australia.	during winter (Riedle-Bauer <i>et al.</i> 2006). BN phytoplasma is considered of quarantine concern by Canada. Presence of this phytoplasma group in Australia would impact upon Australia's ability to access overseas markets. Therefore, this phytoplasma group has the potential for economic consequences in Australia.	
<i>Candidatus</i> Phytoplasma ulmi [16SrV–A] (Elm yellows group EY group) <sup>24</sup>	Not known to occur	<b>Yes:</b> Phytoplasmas are found in the phloem sieve tubes of plants (Hren <i>et al.</i> 2009) causing grapevine yellows. Several molecularly distinct phytoplasma groups which cause grapevine yellows have been identified (Hren <i>et al.</i> 2009). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994); therefore dormant cuttings provide a pathway for these phytoplasmas.	<b>Yes:</b> EY group infection of grapevines has established in areas with a wide range of climatic conditions in different regions of the world (Botti and Bertaccini 2007) and can spread naturally in infected propagative material (Constable 2010; Zorloni <i>et al.</i> 2011). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, this phytoplasma group has the potential to establish and spread in Australia.	<b>Yes:</b> Many diseases inflicted by the EY group Phytoplasmas are economically important and are quarantine pathogens internationally (Lee <i>et al.</i> 2004b). Phytoplasmas generally reduce fruit yield and infected clusters have high levels of acid and low sugar content (Boudon-Padiou <i>et al.</i> 1989). Therefore, this phytoplasma group has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Candidatus</i> Phytoplasma vitis [16SrV]	Not known to	<b>Yes:</b> FD Phytoplasma is	<b>Yes:</b> FD Phytoplasma has	<b>Yes:</b> Flavescence dorée is one	<b>Yes</b>

<sup>24</sup> The EY phytoplasma (16SrV) group consists of diverse phytoplasma strains, representing the third largest phytoplasma cluster after the aster yellows and X-disease phytoplasma groups (Gundersen *et al.* 1996, Lee *et al.* 2000). Other EY group phytoplasmas associated with diseases in grapevines include flavescence dorée (FD) and grapevine yellows phytoplasmas in the European grapevine (Bertaccini *et al.* 1997, Daire *et al.* 1997, Martini *et al.* 2002, Seemuller *et al.* 1994). Strains of 16SrV–A detected in grapevines are distinguishable from strains detected in elms indicating that the phytoplasma in the 16SrV group are able to modify their genome according to environmental conditions (Botti and Bertaccini 2007).



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
(Elm yellows group) (strains: Grapevine Flavescence dorée (FD) phytoplasma; German Palatinat grapevine yellows phytoplasma) <sup>25</sup>	occur	phloem restricted (Hren <i>et al.</i> 2009) and symptoms include downward leaf rolling, yellowing or reddening of the leaves and incomplete shoot lignification (Gajardo <i>et al.</i> 2009). FD and BN Phytoplasma has been reported in grapevine (Bertaccini <i>et al.</i> 1995; Daire <i>et al.</i> 1997). Most grapevine rootstocks are potentially symptomless (Caudwell <i>et al.</i> 1994). This may lead to collection of budwood from symptomless parts of infected vines or from recently infected vines that have not developed symptoms (Martelli and Boudon-Padieu 2006). Propagative material therefore provides a pathway for these phytoplasmas.	established in areas with a wide range of climatic conditions in different regions of the world (Constable 2010) and can spread naturally in infected propagative material (Caudwell <i>et al.</i> 1994; Rott <i>et al.</i> 2007; Matus <i>et al.</i> 2008; Constable 2010). Phloem-feeding hemipterans acquire the pathogen for subsequent transmission (Boudon-Padieu <i>et al.</i> 1989). The symptomless nature of phytoplasmas may contribute to the inadvertent propagation and distribution of infected material that will help spread grape infecting phytoplasmas within Australia. Therefore, grape infecting phytoplasmas have the potential to establish and spread in Australia.	of the most serious diseases of grapevine (Margaria <i>et al.</i> 2007). Phytoplasmas generally reduce fruit yield and infected clusters have high acid levels and low sugar content (Boudon-Padieu <i>et al.</i> 1989). FD Phytoplasma is considered of quarantine concern by COSAVE and Canada. The presence of this phytoplasma group in Australia would impact upon Australia's ability to access overseas markets. Therefore, this phytoplasma group has the potential for economic consequences in Australia.	
European stone fruit yellows Phytoplasma 16SrX-B (Apple proliferation group)	Not known to occur	<b>Yes:</b> Phytoplasmas are found in the phloem sieve tubes of plants (Duduk <i>et al.</i> 2003; Hren <i>et al.</i> 2009) and cause leaf yellowing, leaf rolling and	<b>Yes:</b> This phytoplasma has established in areas with a wide range of climatic conditions in different regions of the world (Varga <i>et al.</i>	<b>Yes:</b> European stone fruit yellows cause various diseases in European stone fruit (Laimer Da Câmara Machado <i>et al.</i> 2001). In apricots, it causes leaf	<b>Yes</b>

<sup>25</sup> Flavescence dorée is caused by several isolates which belong to the 16SrV-C and -D phytoplasma phylogenetic subgroups (Filippin *et al.* 2009). Based on sequence analysis three strain clusters of FD phytoplasma (FD-1, FD-2, FD-3) have been recognized (Arnaud *et al.* 2007). FD-1 is restricted to France and Italy, FD-2 is detected in France, Italy and Spain, whereas FD-3 has been detected in Italy, Serbia and Slovenia (Constable 2010). Recent evidence indicates that the German Palatinat grapevine yellows phytoplasma is related to alder-infecting strains and is a member of the flavescence dorée phytoplasma phylogenetic subclade (Arnaud *et al.* 2007). Alder yellows and Palatinat grapevine yellows diseases in Europe are also attributed to 'Ca. Phytoplasma vitis'-related strains. Phytoplasma FD-associated strains belong to ribosomal subgroups 16SrV-C and 16SrV-D (Botti and Bertaccini 2007).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		shoot drop (Varga <i>et al.</i> 2000). Phytoplasmas are transmitted by propagative material (Caudwell <i>et al.</i> 1994); therefore dormant cuttings provide a pathway for these phytoplasmas.	2000; Duduk <i>et al.</i> 2003) and can spread naturally in infected propagative material (Caudwell <i>et al.</i> 1994; Constable 2010). Distribution of infected propagative material will help spread grape infecting phytoplasmas within Australia. Therefore, grape infecting phytoplasmas have the potential to establish and spread in Australia.	rolling, leaf chlorosis, leaf reddening, phloem necrosis and sudden dieback (Laimer Da Câmara Machado <i>et al.</i> 2001). In addition, affected apricot trees produce shrunken, tasteless fruit that fall prematurely from the tree (Laimer Da Câmara Machado <i>et al.</i> 2001). Therefore, this phytoplasma group has the potential for economic consequences in Australia.	
Tomato big bud Phytoplasma [16SrII-D] <sup>26</sup>	Yes (Constable 2010)	Assessment not required			
<b>VIROIDS</b>					
<i>Australian grapevine viroid</i> (AGVd) [Pospiviroidae: <i>Apescaviroid</i> ]	Yes (Rezaian 1990)	Assessment not required			
<i>Citrus exocortis viroid</i> – grapevine (CEVd-g) [Pospiviroidae: <i>Pospiviroid</i> ] (synonym: Grapevine viroid – slow (Gvd-s))	Yes (Hardy <i>et al.</i> 2008)	Assessment not required			
Grapevine yellow speckle viroid 1 (GYSVd1) [Pospiviroidae: <i>Apescaviroid</i> ]	Yes (Koltunow <i>et al.</i> 1989)	Assessment not required			
Grapevine yellow speckle viroid 2 (GYSVd2) [Pospiviroidae: <i>Aspcaviroid</i> ] synonym: Grapevine viroid (GV1B), Grapevine viroid-fast (Gvd-f)	Yes (Koltunow <i>et al.</i> 1989)	Assessment not required			
Grapevine yellow speckle viroid 3 (GYSVd3) [Pospiviroidae: <i>Aspcaviroid</i> ] (synonym: Chinese grapevine viroid)	Yes (Benson <i>et al.</i> 2008).	Assessment not required			

<sup>26</sup> The classification of phytoplasmas is continuously reviewed resulting in the reclassification of some of these phytoplasmas.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
Hop stunt viroid – grapevine (HSVd-g) [Pospiviroidae: <i>Hostuviroid</i> ]	Yes (Koltunow <i>et al.</i> 1988)	Assessment not required			
<b>VIRUSES</b>					
<i>Alfalfa mosaic virus</i> (AMV) [Bromoviridae: Alfamovirus]	Yes (Garraan and Gibbs 1982)	Assessment not required			
<i>Arabis mosaic virus</i> (ArMV) – grape strain [Secoviridae: Nepovirus]	Not known to occur <sup>27</sup>	<b>Yes:</b> ArMV-grape strain infections are often symptomless and expression varies based on type of rootstock, grape variety, and environmental conditions (Anon 2011). ArMV is also seed-borne in grapevines (Lazar <i>et al.</i> 1990). ArMV-grape strains may cause mottling and flecking on the leaves and leaf deformation, including enations (Anon 2011; Oklahoma State University 2011). This may lead to the propagation and distribution of infected propagative material, suggesting that ArMV-grape strains could enter Australia on propagative material.	<b>Yes:</b> ArMV-grape strains have established in areas with a wide range of climatic conditions (Cadman <i>et al.</i> 1960; Kearns and Mossop 1984; MacKenzie <i>et al.</i> 1996; Delibašić <i>et al.</i> 2000; Abelleira <i>et al.</i> 2010) and can spread naturally in infected propagative material (Anon 2011). Distribution of infected propagative material will help spread ArMV-grape strains within Australia. Therefore, ArMV-grape strains have the potential to establish and spread in Australia.	<b>Yes:</b> Infected plants may have shortened internodes and exhibit vine decline symptoms (Oklahoma State University 2011). This virus can also cause very poor fruit set in affected vines (Abelleira <i>et al.</i> 2010). ArMV can be present in a mixed infection with GFLV (Weber <i>et al.</i> 2002). ArMV-grape strains are considered of quarantine significance by some trading partners. Presence of ArMV-grape strains in Australia would impact upon Australia's ability to access overseas markets. Therefore, ArMV-grape strains have the potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Artichoke Italian latent virus</i> (AILV) [Secoviridae: Nepovirus]	Not known to occur	<b>Yes:</b> AILV is soil-borne (Kyriakopoulou 2008) and causes fanleaf symptoms in grapevine (Jankulova <i>et al.</i>	<b>Yes:</b> AILV has established in areas with a wide range of climatic conditions (Roca <i>et al.</i> 1975; Savino <i>et al.</i> 1977;	<b>Yes:</b> AILV is an economically important virus due to its extensive host range and the yield losses it can cause in some	<b>Yes</b>

<sup>27</sup> *Arabis* mosaic virus (ArMV) has once been recorded on *Narcissus* species in Australia; however, ArMV has not been recorded in grapes in Australia (Constable and Drew 2004; Constable *et al.* 2010). ArMV strains may differ in host range, symptom expression and transmissibility by nematode vectors (Jones *et al.* 1989).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		1978; Martelli and Boudon-Padieu 2006). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material does provide a pathway for AILV.	Gallitelli <i>et al.</i> 2004; Kyriakopoulou 2008) and it can spread naturally in infected propagative material. Distribution of infected propagative material will help spread AILV within Australia. Therefore, AILV has the potential to establish and spread in Australia.	hosts (Gallitelli <i>et al.</i> 2004). No information is available on economic losses caused by this virus in grapes, but AILV causes patchy chlorotic stunting disease in artichokes. Infected crops are rendered unproductive (Brown <i>et al.</i> 1997). Therefore, AILV has the potential for economic consequences in parts of Australia.	
<i>Blueberry leaf mottle virus</i> (BLMoV) New York (NY) strain [Secoviridae: Nepovirus] <sup>28</sup>	Not known to occur	<b>Yes:</b> NY strain is associated with fanleaf like symptoms (Oliver and Fuchs 2011) and is seed-borne in grapevines (Uyemoto <i>et al.</i> 1977). BLMoV-NY strain symptoms include pale green foliage and irregular elongation of shoots (Uyemoto <i>et al.</i> 1977). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for this virus.	<b>Yes:</b> BLMoV-NY strain has established in areas with a wide range of climatic conditions (Uyemoto <i>et al.</i> 1977) and it can spread naturally in infected propagative material. Distribution of infected propagative material will help spread BLMoV-NY strain within Australia. Therefore, BLMoV-NY strain has the potential to establish and spread in Australia.	<b>Yes:</b> Information on the economic consequences of BLMoV-NY strain on grapes is limited. However, as it is a part of the virus complex associated with fanleaf degeneration/decline disease (Oliver and Fuchs 2011), it may cause significant crop losses. BLMoV-NY strain induces delayed bud break and straggly fruit clusters (Uyemoto <i>et al.</i> 1977). This may reduce yield and fruit quality. Therefore, BLMoV-NY strain has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Broad bean wilt virus</i> (BBWV) [Secoviridae: Fabavirus]	Yes (Schwinghamer <i>et al.</i> 2007)	Assessment not required			

<sup>28</sup> A strain of *Blueberry leaf mottle virus* (BLMoV) related to but different from *Grapevine Bulgarian latent virus* has been reported to infect grapevines in the USA (Uyemoto *et al.* 1977).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Carnation mottle carmovirus</i> (CarMV) [Tombusviridae: Carmovirus]	Yes (Moran 1994)	Assessment not required			
<i>Cherry leafroll virus</i> (CLRV) – grape isolate [Secoviridae: Nepovirus]	Nor known to occur <sup>29</sup>	<b>Yes:</b> CLRV is associated with fanleaf like symptoms (Martelli and Boudon-Padieu 2006). The symptoms caused by CLRV on grapes include leaf yellowing, leaf chlorosis and yellow leaf mosaic symptoms (Ipach <i>et al.</i> 2003). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for CLRV.	<b>Yes:</b> CLRV has established in areas with a wide range of climatic conditions (Herrera and Madariaga 2001; Ipach <i>et al.</i> 2003) and it can spread naturally in infected propagative material. Distribution of infected propagative material will help spread CLRV within Australia. Therefore, CLRV has the potential to establish and spread in Australia.	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent. However, as it is a part of the virus complex associated with fanleaf degeneration/decline disease (Oliver and Fuchs 2011), it may cause significant crop losses. CLRV causes leaf yellowing, leaf chlorosis, yellow leaf mosaic symptoms, small fruit and premature berry abscission (Ipach <i>et al.</i> 2003). This may reduce yield and fruit quality. Therefore, this virus has potential for economic consequences in Australia.	<b>Yes</b>
<i>Cucumber mosaic virus</i> (CMV) – grape isolate (CMV-YA200) [Bromoviridae: Cucumovirus]	Nor known to occur <sup>30</sup>	<b>Yes:</b> CMV grape isolate naturally infects grapevine (Koklu <i>et al.</i> 1998) and infections are symptomless (Koklu <i>et al.</i> 1999). This may lead to the propagation and distribution of infected	<b>Yes:</b> CMV grape isolate has established in areas with a wide range of climatic conditions (Paradies <i>et al.</i> 2000). The symptomless nature of this virus may contribute to the inadvertent	No: Information on the economic consequences of this virus is almost non-existent. CMV does not appear to be a threatening pathogen to grapes as infections are apparently symptomless (Paradies <i>et al.</i> 2000) and	

<sup>29</sup> *Cherry leafroll virus* (CLRV) has been reported from rhubarb in Australia (Parmenter *et al.* 2009); however, CLRV has not been recorded in grapes in Australia (Constable and Drew 2004; Constable *et al.* 2010). The rhubarb isolate was identified using sequencing; the Australian isolate is substantially different from other important strains (Parmenter *et al.* 2009). CLRV isolates from different hosts may differ in their serological and molecular traits (Jones 1985; Jones *et al.* 1990; Rebenstorf *et al.* 2006) as well as in their host specificity and ability to induce symptoms (Jones 1973; Rowhani and Mircetich 1988). CLRV isolates segregate into six major groups based on the primary host: birch and cherry (group A); rhubarb, ash and ground elder (group B); raspberry, sorrel and chive (group C); walnut (groups D1 and D2); and elderberry (group E) (Rebenstorf *et al.* 2006).

<sup>30</sup> *Cucumber mosaic virus* (CMV) is recorded in Australia (Carpenter and Lockett 2003, Persley and Gambley 2010). However, this virus has not been recorded on grapevines in Australia. Grapevine isolates possesses a number of properties differing enough from those of other characterized CMV isolates (Paradies *et al.* 2000).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		propagative material, suggesting that CMV grape isolate could enter Australia on propagative material.	propagation and distribution of infected material that will help spread CMV grape isolate within Australia. Therefore, CMV grape isolate has the potential to establish and spread in Australia.	economic consequences are not reported. Therefore, this virus does not have the potential for significant economic consequences in Australia.	
<i>Grapevine ajinashika virus</i> (GAgV) [Luteoviridae: Luteovirus]	Nor known to occur	<b>Yes:</b> GAgV is symptomless in grapevines (Namba <i>et al.</i> 1991b) and this may lead to the propagation and distribution of infected propagative material. GAgV is graft transmissible (Namba <i>et al.</i> 1991b). Therefore, propagative material may provide a pathway for GAgV.	<b>Yes:</b> GAgV has established in areas with a wide range of climatic conditions (Namba <i>et al.</i> 1979) The symptomless nature of this virus may contribute to the inadvertent propagation and distribution of infected material that will help spread GAgV within Australia. Therefore, GAgV has the potential to establish and spread in Australia.	<b>Yes:</b> Grapevine ajinashika is the most important graft transmissible disease in Japan since the 1970s (Namba <i>et al.</i> 1991b). GAgV reduces the sugar content of grape berries, rendering table and wine grapes unmarketable (Namba <i>et al.</i> 1991b). Therefore, GAgV has potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine Algerian latent virus</i> (GALV) [Tombusviridae: Tombusvirus]	Not known to occur	<b>Yes:</b> GALV infections are symptomless in grapevines (Gallitelli <i>et al.</i> 1989; Brunt <i>et al.</i> 1996). This may lead to the propagation and distribution of infected propagative material. Therefore, GALV could enter Australia on propagative material.	<b>Yes:</b> GALV has established in areas with a wide range of climatic conditions (Gallitelli <i>et al.</i> 1989; Cannizzaro <i>et al.</i> 1990; Fuchs <i>et al.</i> 1994; Fujinaga <i>et al.</i> 2009). Trade of infected propagative material will help spread GALV within Australia. Therefore, GALV has the potential to establish and spread in Australia.	No: Information on the economic consequences of this virus is almost non-existent. (Gallitelli <i>et al.</i> 1989). GALV does not appear to be a threatening pathogen to grapes as infections are apparently symptomless (Gallitelli <i>et al.</i> 1989) and economic consequences are not reported. Therefore, this virus does not have the potential for significant economic consequences in Australia.	
<i>Grapevine Anatolian ringspot virus</i> (GARSV) [Secoviridae: Nepovirus]	Not known to occur	<b>Yes:</b> GARMV is associated with fanleaf degeneration/	<b>Yes:</b> GARMV has established in areas with a wide range of	<b>Yes:</b> Information on the economic consequences of this	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		decline disease (Gokalp <i>et al.</i> 2003; Oliver and Fuchs 2011). The symptoms consist of vein clearing, mottling and leaf deformation preceded by chlorotic or necrotic local lesions (Gokalp <i>et al.</i> 2003). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GARMV.	climatic conditions (Cigsar <i>et al.</i> 2002; Gokalp <i>et al.</i> 2003) and it can spread naturally in infected propagative material (Andret-Link <i>et al.</i> 2004; Oliver and Fuchs 2011). Distribution of infected propagative material will help spread GARMV within Australia. Therefore, GARMV has the potential to establish and spread in Australia.	virus is almost non-existent as it has only recently been described (Gokalp <i>et al.</i> 2003). However, as it is a part of the virus complex associated with fanleaf degeneration/decline disease (Gokalp <i>et al.</i> 2003; Oliver and Fuchs 2011), it may cause significant crop losses. Fanleaf diseases in grapevines are important diseases (Andret-Link <i>et al.</i> 2004) and cause substantial crop loss; reduced fruit quality and shortened longevity (Laimer <i>et al.</i> 2009; Oliver and Fuchs 2011). Therefore, this virus has the potential for economic consequences in Australia.	
<i>Grapevine angular mosaic-associated virus</i> (GAMaV) [Bromoviridae: Ilarvirus]	Nor known to occur	<b>Yes:</b> GAMaV naturally infects grapevine, causing angular mosaic on leaves and gradual decline and stunting of vines (Girgis <i>et al.</i> 2000, 2009). This virus is also transmitted through seed, pollen and grafting (Girgis <i>et al.</i> 2009). Therefore, propagative material provides a pathway for GAMaV.	<b>Yes:</b> GAMaV has established in areas with a wide range of climatic conditions (Girgis <i>et al.</i> 2000, 2009). It is graft transmissible (Girgis <i>et al.</i> 2009) and may therefore spread by propagative material. Multiplication and distribution of infected propagative material will help spread GAMaV within Australia. Therefore, GAMaV has the potential to establish and spread in Australia.	<b>Yes:</b> GAMaV causes a reduction in inflorescences, flower abortion, reduced berry size, gradual decline and stunting of the vine and can ultimately lead to the death of the plant (Girgis <i>et al.</i> 2009). Therefore, GAMaV has potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine asteroid mosaic associated</i>	Nor known to	<b>Yes:</b> GAMV naturally infects	<b>Yes:</b> GAMV has established	<b>Yes:</b> Plants infected with this	<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>virus</i> (GAMV) [Tymoviridae: Marafivirus]	occur	grapevines, causing leaf spot and the formation of asymmetrical leaves (Martelli and Boudon-Padieu 2006). Grapevine varieties and rootstocks infected with a Marafivirus may be symptomless (Constable and Rodoni 2011a). This may lead to the propagation and distribution of infected propagative material, suggesting that GAMV could enter Australia on propagative material.	in areas with a wide range of climatic conditions (Martelli and Boudon-Padieu 2006) and it may spread naturally in infected propagative material (Martelli and Boudon-Padieu 2006). Multiplication and distribution of infected propagative material will help spread GAMV within Australia. Therefore, GAMV has the potential to establish and spread in Australia.	virus are stunted and can be damaged quite severely (Frazier 1970). GAMV, in combination with other viruses like <i>Grapevine rupestris vein feathering virus</i> , <i>Grapevine angular mosaic-associated virus</i> or <i>Grapevine Syrah virus-1</i> , may impact grapevine health. Therefore, GAMV has the potential for economic consequences in parts of Australia.	
<i>Grapevine berry inner necrosis virus</i> (GINV) [Betaflexividae: Trichovirus]	Nor known to occur	<b>Yes:</b> GINV naturally infects grapevines resulting in poor growth (Yoshikawa <i>et al.</i> 1997). The virus causes a reduction in vigour, late sprouting, inner necrosis of shoots, and mosaic patterns on leaves (Yoshikawa <i>et al.</i> 1997). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GINV.	<b>Yes:</b> GINV has established in areas with a wide range of climatic conditions (Terai <i>et al.</i> 1993; Yoshikawa <i>et al.</i> 1997) and it may spread naturally in infected propagative material (Nishijima <i>et al.</i> 2000). Multiplication and distribution of infected propagative material and its vector <i>Colomerus vitis</i> (Kunugi <i>et al.</i> 2000) will help spread GINV within Australia. Therefore, GINV has the potential to establish and spread in Australia.	<b>Yes:</b> In Japan, GINV is considered to be one of the most important viruses of certain varieties of grapevines (Martelli and Boudon-Padieu 2006). The virus has a significant impact on the health of the grapevines, resulting in poor growth and necrosis of berries (Yoshikawa <i>et al.</i> 1997). Therefore, this virus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Grapevine Bulgarian latent virus</i> (GBLV) [Secoviridae: Nepovirus]	Nor known to occur	<b>Yes:</b> GBLV is associated with fanleaf degeneration/ decline disease (Oliver and Fuchs	<b>Yes:</b> GBLV has established in areas with a wide range of climatic conditions (Martelli <i>et</i>	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent.	<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		2011) and is seed-borne in grapes (Richardson 1990). GBLV infections are symptomless (Martelli <i>et al.</i> 1977) and this may lead to the propagation and distribution of infected propagative material. Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GBLV.	<i>al.</i> 1978; Uyemoto <i>et al.</i> 1977; Sequeira and Mendonça 1992) and it can spread naturally in infected propagative material. The symptomless nature of this virus may contribute to the inadvertent propagation and distribution of infected material that will help spread GBLV within Australia. Therefore, GBLV has the potential to establish and spread in Australia.	However, as it is a part of the virus complex associated with fanleaf degeneration/decline disease (Oliver and Fuchs 2011), it may cause significant crop losses. A New York isolate caused delayed bud break and differential elongation of bud shoots and smaller fruit clusters with many aborted berries (Uyemoto <i>et al.</i> 1977). Therefore, this virus has potential for economic consequences in Australia.	
<i>Grapevine chrome mosaic virus</i> (GCMV) [Secoviridae: Nepovirus]	Nor known to occur	<b>Yes:</b> GCMV is associated with fanleaf degeneration/ decline disease (Oliver and Fuchs 2011). GCMV is seed-borne in grapevines (Lazar <i>et al.</i> 1990; Lehoczy 1991) and causes chrome yellow or white discolouration of the leaves with leaf and cane deformations (Martelli <i>et al.</i> 1970; Dimou <i>et al.</i> 1994). However, symptomless infection may occur (Martelli and Boudon-Padiou 2006). Therefore, propagative material provides a pathway for GCMV.	<b>Yes:</b> GCMV has established in areas with a wide range of climatic conditions (Uyemoto <i>et al.</i> 2009) and it can spread naturally in infected propagative material (Dimou <i>et al.</i> 1994). The symptomless nature of this virus may contribute to the inadvertent propagation and distribution of infected material that will help spread GCMV within Australia. Therefore, GCMV has the potential to establish and spread in Australia.	<b>Yes:</b> Infected vines show a remarkable reduction in vigour and progressive decline leading to low fruit yield (Martelli <i>et al.</i> 1970) and eventual death of the plants 5–6 years after infection (Martelli <i>et al.</i> 1970; Pozsár <i>et al.</i> 1969). This pathogen can also reduce chlorophyll production and CO <sub>2</sub> fixation (Pozsár <i>et al.</i> 1969), causing grapevine yield to decline by 66% and reducing grape sugar content (Lehoczy and Tasnády 1971). Therefore, GCMV has the potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine deformation virus</i> (GDefV) [Secoviridae: Nepovirus]	Nor known to occur	<b>Yes:</b> GDefV is associated with fanleaf-like symptoms	<b>Yes:</b> GDefV has established in areas with a wide range of	<b>Yes:</b> Information on the economic consequences of this	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Martelli and Boudon-Padieu 2006). GDefV does not always display easily detectable symptoms (Cigsar <i>et al.</i> 2003). This virus can spread naturally in infected propagative material (Cigsar <i>et al.</i> 2003). Therefore, propagative material provides a pathway for GDefV.	climatic conditions (Cigsar <i>et al.</i> 2003) and it can spread naturally in infected propagative material (Cigsar <i>et al.</i> 2003). Multiplication and distribution of infected propagative material will help spread GDefV within Australia. Therefore, GDefV has the potential to establish and spread in Australia.	virus is almost non-existent. However, as it is a part of the virus complex associated with fanleaf degeneration/decline disease (Oliver and Fuchs 2011), it may cause significant crop losses. Affected plants have depressed growth and straggly fruit clusters (Cigsar <i>et al.</i> 2003). This may reduce fruit yield and quality. Therefore, GCMV has the potential for economic consequences in parts of Australia.	
<i>Grapevine fanleaf virus</i> (GFLV) [Secoviridae: Nepovirus]	Not known to occur <sup>31</sup>	<b>Yes:</b> GFLV is associated with fanleaf (Martelli and Boudon-Padieu 2006) and is seed-borne in grapes (Richardson 1990). GFLV causes a variety of symptoms that differ in type and severity (Martelli 1993). Typical symptoms include distorted leaves, chlorotic mottling, yellow mosaic and cane malformation (Raski <i>et al.</i> 1983). However, leaf and cane malformation symptoms may not always be prominent	<b>Yes:</b> GFLV has established in areas with a wide range of climatic conditions (Andret-Link <i>et al.</i> 2004) and it can spread naturally in infected propagative material. Multiplication and distribution of infected propagative material will help spread GFLV within Australia. Therefore, GFLV has the potential to establish and spread in Australia.	<b>Yes:</b> GFLV is associated with fanleaf degeneration, causing substantial crop losses, reduced fruit quality and shortened longevity of vineyards (Andret-Link <i>et al.</i> 2004). Crop losses depend on the virulence of the virus isolate, the susceptibility of the cultivar and environmental factors (Bovey <i>et al.</i> 1990). GFLV also reduces fruit quality, with a substantial decrease in sugar content and titratable acidity (Andret-Link <i>et al.</i> 2004).	<b>Yes</b>

<sup>31</sup> Grapevine fanleaf virus (GFLV) has been reported from South Australia and Victoria (Taylor 1962; Taylor and Hewitt 1964; Meagher *et al.* 1976; Cirami *et al.* 1988). In South Australia, GFLV affected only a small number of grapevines and occurred in the absence of the vector (Cirami *et al.* 1988); and in Victoria, GFLV and its vector occurred only in the Rutherglen district and quarantine restriction (due to *Phylloxera*) prevented their movement to other regions (Krake *et al.* 1999). In recent years, there have been no reports of fanleaf disease in South Australia and Victoria (Constable *et al.* 2010). Specific strains of GFLV cause fanleaf, yellow mosaic and veinbanding diseases. Some isolates are associated with leaf enation, bark pitting, wood pitting and flat trunk diseases (Hewitt *et al.* 1970).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Martelli 1993). Therefore, propagative material provides a pathway for GFLV.		Therefore, GFLV has the potential for economic consequences in parts of Australia.	
<i>Grapevine fleck virus</i> (GFKV) [Tymoviridae: Maculavirus]	Yes (Habili <i>et al.</i> 2003)	Assessment not required			
<i>Grapevine labile rod-shaped virus</i> (GLRSV) [Unassigned: Unassigned]	Not known to occur	<b>Yes:</b> This virus is reported to infect grapevines (Faggioli <i>et al.</i> 1992). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GLRSV.	<b>Yes:</b> This virus is reported to occur in Italy (Faggioli <i>et al.</i> 1992). There are similar climates in Australia that would be suitable for its establishment and spread. The symptomless nature of this virus in grapevine (Faggioli <i>et al.</i> 1992) would facilitate its inadvertent distribution with nursery stock. Multiplication and distribution of infected propagative material will help spread GLRSV within Australia. Therefore, GLRSV has the potential to establish and spread in Australia.	No: There is little information available on the economic importance of this species. It is symptomless in grapevines (Faggioli <i>et al.</i> 1992) and is not reported to infect other hosts. It is not reported to play a significant role in causing grapevine diseases (Constable <i>et al.</i> 2010). Therefore, it is unlikely to cause significant economic consequences in Australia.	
<i>Grapevine leafroll associated virus 1</i> (GLRaV-1) [Closteroviridae: Ampelovirus]	Yes (Habili <i>et al.</i> 2007)	Assessment not required			
<i>Grapevine leafroll associated virus 2</i> (GLRaV-2) [Closteroviridae: Closterovirus]	Yes (Constable <i>et al.</i> 2010)	Assessment not required			
<i>Grapevine leafroll associated virus 3</i> (GLRaV-3) [Closteroviridae: Ampelovirus]	Yes (Habili and Symons 2000)	Assessment not required			
<i>Grapevine leafroll associated virus 4</i>	Yes (Constable <i>et</i>	Assessment not required			

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
(GLRaV-4) [Closteroviridae: Ampelovirus]	<i>al.</i> 2010)				
<i>Grapevine leafroll associated virus 5</i> (GLRaV-5) [Closteroviridae: Ampelovirus]	Yes (Constable <i>et al.</i> 2010)	Assessment not required			
<i>Grapevine leafroll associated virus 6</i> (GLRaV-6) [Closteroviridae: Ampelovirus] <sup>32</sup>	Not known to occur	<b>Yes:</b> GLRaVs colonize and reproduce in the grapevine phloem tissue (Martinson <i>et al.</i> 2008) and mixed infections of GLRaV are common (Hu <i>et al.</i> 1990; Zimmerman <i>et al.</i> 1990). Symptoms are not expressed on all infected vines (Fuchs 2007; Martinson <i>et al.</i> 2008). This may lead to the propagation and distribution of infected propagative material, suggesting that GLRaVs could enter Australia on propagative material.	<b>Yes:</b> GLRaVs have established in areas with a wide range of climatic conditions (Cigsar <i>et al.</i> 2002; Kuniyuki <i>et al.</i> 2006; Martinson <i>et al.</i> 2008; Eddin <i>et al.</i> 2008; Mahfoudhi <i>et al.</i> 2009) and spread by propagative material (Weber <i>et al.</i> 1993). Distribution of infected propagative material will help spread GLRaVs within Australia. Therefore, GLRaVs have the potential to establish and spread in Australia.	<b>Yes:</b> GLRaVs pose a significant threat to the grape industry through yield reduction, reduced fruit quality and the need to introduce control measures such as replanting vineyards (Maliogka <i>et al.</i> 2008a). Infected vines often have fewer clusters, lower yield (up to 30-50%) and delayed fruit ripening (Martinson <i>et al.</i> 2008). Therefore, GLRaVs have the potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine leafroll associated virus 7</i> (GLRaV-7) [Closteroviridae: Unassigned]	Not known to occur				<b>Yes</b>
<i>Grapevine leafroll associated virus 9</i> (GLRaV-9) [Closteroviridae: Ampelovirus]	Yes (Habibi <i>et al.</i> 2003)	Assessment not required			
<i>Grapevine leafroll associated virus 10</i> (GLRaV-10) [Closteroviridae: Ampelovirus] <sup>33</sup>	Not known to occur	<b>Yes:</b> GLRaVs colonize and reproduce in the grapevine phloem tissue (Martinson <i>et</i>	<b>Yes:</b> GLRaVs have established in areas with a wide range of climatic	<b>Yes:</b> GLRaVs pose a significant threat to the grape industry through yield reductions, reduced	<b>Yes</b>

<sup>32</sup> The grapevine leafroll-associated viruses (GLRaVs) are a group of viruses (at least 9) that cause similar symptoms in infected grapevines (Martinson *et al.* 2008). GLRaVs most likely originated in the Eastern Mediterranean region and co-evolved with grapevines, later spreading throughout the world by the movement of infected vines and cuttings (Weber *et al.* 1993). Currently the GLRaV-4, -5, -6 and -9 are considered distinct Ampelovirus species. However based on their genome structure, serological relationships and biology there is a suggestion that the taxonomy will be contracted and that these GLRaV species along with GLRaV-Pr and -De will be considered strains of one species (Martelli 2009).

<sup>33</sup> GLRaV-De is referred to as GLRaV-10 (Maliogka *et al.* 2008a).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Grapevine leafroll associated virus 11</i> (GLRaV-11) [Closteroviridae: Ampelovirus] <sup>34</sup>	Not known to occur	<i>al.</i> 2008) and mixed infections of GLRaV are common (Hu <i>et al.</i> 1990; Zimmerman <i>et al.</i> 1990). Symptoms are not expressed on all infected vines (Fuchs 2007; Martinson <i>et al.</i> 2008). This may lead to the propagation and distribution of infected propagative material, suggesting that GLRaVs could enter Australia on propagative material.	conditions (Cigsar <i>et al.</i> 2002; Kuniyuki <i>et al.</i> 2006; Martinson <i>et al.</i> 2008; Mahfoudhi <i>et al.</i> 2009) and are spread by propagative material (Weber <i>et al.</i> 1993). Distribution of infected propagative material will help spread GLRaVs within Australia. Therefore, GLRaVs have the potential to establish and spread in Australia.	fruit quality and the need to introduce control measures such as replanting vineyards (Maliogka <i>et al.</i> 2008a). Infected vines often have fewer clusters, lower yield (up to 30-50% yield reduction) and delayed fruit ripening (Martinson <i>et al.</i> 2008). Therefore, GLRaVs have the potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine line pattern virus</i> (GLPV) [Bromoviridae: Ilarvirus]	Not known to occur	<b>Yes:</b> GLPV naturally infects grapevines (Martelli and Boudon-Padieu 2006). GLPV is seed-borne in grapevines (Lehoczy <i>et al.</i> 1992). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GLPV.	<b>Yes:</b> GLPV has established in areas with a wide range of climatic conditions and spreads by propagative material (Martelli and Boudon-Padieu 2006). Distribution of infected propagative material and seed will help spread GLPV within Australia. Therefore, GLPV has the potential to establish and spread in Australia.	<b>Yes:</b> GLPV is known to affect vine vigour and yield is progressively reduced (Martelli 1993). Infected vines show small yellow spots and flecks on the leaf margins (Martelli 1993). Therefore, GLPV has the potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine Pinot gris virus</i> (PGPV) [Betaflexiviridae: Trichovirus]	Not known to occur	<b>Yes:</b> This recently discovered virus is associated with chlorotic mottling, puckering, leaf deformation and berry necrosis in grapevines	<b>Yes:</b> Although this virus has only recently been described, it has been reported to occur in a range of climates (Giampetruzzi <i>et al.</i> 2012;	<b>Yes:</b> There is little information available on the economic consequences of this virus as it has only recently been described. However, in Italy it	<b>Yes</b>

<sup>34</sup> GLRaV-Pr is referred to as GLRaV-11 (Maliogka *et al.* 2008a).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Giampetruzzi <i>et al.</i> 2012; Cho <i>et al.</i> 2013). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GPGV.	Cho <i>et al.</i> 2013). Closely related viruses are transmitted by mites (Giampetruzzi <i>et al.</i> 2012). Distribution of infected propagative material will help spread GPGV within Australia. Therefore, GPGV has the potential to establish and spread in Australia.	appears to cause chlorotic mottling, puckering and leaf deformation in grapevines (Giampetruzzi <i>et al.</i> 2012). In Korea, this virus is reported to cause inner necrosis of grape berries with an incidence of 1.7% in cv. Tamnara (Cho <i>et al.</i> 2013). Therefore, GPGV has the potential for economic consequences in Australia.	
<i>Grapevine red blotch-associated virus</i> (GRBaV) [Geminiviridae: Not assigned] (synonym <i>Grapevine cabernet franc-associated virus</i> )	Not known to occur	<b>Yes:</b> GRBaV infects rootstock and grafted scions of multiple grapevine varieties (Stamp and Wei 2013). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GRBaV.	<b>Yes:</b> GRBaV appears to be widespread in California and other wine growing regions of the United States (Stamp and Wei 2013). This virus is graft transmissible (Stamp and Wei 2013). It is also thought to be vectored by leafhoppers and aphids as it is related to geminiviruses, but evidence is lacking due to the recent discovery of this virus (Stamp and Wei 2013). Distribution of infected propagative material will help spread GRBaV within Australia. Therefore, GRBaV has the potential to establish and spread in Australia.	<b>Yes:</b> This virus causes red blotch disease symptoms, similar to grapevine leafroll, including irregular blotchy leaves and red veins (Al-Rwahnih <i>et al.</i> 2012a; Stamp and Wei 2013). As this virus has only recently been discovered, its affect on yield has not been determined. However, this virus has been implicated in reducing the quality of grape berries by lowering its sugar content (Al-Rwahnih <i>et al.</i> 2012a; Stamp and Wei 2013). It may also retard maturation (Zwadlo 2012). Therefore, this virus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Grapevine red globe virus</i> (GRGV) [Tymoviridae: Maculavirus]	Not known to occur	<b>Yes:</b> GRGV is part of the fleck complex of grapevines (Martelli and Boudon-Padieu 2006) causing latent or semi-latent infections in <i>Vitis</i>	<b>Yes:</b> GRGV has established in areas with a wide range of climatic conditions (Martelli and Boudon-Padieu 2006) and may spread naturally with	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent. However, as it is a part of the fleck complex (Martelli and	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		<i>vinifera</i> and most American <i>Vitis</i> species and rootstock hybrids (Martelli and Boudon-Padieu 2006). This may lead to the propagation and distribution of infected propagative material, suggesting that GRGV could enter Australia on propagative material.	propagative material. Distribution of infected propagative material will help spread GRGV within Australia. Therefore, GRGV has the potential to establish and spread in Australia.	Boudon-Padieu 2006), it may cause significant crop losses. Adverse effects on vine vigour and rooting ability of root stocks have been reported as a result of fleck complex (Martelli and Boudon-Padieu 2006). Therefore, GRGV has the potential for economic consequences in parts of Australia.	
<i>Grapevine rootstock stem lesion closterovirus</i> (GRSLaV = strain of GLRaV-2)	Yes (Constable and Drew 2004)	Assessment not required			
<i>Grapevine rupestris stem pitting associated virus</i> (GRSPaV) [Betaflexiviridae: Foveavirus]	Yes (Habibi and Symons 2000)	Assessment not required			
<i>Grapevine rupestris vein feathering virus</i> (GRVFV) [Tymoviridae: Marafivirus]	Not known to occur	<b>Yes:</b> GRVFV in association with other viruses causes grapevine fleck complex or Syrah Decline (Al Rwahnih <i>et al.</i> 2009; Uyemoto <i>et al.</i> 2009). In the absence of other viruses, GRVFV induces mild chlorosis of primary and secondary leaf veins (Uyemoto <i>et al.</i> 2009). Grapevine infected with a Marafivirus may be symptomless (Constable and Rodoni 2011a). Therefore, propagative material provides a pathway for GRVFV.	<b>Yes:</b> GRVFV has established in areas with a wide range of climatic conditions (Al Rwahnih <i>et al.</i> 2009; Uyemoto <i>et al.</i> 2009) and spread by propagative material (Martelli and Boudon-Padieu 2006; Constable and Rodoni 2011a). Distribution of infected propagative material will help spread GRVFV within Australia. Therefore, GRVFV has the potential to establish and spread in Australia.	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent. However, this virus is associated with Syrah decline, which causes leaf reddening and scorching, swelling of the graft union, superficial cracking and pitting of woody tissue, stem necrosis, and the eventual death of the vines (Al-Rwahnih <i>et al.</i> 2009). Therefore, GRVFV has the potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine stunt virus</i> (GSV)	Not known to	<b>Yes:</b> This virus infects	<b>Yes:</b> This virus occurs on	No: GSV causes leaf rolling and	



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
[Unassigned: Unassigned]	occur	grapevines in Japan (Martelli 1993). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GSV.	grapevine in Japan and is transmitted by <i>Aboridia apicalis</i> (Martelli 1993; Büchen-Osmond 2006). Multiplication and distribution of infected propagative material will help spread GRV-FV within Australia. Therefore, GRV-FV has the potential to establish and spread in Australia.	stunting of grapevines, although signs and symptoms vary seasonally (Büchen-Osmond 2006). Although this virus has been described since 1993, significant economic consequences have not been reported. Therefore, GSV is unlikely to have significant economic consequences in Australia.	
<i>Grapevine syrah virus I</i> (GSyV-I) [Tymoviridae: Marafivirus]	Not known to occur	<b>Yes:</b> GSyV-I, in association with other viruses, causes Syrah decline disease (Al-Rwahnih <i>et al.</i> 2009). Grapevine infected with a Marafivirus may be symptomless (Constable and Rodoni 2011a). Therefore, propagative material provides a pathway for GSyV-I.	<b>Yes:</b> GSyV-I has established in areas with a wide range of climatic conditions (Al-Rwahnih <i>et al.</i> 2009; Engel <i>et al.</i> 2010) and spreads by propagative material (Engel <i>et al.</i> 2010; Constable and Rodoni 2011a). Distribution of infected propagative material will help spread GSyV-I within Australia. Therefore, GSyV-I has the potential to establish and spread in Australia.	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent. However, this virus is associated with Syrah decline, which causes leaf reddening and scorching, swelling of the graft union, superficial cracking and pitting of woody tissue, stem necrosis and eventual death of the vines (Al-Rwahnih <i>et al.</i> 2009). Therefore, GSyV-I has potential for economic consequences in parts of Australia.	<b>Yes</b>
<i>Grapevine Tunisian ringspot virus</i> (GTRSV) [Secoviridae: Nepovirus]	Not known to occur	<b>Yes:</b> GTRSV is found in vines with mild fanleaf-like symptoms (Mahfoudhi <i>et al.</i> 1998). Symptoms include mild mottling and leaf deformation (Ouertani <i>et al.</i> 1992). Viruses, as a rule, infect host plants systemically and all plant parts, including parts	<b>Yes:</b> GTRSV has established in areas with a wide range of climatic conditions (Ouertani <i>et al.</i> 1992) and it can spread in infected propagative material. Multiplication and distribution of infected propagative material will help spread GTRSV within	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent. However, as a part of virus complex associated with fanleaf degeneration/decline disease (Oliver and Fuchs 2011), it may cause significant crop losses. Affected plants have depressed	<b>Yes</b>



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GTRSV.	Australia. Therefore, GTRSV has the potential to establish and spread in Australia.	growth and straggly fruit clusters (Cigsar <i>et al.</i> 2003). This may reduce fruit yield and quality. Therefore, GCMV has potential for economic consequences in parts of Australia.	
<i>Grapevine vein clearing virus</i> (GVCV) [Caulimoviridae: Badnavirus]	Not known to occur	<b>Yes:</b> This virus is reported to cause vein clearing and plant decline in grapevines (Zhang <i>et al.</i> 2011). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GVCV.	Yes: This virus occurs in a range of climates in the United States (Zhang <i>et al.</i> 2011). There is little information available on the transmission of this virus due to its recent discovery, but transmission by mealybugs and whiteflies is suspected (Zhang <i>et al.</i> 2011). This virus can also be transmitted by vegetative propagation of infected source vines (Zhang <i>et al.</i> 2011). Multiplication and distribution of infected propagative material will help spread GVCV within Australia. Therefore, GVCV has the potential to establish and spread in Australia.	No: There is little information relating to the economic importance of this virus. Although it is reported to cause vein clearing, short internodes and a decline in grapevine vigour (Zhang <i>et al.</i> 2011), yield loss and economic consequences are not reported. Therefore, this virus is unlikely to cause significant economic consequences in Australia.	
<i>Grapevine virus A</i> (GVA) [Betaflexiviridae: Vitivirus]	Yes (Habili and Symons 2000)	Assessment not required			
<i>Grapevine virus B virus</i> (GVB) [Betaflexiviridae: Vitivirus]	Yes (Habili 2009)	Assessment not required			
<i>Grapevine virus B</i> (GVB) (strains associated with grapevine corky bark) [Betaflexiviridae: Vitivirus]	Not known to occur	<b>Yes:</b> This phloem limited virus associated with grapevine corky bark is latent (Golino 1993; Abdallah <i>et al.</i> 2003) or	<b>Yes:</b> GVB strains associated with grapevine corky bark have established in areas with a wide range of climatic	<b>Yes:</b> GVB strains associated with grapevine corky bark are of major importance to viticulture worldwide (Constable and	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		produces a mild reduction in plant vigour (Namba <i>et al.</i> 1991a). This may lead to the inadvertent propagation and distribution of infected propagative material. Therefore, propagative material may provide a pathway for GVB strains associated with corky bark.	conditions (Namba <i>et al.</i> 1991a; Abdallah <i>et al.</i> 2003) and it can spread naturally in infected propagative material (Abdallah <i>et al.</i> 2003). Multiplication and distribution of infected propagative material will help spread this virus within Australia. Therefore, this virus has the potential to establish and spread in Australia.	Rodoni 2011b). This virus is associated with grapevine degeneration where grapevine yield is decreased by 66% and the grapes have reduced sugar content (Lehoczky and Tasnady 1971). Therefore, GVB strains associated with grapevine corky bark have the potential for economic consequences in parts of Australia.	
<i>Grapevine virus C</i> (GVC) (strain of GLRaV-2) <sup>35</sup> [Betaflexiviridae: Vitivirus]	Yes (Constable <i>et al.</i> 2010)	Assessment not required			
<i>Grapevine virus D</i> (GVD) [Betaflexiviridae: Vitivirus]	Yes (Habibi pers. comm. 2009)	Assessment not required			
<i>Grapevine virus E</i> (GVE) [Betaflexiviridae: Vitivirus]	Not known to occur	<b>Yes:</b> This virus is associated with grapevine causing typical Shiraz disease symptoms including canes lacking lignifications, delayed leaf fall and reduced vigour (Coetzee <i>et al.</i> 2010). Canes may also not show symptoms and this may lead to the inadvertent propagation and distribution of infected propagative material. Therefore, propagative material may provide a pathway for GVE.	<b>Yes:</b> GVE has established in areas with a wide range of climatic conditions (Nakaune <i>et al.</i> 2008; Coetzee <i>et al.</i> 2010) and it may spread naturally in infected propagative material. Propagation and distribution of infected material will help spread GVE within Australia. Therefore, GVE has the potential to establish and spread in Australia.	<b>Yes:</b> Vitiviruses may display delayed bud burst, and thick, rough bark with an enlarged scion trunk (Uyemoto <i>et al.</i> 2009). Information on the economic consequences of this virus is almost non-existent. However, as it is a part of the virus complex associated with rugose wood (Martelli <i>et al.</i> 2007), it may cause significant crop losses. Therefore, GVE has the potential for economic consequences in parts of Australia.	<b>Yes</b>

<sup>35</sup> GVC was considered to be related to the Vitiviruses but it is now considered to be a strain of GLRaV-2 (Masri *et al.* 2006). GLRaV-2 is present in Australia (Constable *et al.* 2010)

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Grapevine virus F</i> (GVF) [Betaflexiviridae: Vitivirus]	Not known to occur	<b>Yes:</b> GVF infects grapevines, including Cabernet Sauvignon plants (Al-Rwahnih <i>et al.</i> 2012b). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for GVF.	<b>Yes:</b> There is little information on the establishment and spread of this species as it was only described in 2012. However, propagation and distribution of infected material will help spread GVF within Australia. Therefore, GVF has the potential to establish and spread in Australia.	<b>Yes:</b> There is little information on the economic consequences of this species as it has only been recently described. However, bioassays of this virus caused grapevine death in 1–2 years (Al-Rwahnih <i>et al.</i> 2012b). Vitiviruses infecting grapevines are associated with rugose woody complex, which includes several important diseases that produce woody cylinder modifications (Al-Rwahnih <i>et al.</i> 2012b). Therefore, GVF has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Peach rosette mosaic virus</i> (PRMV) [Secoviridae: Nepovirus]	Not known to occur	<b>Yes:</b> PRMV is seed-borne and soil-borne (Richardson 1990). It is associated with symptoms similar to those of fanleaf degeneration and decline (Martelli and Boudon-Padieu 2006). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for PRMV.	<b>Yes:</b> PRMV has established in areas with a wide range of climatic conditions (Uyemoto <i>et al.</i> 2009) and it can spread naturally in infected propagative material (Martelli and Boudon-Padieu 2006). Propagation and distribution of infected material will help spread PRMV within Australia. Therefore, PRMV has the potential to establish and spread in Australia.	<b>Yes:</b> PRMV causes delayed bud burst, small sized berries, stunted vines and a progressive decline in plant health, which can lead to grapevine death (Martelli and Boudon-Padieu 2006). Crop losses of up to 60% and death of susceptible <i>Vitis labrusca</i> cultivars and a number of American-French hybrids have been recorded (Martelli and Boudon-Padieu 2006). Therefore, this virus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Petunia asteroid mosaic virus</i> (PeAMV) [Tombusviridae: Tombusvirus]	Not known to occur	<b>Yes:</b> PeAMV is a soil-borne virus and infects plant systemically via roots (Kegler	<b>Yes:</b> PeAMV has established in areas with a wide range of climatic conditions (Bercks	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent.	<b>Yes</b>

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		and Kontzog 1990; Lovisolo 1990). The infections may be latent (Kegler and Kontzog 1990). This may lead to the inadvertent propagation and distribution of infected propagative material. Therefore, propagative material provides a pathway for PeAMV.	1967; Novák and Lanzová 1976; Smith <i>et al.</i> 1988; Koenig <i>et al.</i> 1989; Martelli 1993; Constable <i>et al.</i> 2010) and it can spread naturally in infected propagative material. Propagation and distribution of infected material will help spread PeAMV within Australia. Therefore, PeAMV has the potential to establish and spread in Australia.	However, PeAMV generally occur in mixed infections (Constable <i>et al.</i> 2010). PeAMV is associated with a serious disease—viral necrosis of sweet cherry—that causes heavy damage due to canker-like deformations on the shoots as well as bark splits, necrosis of leaf mid-veins and misshapen fruits with necrotic spots (Pfeilstetter <i>et al.</i> 1992). Therefore, this virus has the potential for economic consequences in Australia.	
<i>Potato virus X</i> (PVX)—grapevine strains [Alphaflexiviridae: Potexvirus]	Not known to occur <sup>36</sup>	<b>Yes:</b> PVX is reported to infect grapevine in Tunisia and Italy (Chabbouh <i>et al.</i> 1993; Martelli 2012). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for PVX.	<b>Yes:</b> PVX grapevine strains occur in Tunisia and Italy (Chabbouh <i>et al.</i> 1993). There are similar climates in parts of Australia that will be suitable for its establishment and spread. Other PVX strains are already distributed in Australia (Büchen-Osmond <i>et al.</i> 1988). Propagation and distribution of infected material will help spread PVX grapevine strains within Australia. Therefore, PVX has the potential to establish and spread in Australia.	No: Grapevines infected with PVX are asymptomatic or at most show faint chlorotic spots or veinbanding (Chabbouh <i>et al.</i> 1993). Although other PVX strains can cause significant economic consequences on some hosts, PVX strains infecting grapevine are not reported to cause significant economic consequences. Therefore, this virus does not have the potential for significant economic consequences in Australia.	

<sup>36</sup> Although PVX is known to occur in Australia (Büchen-Osmond *et al.* 1988), it is not reported to infect grapevine.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Raspberry bushy dwarf virus</i> (RBDV) [Unassigned: Ideaeovirus]	Yes (McGregor <i>et al.</i> 1996)	Assessment not required			
<i>Raspberry ringspot virus</i> (RpRSV) grapevine strain <sup>37</sup> [Secoviridae: Nepovirus]	Not known to occur	<b>Yes:</b> RpRSV grapevine strain causes symptoms similar to those of fanleaf degeneration disease (Stellmach and Querfurth 1978; Wetzel <i>et al.</i> 2006, Wetzel and Kraczel 2007). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for RpRSV grapevine strain.	<b>Yes:</b> RpRSV grapevine strain has established in areas with a wide range of climatic conditions (Martelli and Boudon-Padieu 2006; Wetzel <i>et al.</i> 2006) and it can spread naturally in infected propagative material. Propagation and distribution of infected material will help spread RpRSV within Australia. Therefore, RpRSV grapevine strain has the potential to establish and spread in Australia.	<b>Yes:</b> RpRSV is a causal agent of grapevine fanleaf disease, one of the most widespread and damaging diseases of grapevine (Wetzel <i>et al.</i> 2006). Crop losses caused by RpRSV grapevine strain can be higher than 30% (Martelli and Boudon-Padieu 2006). Therefore, this virus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Sowbane mosaic virus</i> (SoMV) – grape infecting strain [Unassigned: Sobemovirus]	Not known to occur <sup>38</sup>	Yes: SoMV grape infecting strain may be latent in naturally infected grapevines (Bercks and Querfurth 1969). This may lead to the inadvertent propagation and distribution of infected propagative material. Therefore, propagative	<b>Yes:</b> SoMV grape infecting strain has established in areas with a wide range of climatic conditions (Bercks and Querfurth 1969; Jankulova 1972; Pozdena <i>et al.</i> 1977) and it can spread naturally in infected propagative material.	<b>Yes:</b> Information on the economic consequences of this virus is almost non-existent. However, SoMV grape infecting strain is considered of quarantine significance by some trading partners. Presence of SoMV grape infecting strain in Australia would impact upon Australia's	<b>Yes</b>

<sup>37</sup> The grapevine strain of RsRSV is serologically very distantly related to the main serotypes Scottish and English. These differences strongly suggest that the grapevine infecting RsRSV may be a different viral species (Jones *et al.* 1994; Ebel *et al.* 2003). The type strain is transmitted by *Longidorus macrosoma* whereas grapevine strain is transmitted by *Paralongidorus maximus* (Jones *et al.* 1994). Two strains of different virulence occur (Ebel *et al.* 2003): Raspberry ringspot virus – cherry isolate (RpRSV - ch) in Germany and Raspberry ringspot virus – RAC815 isolate (RpRSV-RAC815) in Switzerland; both have also been recorded from grapevines (Wetzel *et al.* 2006).

<sup>38</sup> Sowbane mosaic virus (SoMV) naturally occurs on *Atriplex suberecta*, *Chenopodium album* and *Chenopodium trigonon* in Australia (Teakle 1968; Guy 1982). However, this virus has not been recorded on grapevines in Australia (Constable and Drew 2004; Constable *et al.* 2010).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		material provides a pathway for SoMV grape infecting strain.	Therefore, SoMV grape infecting strain has the potential to establish and spread in Australia.	ability to access overseas markets. Therefore, SoMV grape infecting strain has potential for economic consequences in parts of Australia.	
<i>Strawberry latent ringspot virus</i> (SLRSV) [Comoviridae: Unassigned]	Not known to occur <sup>39</sup>	<b>Yes:</b> SLRSV is associated with symptoms similar to those of fanleaf degeneration (Martelli and Walter 1993; Constable <i>et al.</i> 2010; Oliver and Fuchs 2011). SLRSV infections are generally latent, but SLRSV may induce leaf deformity, chlorotic mottling on leaf, leaf roll symptoms, reddish discoloration of the tip of the spring shoots and reduced or stunted growth (Savino <i>et al.</i> 1987; Martelli and Walter 1993). Therefore, propagative material provides a pathway for SLRSV.	<b>Yes:</b> SLRSV has established in areas with a wide range of climatic conditions (Murant 1983; EPPO 2010a) and it can spread naturally in infected propagative material (Savino <i>et al.</i> 1987; Holleínova <i>et al.</i> 2009). The symptomless nature of this virus may contribute to the inadvertent propagation and distribution of infected material that will help spread SLRSV within Australia. Therefore, SLRSV has the potential to establish and spread in Australia.	<b>Yes:</b> SLRSV is an economically important virus due to its extensive host range and the yield losses it can cause (Tzanetakis <i>et al.</i> 2006). SLRSV occurrence varies from 3% to 18% in grapevines (Akbas and Erdiller 1993; Komínek 2008; Holleínovà <i>et al.</i> 2009). Heavy yield losses (up to 80% of the crop) are associated with SLRSV infections in grapevine (Rudel 1985; Martelli and Walter 1993). Therefore, this virus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Tobacco mosaic virus</i> (TMV) [Unassigned: Tobamovirus]	Yes (Randles 1986)	Assessment not required			

<sup>39</sup> In Australia, SLRSV has only once been reported from Rhubarb in South Australia (Cook and Dubé 1989). As there have been no further reports of this virus in Australia, it is considered to be eradicated. The natural vector of SLRSV is also absent from Australia.

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
<i>Tobacco necrosis virus</i> (TNV) grape strain [Tombusviridae: Necrovirus]	Not known to occur <sup>40</sup>	<b>Yes:</b> TNV grape strain causes yellowing and mottling on grapevine leaves (Cesati and Van Regenmortel 1969) and infections are systemic (Cesati and Van Regenmortel 1969). Therefore, propagative material provides a pathway for TNV grape strain.	<b>Yes:</b> TNV grape strain has established in areas with a wide range of climatic conditions (Cesati and Van Regenmortel 1969) and it is graft transmissible (Cesati and Van Regenmortel 1969). Propagation and distribution of infected material, and the presence of efficient vectors ( <i>Olpidium</i> species), will help spread TNV grape strain within Australia. Therefore, TNV grape strain has the potential to establish and spread in Australia.	<b>Yes:</b> Information on the economic consequences of this virus on grapevines is almost non-existent. However, in other hosts TNVs cause significant yield losses. In strawberry in the Czech Republic, TNV has caused dwarfing and leaf and root necrosis (Martin and Tzanetakis 2006). Losses as high as 50% have been recorded in tulips and glasshouse grown cucumbers (CABI 2012a). Therefore, this virus has the potential for economic consequences in Australia.	<b>Yes</b>
<i>Tobacco ringspot virus</i> (TRSV) [Secoviridae: Nepovirus]	Yes (Randles 1986)	Assessment not required			
<i>Tomato black ring virus</i> (TBRV) [Secoviridae: Nepovirus] (synonym <i>Grapevine Joannes-Seyve virus</i> [GJSV])	Not known to occur	<b>Yes:</b> TBRV naturally infects grapevines and produces chlorotic spots, rings and lines on newly infected vines, and mottling of the older leaves (Stobbs and van Schagen 1984; Walker 2006). TBRV is seed-borne in grapevines	<b>Yes:</b> TBRV has established in areas with a wide range of climatic conditions (Harper <i>et al.</i> 2010) and it can spread naturally in infected propagative material. Multiplication and distribution of infected propagative	<b>Yes:</b> Production losses caused by TBRV in grapevine are not known precisely, but they can be high (Uyemoto <i>et al.</i> 2009). Vines infected with TBSV are generally stunted with older leaves showing mottling, yellowing of leaf margins, vein	<b>Yes</b>

<sup>40</sup> The taxonomy of TNV has been revised to recognise that what was originally named TNV is actually a group of related virus species. *Tobacco necrosis virus A* (TNV-A) and *Tobacco necrosis virus D* (TNV-D) have been recognised as distinct species in the *Necrovirus* genus (Coutts *et al.* 1991; Meulewaeter *et al.* 1990), as have *Chenopodium necrosis virus* (ChNV) and *Olive mild mosaic virus* (OMMV), which were previously considered TNV isolates (Tomlinson *et al.* 1983). TNV isolates from Nebraska and Toyama (TNV-NE and TNV-Toyama) are likely to represent two new species in the genus, but have not yet been officially recognised (Saeki *et al.* 2001; Zhang *et al.* 1993). Molecular sequence data indicates that other necroviruses originally labelled '*Tobacco necrosis virus*' are likely to be confirmed as distinct species (NCBI 2010). Viruses likely to be strains of TNVs A and D have been recorded in Victoria and Queensland (Finlay and Teakle 1969; Teakle 1988). TNV Nebraska isolate and grape infecting strain has not been recorded in Australia, nor have other TNVs that have since been renamed or have not yet been formally classified (Tomlinson *et al.* 1983; Zhang *et al.* 1993; Cardoso *et al.* 2005; NCBI 2010).

Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		(Martelli 1978). Viruses, as a rule, infect host plants systemically and all plant parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for TBRV.	material, and the presence of its nematode vectors in Australia (Stirling <i>et al.</i> 1992), will help spread TBRV within Australia. Therefore, TBRV has the potential to establish and spread in Australia.	bunching, leaf deformation, and small, poorly set berries (Stobbs and van Schagen 1984). This may reduce yield and fruit quality. Yield losses of up to 20% in raspberry (Taylor <i>et al.</i> 1965) and up to 40% on artichoke (Harper <i>et al.</i> 2010) have been reported due to TBRV. TBRV is of quarantine significance for NAPPO and New Zealand (Harper <i>et al.</i> 2010). Therefore, TBRV has the potential for economic consequences in parts of Australia.	
<i>Tomato mosaic virus</i> (ToMV) [Unassigned: Tobamovirus]	Yes (PHA 2001)	Assessment not required			
<i>Tomato ringspot virus</i> (ToRSV) [Secoviridae: Nepovirus]	Not known to occur <sup>41</sup>	<b>Yes:</b> ToRSV naturally infects grapevines causing faint chlorotic mottling, small, distorted leaves, irregular, ringlike line patterns on leaves and shortened internodes (Uyemoto <i>et al.</i> 2009; Schilder 2011). ToRSV is seed-transmitted in grapes (Uyemoto 1975). Viruses, as a rule, infect host plants systemically and all plant	<b>Yes:</b> ToRSV has established in areas with a wide range of climatic conditions (EPPO 2010b) and it can spread naturally in infected propagative material (Gooding and Téliv 1970; Schilder 2011). Multiplication and distribution of infected propagative material, and the presence of nematode vectors in Australia (Stirling <i>et al.</i>	<b>Yes:</b> ToRSV is an economically important pathogen. Vines infected with ToRSV show shortened internodes, distorted leaves and sparse fruit clusters with many berries aborting (Uyemoto 1975). Infected raspberries experience a gradual decline and up to 80% of fruiting canes may be killed in the third year of infection (EPPO 2010b). TomRSV is an A2 quarantine	<b>Yes</b>

<sup>41</sup> *Tomato ring spot virus* was reported more than two decades ago in *Pentas lanceolata* (Egyptian starflower) and *Cymbidium* orchid species in South Australia (Chu *et al.* 1983; Cook and Dubé 1989). The infected plants were removed and it has not since been reported to occur in South Australia (Cartwright 2009), suggesting the virus has not spread and is probably absent from Australia.



Pest	Present within Australia	Potential to be on pathway	Potential for establishment and spread	Potential for economic consequences	Quarantine pest
		parts, including parts used for vegetative propagation, are infected (Bos 1999). Therefore, propagative material provides a pathway for ToRSV.	1992), will help spread ToRSV within Australia. Therefore, ToRSV has the potential to establish and spread in Australia.	pest for EPPO (OEPP/EPPO 1982) and has quarantine significance for the Inter-African Phytosanitary Council (IAPSC). Therefore, ToRSV has the potential for economic consequences in parts of Australia.	
<i>Tomato spotted wilt virus</i> (TSWV) [Bunyaviridae: Tospovirus]	Yes (Persely <i>et al.</i> 2006)	Assessment not required			
<b>DISEASES OF UNKNOWN AETIOLOGY</b>					
Grapevine enation disease	Present (Krake <i>et al.</i> 1999)	Assessment not required			
Grapevine vein mosaic	Present (Uyemoto <i>et al.</i> 2009)	Assessment not required			
Grapevine vein necrosis	Present (Woodham and Krake 1984)	Assessment not required			
Summer mottle	Present (Woodham and Krake 1984)	Assessment not required			

## Appendix B: Additional quarantine pest data

ARTHROPODS	
<b>Quarantine pest</b>	<i>Brevipalpus chilensis</i> Baker
<b>Synonyms</b>	
<b>Common name(s)</b>	Chilean false red mite
<b>Main hosts</b>	<i>Actinidia chinensis</i> ; <i>Ampelopsis</i> sp.; <i>Annona cherimola</i> ; <i>Antirrhinum</i> sp.; <i>Catalpa speciosa</i> ; <i>Chrysanthemum</i> sp.; <i>Citrus limon</i> ; <i>Citrus sinensis</i> ; <i>Cydonia oblonga</i> ; <i>Diospyros kaki</i> ; <i>Ficus carica</i> ; <i>Garcinia</i> sp.; <i>Jasminum angustifolium</i> ; <i>Lugustrum sinensis</i> ; <i>Malus pumila</i> ; <i>Pelagonium</i> sp.; <i>Prunus armeniaca</i> ; <i>Prunus dulcis</i> ; <i>Pyrus communis</i> ; <i>Rubus idaeus</i> ; <i>Strongylodon macrobotrys</i> ; <i>Viburnum</i> sp.; <i>Vinca</i> sp.; <i>Vitis vinifera</i> (Gonzalez 1983; Klein Koch and Waterhouse 2000; SAG/USDA 2002; CABI 2012a)
<b>Distribution</b>	Argentina, Chile
<b>Quarantine pest</b>	<i>Colomerus vitis</i> Pagenstecher - strain c
<b>Synonyms</b>	<i>Phytoptus vitis</i> Pagenstecher, <i>Eriophyes vitis</i> Pagenstecher
<b>Common name(s)</b>	Grapeleaf bud mite – leaf curl strain
<b>Main hosts</b>	<i>Diospyros</i> spp.; <i>Vitis</i> spp. (CABI 2012a)
<b>Distribution</b>	California, USA (Smith and Stafford 1948), South Africa (Schwartz 1986)
<b>Quarantine pest</b>	<i>Daktulosphaira vitifoliae</i> (Fitch 1855)
<b>Synonyms</b>	<i>Vitis vitifoliae</i> (Fitch); <i>Phylloxera vastatrix</i> Planchon; <i>Phylloxera vitifoliae</i> (Fitch)
<b>Common name(s)</b>	Grape phylloxera
<b>Main hosts</b>	<i>Vitis</i> spp. (CABI 2012a)
<b>Distribution</b>	Present in Australia but under official control. Also occurs in Algeria, Argentina, Armenia, Austria, Azerbaijan, Bermuda, Bolivia, Bosnia-Herzegovina, Brazil, Bulgaria, Canada, China, Colombia, Croatia, Czech Republic, France, Georgia, Germany, Hungary, India, Israel, Italy, Japan, Jordan, Korea, Lebanon, Luxembourg, Macedonia, Malta, Mexico, Montenegro, Morocco, Panama, Peru, Poland, Portugal, Romania, Russian Federation, Serbia, Slovakia, Slovenia, South Africa, Spain, Syria, Tunisia, Turkey, United States, Ukraine, Uruguay, Venezuela, Zimbabwe (CABI 2012a).
<b>Quarantine pest</b>	<i>Paranthrene regalis</i> Butler
<b>Synonyms</b>	<i>Paranthrene regale</i> , <i>Sciapteron regalis</i>
<b>Common name(s)</b>	grape clearwing moth
<b>Main hosts</b>	<i>Vitis vinifera</i> (grapevine) (CABI 2012a; Shao-Hua 2012)
<b>Distribution</b>	China (CABI 2012a; Shao-Hua 2012)
<b>Quarantine pest</b>	<i>Planococcus ficus</i> Signoret
<b>Synonyms</b>	<i>Dactylopius subterraneus</i> , <i>Pseudococcus vitis</i> , <i>Pseudococcus citriodes</i> , <i>Planococcus citriodes</i> , <i>Pseudococcus praetermissus</i> (Walton and Pringle 2004)
<b>Common name(s)</b>	Subterranean vine mealybug, Vine mealybug (Walton and Pringle 2004)
<b>Main hosts</b>	<i>Bambusa</i> spp.; <i>Cydonia oblonga</i> ; <i>Dahlia</i> spp.; <i>Dichrostachys glomerata</i> ; <i>Ficus benjamina</i> ; <i>Juglans</i> spp.; <i>Malus domestica</i> ; <i>Malus pumila</i> ; <i>Mangifera indica</i> ; <i>Nerium oleander</i> ; <i>Persea americana</i> ; <i>Phoenix dactylifera</i> ; <i>Platanus orientalis</i> ; <i>Prosopis farcata</i> ; <i>Salix</i> spp.; <i>Styrax officinalis</i> ; <i>Theobroma cacao</i> ; <i>Vitis vinifera</i> ; <i>Zizyphus spina-christi</i> (Ezzat and McConnel 1956; Cox 1989; Walton and Pringle 2004).
<b>Distribution</b>	Found in most grape-production areas throughout the world with particular economic importance on grapevines in Argentina, the Mediterranean region, Pakistan and South

	Africa (Ben-Dov 1994; Walton and Pringle 2004).
<b>Quarantine pest</b>	<i>Planococcus lilacinus</i> Cockerell
<b>Synonyms</b>	<i>Dactylopius crotonis</i> , <i>Planococcus citri</i> , <i>P. crotonis</i> , <i>P. lilacinus</i> , <i>P. tavabanus</i> , <i>Pseudococcus lilacinus</i> , <i>P. tavabanus</i> , <i>P. crotonis</i> , <i>P. deceptor</i> , <i>Tylococcus mauritiensis</i> (USDA 1995).
<b>Common name(s)</b>	Coffee mealybug, Cocoa mealybug
<b>Main hosts</b>	<i>P. lilacinus</i> is extremely polyphagous, feeding on tropical and sub-tropical fruit and shade trees within 35 families (Williams 1982; Cox 1989; Ben-Dov 1994). Hosts include <i>Adenophyllum</i> spp.; <i>Ailanthus</i> spp.; <i>Albizia lebbek</i> ; <i>Alphitonia incana</i> ; <i>Annona</i> spp.; <i>Apium graveolens</i> ; <i>Arachis hypogaea</i> ; <i>Asteraceae</i> ; <i>Bauhinia monandra</i> ; <i>Caianus</i> spp.; <i>Calophyllum inophyllum</i> ; <i>Cananaa oderata</i> ; <i>Castilloa elastic</i> ; <i>Citrus aurantium</i> ; <i>C. grandis</i> ; <i>Cocos nucifera</i> ; <i>Codiaeum</i> spp.; <i>Coffea canephora</i> ; <i>C. sepihiiala</i> ; <i>Cordia myxa</i> ; <i>Couroupita quianensis</i> ; <i>Dioscorea</i> spp.; <i>Dipterocarpus</i> spp.; <i>Ervthrin lithosperma</i> ; <i>E. indica</i> , <i>E. variegata</i> , <i>Euphorbia pyrifolia</i> , <i>Euqenia mespiloides</i> , <i>Ficus rubra</i> , <i>Gladiolus carmels</i> ; <i>Hibiscus rosa-sinensis</i> ; <i>Hymenaea</i> spp.; <i>Litchi</i> spp.; <i>Mallotus japonicus</i> ; <i>Mangifera indica</i> ; <i>Nicotiana tabacum</i> ; <i>Ochroma</i> sp; <i>Pandanus</i> spp.; <i>Phoenix dactylifera</i> ; <i>Pongamia pinnata</i> ; <i>Prosopis juliflora</i> ; <i>Psidium quaiava</i> ; <i>Punica granatum</i> ; <i>Tamarindus indica</i> ; <i>Tectona grandis</i> ; <i>Theobroma cacao</i> ; <i>Vitis vinifera</i> ; <i>Zizyphus juiuba</i> (Williams 1982; Cox 1989; Ben-Dov 1994; USDA 1995).
<b>Distribution</b>	Aden, Bangladesh, Borneo, Burma, Cambodia, Cocos Keeling Island, China, Comoros, Dominican Republic, El Salvador, Guyana, Haiti, India, Indonesia, Japan, Java, Madagascar, Mauritius, Papua New Guinea, Philippines, Rodriguez Island, Seychelles, Sri Lanka, Taiwan, Thailand, Vietnam, West Malayasia (USDA 1995).
<b>Quarantine pest</b>	<i>Planococcus kraunhiae</i> (Kuwana, 1902)
<b>Synonyms</b>	<i>Dactylopius kraunhiae</i> Kuwana 1902, <i>Planococcus siakwanensis</i> Borchsenius 1962, <i>Dactylopius krounhiae</i> Kuwana 1917, <i>Planococcus kraunhiae</i> Ferris 1950, <i>Pseudococcus kraunhiae</i> Fernald, 1903
<b>Common name(s)</b>	Japanese mealybug
<b>Main hosts</b>	<i>Actinidia</i> (kiwifruit), <i>Agave americana</i> (Century plant), <i>Artocarpus lanceolata</i> , <i>Broussonetia kazinoki</i> (Japanese paper mulberry), <i>Casuarina stricta</i> (she oak), <i>Citrus junos</i> (yuzu), <i>Citrus nobilis</i> (tangor), <i>Citrus paradisi</i> (grapefruit), <i>Codiaeum variegatum pictum</i> (variegated laurel), <i>Coffea arabica</i> (coffee), <i>Crinum asiaticum</i> (poison bulb), <i>Cucurbita moschata</i> (pumpkin), <i>Cydonia sinensis</i> (quince), <i>Digitaria sanguinalis</i> (crab-grass), <i>Diospyros kaki</i> (Japanese kaki), <i>Ficus carica</i> (fig), <i>Gardenia jasminoides</i> (common gardenia), <i>Ilex</i> (holly), <i>Magnolia grandiflora</i> (magnolia), <i>Mallotus japonicus</i> (green tiger lotus), <i>Morus alba</i> (white mulberry), <i>Musa basjoo</i> (Japanese banana), <i>Nandina domestica</i> (heavenly bamboo), <i>Nerium indicum</i> (Indian oleander), <i>Olea chrysophylla</i> (African olive), <i>Platanus orientalis</i> (oriental planetree), <i>Portulaca oleracea</i> (pigweeds), <i>Pyrus ussuriensis</i> (ornamental pear), <i>Rhododendron indicum</i> (azalea), <i>Trachycarpus exelsus fortunei</i> (wind-mill palm), <i>Wisteria floribunda</i> (Japanese wisteria) (Ben-Dov 1994).
<b>Distribution</b>	China, Japan, Philippines, South Korea, USA (Ben-Dov et al. 1994).
<b>Quarantine pest</b>	<i>Pseudococcus maritimus</i> (Ehrhorn 1900)
<b>Synonyms</b>	<i>Dactylopius maritimus</i> ; <i>Pseudococcus bakeri</i> ; <i>Pseudococcus omniverae</i>
<b>Common name(s)</b>	American grape mealybug, Baker's mealybug; grape mealybug; ocean mealybug
<b>Main hosts</b>	<i>Acacia julibrissin</i> ; <i>Acer</i> spp.; <i>Alternanthera</i> spp.; <i>Annona hastata</i> ; <i>Arbutus</i> spp.; <i>Astragalus</i> spp.; <i>Berberis compacta gracilis</i> ; <i>Boerhavia nivea</i> ; <i>Carya</i> spp.; <i>Catalpa</i> spp.; <i>Ceanothus</i> spp.; <i>Celtis</i> spp.; <i>Cestrum</i> spp.; <i>Chysis aurea</i> ; <i>Citrus</i> spp.; <i>Cornus florida</i> ; <i>Corylus americana</i> ; <i>Cotoneaster</i> spp.; <i>Cupressus</i> spp.; <i>Cydonia</i> spp.; <i>Cyperus</i> spp.; <i>Diospyros</i> spp.; <i>Erigeron</i> spp.; <i>Eriogonum</i> spp.; <i>Eustoma russellianum</i> ; <i>Fraxinus caroliniana</i> ; <i>Genista</i> spp.; <i>Gleditsia triacanthos</i> ; <i>Grevillea</i> spp.; <i>Haplopappus ericoides</i> ; <i>Illex vomitoria</i> ; <i>Ipomoea</i> spp.; <i>Juglans regia</i> ; <i>Juniperus maritima</i> ; <i>Liquidambar styraciflora</i> ; <i>Maclura</i> spp.; <i>Magnolia</i> spp.;

	<i>Malus</i> spp.; <i>Manihot esculenta</i> ; <i>Medicago sativa</i> ; <i>Mesembryanthemum</i> spp.; <i>Morus</i> spp.; <i>Narcissus</i> spp.; <i>Odontoglossum grande</i> ; <i>Ostrya virginiana</i> ; <i>Parthenium</i> spp.; <i>Persea</i> spp.; <i>Platanus</i> spp.; <i>Polygonum</i> spp.; <i>Prunus</i> spp.; <i>Psoralea macrostachya</i> ; <i>Pyrus communis</i> ; <i>Ramona stachyoides</i> ; <i>Rhododendron</i> spp.; <i>Rhus diversiloba</i> spp.; <i>Robinia</i> spp.; <i>Rubus vitifolius</i> ; <i>Sambucus glauca</i> ; <i>Sambucus</i> spp.; <i>Sassafras</i> spp.; <i>Solanum melongena</i> ; <i>Solidago sempervirens</i> ; <i>Strelitzia</i> spp.; <i>Tapirira edulis</i> ; <i>Taxus</i> spp.; <i>Thuja</i> spp.; <i>Tilia americana</i> ; <i>Trifolium</i> spp.; <i>Ulmus</i> spp.; <i>Vaccinium</i> spp.; <i>Vitis</i> spp.; <i>Zantedeschia</i> spp. (Ben-Dov <i>et al.</i> 2012).
<b>Distribution</b>	Argentina; Armenia; Bermuda; Canada; Chile; Colombia; French Guiana; Guadeloupe; Guatemala; Indonesia; Mexico; Poland; Puerto Rico & Vieques Island; United States of America (Ben-Dov <i>et al.</i> 2012).
<b>Quarantine pest</b>	<i>Sinoxylon perforans</i> Schrank
<b>Synonyms</b>	<i>Bostrichus perforans</i> Shrank, <i>Sinoxylon muricatum</i> Duftschmid
<b>Common name(s)</b>	Branch borer, Twig borer, Vine borer
<b>Main hosts</b>	Hosts include <i>Quercus</i> spp. and <i>Vitis</i> spp., (Filip 1986; Taralashvii 1989; Ragazzini 1996). Other deciduous trees and orchards crops are also likely to be attacked (Solomon 1995)
<b>Distribution</b>	Central Asia, Europe including Russia (Filip 1986; Ragazzini 1996; Taralashvii 1989)
<b>Quarantine pest</b>	<i>Sinoxylon sexdentatum</i> Olivier
<b>Synonyms</b>	-
<b>Common name(s)</b>	-
<b>Main hosts</b>	<i>Vitis</i> spp. (Moleas 1988)
<b>Distribution</b>	Apulia (Italy) (Moleas 1988)
<b>Quarantine pest</b>	<i>Targionia vitis</i> Signoret 1876
<b>Synonyms</b>	<i>Aspidiotus vitis</i> ; <i>Diaspis blanckenhorni</i> ; <i>Diaspis blankenhornii</i> ; <i>Targionia arbutus</i> ; <i>Targionia suberi</i> ; <i>Targionia vitis</i> ; <i>Targionia vitis arbutus</i> ; <i>Targionia vitis suberi</i> , <i>Targionia arbutus</i> , <i>Targionia suberi</i> (Ben-Dov <i>et al.</i> 2012).
<b>Common name(s)</b>	Grapevine black scale
<b>Main hosts</b>	<i>Arbutus unedo</i> ; <i>Castanea crenata</i> ; <i>Castanea sativa</i> ; <i>Fagus sylvatica</i> ; <i>Platanus orientalis</i> ; <i>Quercus cerris</i> ; <i>Q. coccifera</i> ; <i>Q. dentate</i> ; <i>Q. ilex</i> ; <i>Q. lanuginose</i> ; <i>Q. pubescens</i> ; <i>Q. sessiliflora</i> ; <i>Q. suber</i> <i>Salix</i> spp.; <i>Vitis vinifera</i> (CABI 2012a; Ben-Dov <i>et al.</i> 2012)
<b>Distribution</b>	Algeria, Armenia, Azerbaijan, Czech Republic, Corsica, Georgia, Greece, Hungary, Iran, Iraq, Israel, Italy, France, Malta, Morocco, Portugal, Romania, Russia, Sardinia, Spain, Turkey, Ukraine, Yugoslavia (Ben-Dov <i>et al.</i> 2012).
<b>Quarantine pest</b>	<i>Zeuzera coffeae</i> Nietner
<b>Synonyms</b>	<i>Zeuzera roricyanea</i>
<b>Common name(s)</b>	Carpenter worm, cocoa pod and stem borer, coffee leopard moth, red branch borer, red coffee borer, red twig borer, tea stem borer
<b>Main hosts</b>	<i>Z. coffeae</i> is highly polyphagous and has been recorded on over 40 hosts including: <i>Abelmoschus esculentus</i> , <i>Acacia auriculiformis</i> , <i>Acacia mangium</i> , <i>Artocarpus</i> , <i>Camellia sinensis</i> , <i>Carya</i> , <i>Castanea</i> , <i>Ceiba pentandra</i> , <i>Cinnamomum verum</i> , <i>Citrus</i> , <i>Clausena lansium</i> , <i>Coffea</i> , <i>Eucalyptus</i> spp., <i>Gossypium</i> , <i>Juglans regia</i> , <i>Leucaena leucocephala</i> , <i>Malus domestica</i> , <i>Manihot esculenta</i> , <i>Persea americana</i> , <i>Populus</i> , <i>Robinia pseudoacacia</i> , <i>Swietenia</i> , <i>Tectona grandis</i> , <i>Theobroma cacao</i> and <i>Vitis vinifera</i> (Mathew 1987; Chang 1988; Schoorl 1990; Griffiths <i>et al.</i> 2004).
<b>Distribution</b>	Bangladesh, China, Cambodia, India, Malaysia, Sri Lanka, Taiwan, Thailand, Vietnam, Papua New Guinea (Chang 1988; Waterhouse 1993; Griffiths <i>et al.</i> 2004; EPPO 2009)
<b>BACTERIA</b>	
<b>Quarantine pest</b>	<i>Xanthomonas campestris</i> pv. <i>viticola</i> (Nayudu) Dye

<b>Synonyms</b>	<i>Pseudomonas viticola</i> Nayudu sp. nov.
<b>Common name(s)</b>	Bacterial canker of grapevine
<b>Main hosts</b>	<i>Alternanthera tenella</i> , <i>Amaranthus</i> spp., <i>Glycine</i> spp. <i>Senna obtusifolia</i> and <i>Vitis vinifera</i> (Peixoto <i>et al.</i> 2007)
<b>Distribution</b>	Brazil and India (Trindade <i>et al.</i> 2005).
<b>Quarantine pest</b>	<i>Xylella fastidiosa</i> (Wells <i>et al.</i> ) – grapevine strain
<b>Synonyms</b>	
<b>Common name(s)</b>	Pierce's disease
<b>Main hosts</b>	Wide host range
<b>Distribution</b>	Central America, North America, Peru (Janse and Obradovic 2010). Unconfirmed report in Kosovo (Janse and Obradovic 2010).
<b>Quarantine pest</b>	<i>Xylophilus ampelinus</i> (Panagopoulos 1969) Willems <i>et al.</i> 1987
<b>Synonyms</b>	<i>Xanthomonas ampelina</i> Panagopoulos 1969
<b>Common name(s)</b>	Canker of grapevine
<b>Main hosts</b>	<i>Vitis vinifera</i> (Panagopoulos 1988).
<b>Distribution</b>	France (Manceau <i>et al.</i> 2005); Greece, Italy (CABI/EPPO 1999); Slovenia (Dreo <i>et al.</i> 2005); South Africa (Botha <i>et al.</i> 2001)
<b>FUNGI</b>	
<b>Quarantine pest</b>	<i>Alternaria viticola</i> Brunaud
<b>Synonyms</b>	-
<b>Common name(s)</b>	Spike-stalk brown spot of grape, brown blotch, grape rachis blotch
<b>Main hosts</b>	<i>Vitis</i> species including some hybrid grapes (Liu <i>et al.</i> 1996; Ma <i>et al.</i> 2004).
<b>Distribution</b>	China (Liu <i>et al.</i> 1996; Ma <i>et al.</i> 2004)
<b>Quarantine pest</b>	<i>Cadophora luteo-olivacea</i> (J.F.H Beyma) T.C. Harr. & McNew
<b>Synonyms</b>	<i>Phialophora luteo-olivacea</i> J.F.H. Beyma
<b>Common name(s)</b>	-
<b>Main hosts</b>	Grapevines (Gramaje <i>et al.</i> 2011), kiwifruit (Prodi <i>et al.</i> 2008)
<b>Distribution</b>	California, Italy, New Zealand, Northeastern America, South Africa, Spain and Uruguay (Prodi <i>et al.</i> 2008; Gramaje and Armengol 2011).
<b>Quarantine pest</b>	<i>Cadophora melinii</i> Nannf.
<b>Synonyms</b>	<i>Phialophora melinii</i> (Nannf.) Conant
<b>Common name(s)</b>	-
<b>Main hosts</b>	Grapevines (Gramaje <i>et al.</i> 2011), kiwifruit (Prodi <i>et al.</i> 2008)
<b>Distribution</b>	Italy (Prodi <i>et al.</i> 2008), Spain (Gramaje <i>et al.</i> 2011), Uruguay (Navarrete <i>et al.</i> 2010)
<b>Quarantine pest</b>	<i>Eutypella leprosa</i> (Pers.) Berl.
<b>Synonyms</b>	<i>Sphaeria leprosa</i> Pers
<b>Common name(s)</b>	-
<b>Main hosts</b>	<i>Aesculus</i> spp., <i>Corylus</i> spp. <i>Fraxinus</i> spp., <i>Tilia</i> spp., <i>Vitis vinifera</i> L. (Vasilyeva and Stephenson 2006; Diaz <i>et al.</i> 2011; Farr and Rossman 2011).
<b>Distribution</b>	Chile (Diaz <i>et al.</i> 2011), USA (Vasilyeva and Stephenson 2006).
<b>Quarantine pest</b>	<i>Eutypella vitis</i> (Schwein.:Fr.) Ellis & Fischer
<b>Synonyms</b>	<i>Diatrype vitis</i> (Schwein.: Fr.) Berk, <i>Engizostoma vitis</i> (Schwein.: Fr.) Kuntze, <i>Eutypella aequilinaris</i> , <i>Sphaeria vitis</i> Schwein., Schrift, <i>Valsa vitis</i> (Schwein.: Fr.) M.A. Curtis,

	(Vasilyeva and Stephenson 2006; Catal <i>et al.</i> 2007)
<b>Common name(s)</b>	-
<b>Main hosts</b>	<i>Vitis</i> spp. (Catal <i>et al.</i> 2007; Vasilyeva and Stephenson 2006)
<b>Distribution</b>	Eastern United States in North America (Farr <i>et al.</i> 1989; Vasilyeva and Stephenson 2006)
<b>Quarantine pest</b>	<i>Fomitiporia mediterranea</i> M. Fisher
<b>Synonyms</b>	-
<b>Common name(s)</b>	Esca disease
<b>Main hosts</b>	<i>Acer negundo</i> , <i>Actinidia chinensis</i> , <i>Cornus mas</i> , <i>Corylus avellana</i> , <i>Lagerstroemia indica</i> , <i>Laurus nobilis</i> , <i>Ligustrum vulgare</i> , <i>Olea europaea</i> , <i>Quercus</i> spp., <i>Quercus ilex</i> , <i>Robinia pseudoacacia</i> , <i>Vitis vinifera</i> (Fischer 2002; Fisher and Binder 2004; Fischer 2006; Amalfi <i>et al.</i> 2010; Pilotti <i>et al.</i> 2010).
<b>Distribution</b>	Algeria, Austria, France, Germany, Greece, Iran, Italy, Portugal, Slovenia, Spain, Switzerland (Karimi <i>et al.</i> 2001; Fischer 2002; Fischer 2006; Péros <i>et al.</i> 2008; Pilotti <i>et al.</i> 2010)
<b>Quarantine pest</b>	<i>Fomitiporia polymorpha</i> M. Fisher (recently described species, limited information)
<b>Synonyms</b>	-
<b>Common name(s)</b>	-
<b>Main hosts</b>	Hardwoods (Fisher and Binder 2004)
<b>Distribution</b>	North America, USA (Fisher and Binder 2004; Fischer 2006; Pilotti <i>et al.</i> 2010)
<b>Quarantine pest</b>	<i>Guignardia</i> spp. ( <i>G. bidwellii</i> , <i>G. bidwellii</i> f. <i>euvitis</i> , <i>G. bidwellii</i> f. <i>muscadinii</i> )
<b>Synonyms</b>	<i>Botryosphaeria bidwellii</i> , <i>Carlia bidwellii</i> , <i>Depazea labruscae</i> , <i>Greenaria uvicola</i> , <i>Laestadia bidwellii</i> , <i>Naemospora ampellicida</i> , <i>Phoma ustulata</i> , <i>Phoma uvicola</i> var. <i>labruscae</i> , <i>Phoma uvicola</i> , <i>Phyllosticta ampellicida</i> , <i>Phyllosticta ampelopsidis</i> , <i>Phyllosticta viticola</i> , <i>Phyllosticta vulpinae</i> , <i>Phyllostictina clemensae</i> , <i>Phyllostictina uvicola</i> , <i>Phyllostictina viticola</i> , <i>Physalospora bidwellii</i> , <i>Sacidium viticolum</i> , <i>Septoria viticola</i> , <i>Sphaeria bidwellii</i> (Ullrich <i>et al.</i> 2009; CABI 2012a).
<b>Common name(s)</b>	Black rot
<b>Main hosts</b>	<i>Ampelopsis</i> , <i>Asplenium nidus</i> , <i>Cissus</i> , <i>Citrus</i> , <i>Parthenocissus quinquefolia</i> , <i>P. tricuspidata</i> , <i>V. amurensis</i> , <i>Vitis arizonica</i> , <i>Vitis labrusca</i> , <i>Vitis rotundifolia</i> , <i>Vitis vinifera</i> (University of Illinois 2001; Eyres <i>et al.</i> 2006; Ullrich <i>et al.</i> 2009; CABI 2012a).
<b>Distribution</b>	Argentina, Austria, Barbados, Brazil, Bulgaria, Canada, Chile, China, Cuba, Cyprus, El Salvador, Former Yugoslavia, France, Germany, Guyana, Haiti, India, Iran, Italy, Jamaica, Japan, Korea, Martinique, Mexico, Morocco, Mozambique, Pakistan, Panama, Philippines, Romania, Russian Federation, Slovakia, Sudan, Switzerland, Turkey, Ukraine, Virgin Islands, Uruguay, USA and Venezuela (AQSIQ 2006; Eyres <i>et al.</i> 2006; AQSIQ 2007; Ullrich <i>et al.</i> 2009; CABI 2012a).
<b>Quarantine pest</b>	<i>Inocutis jamaicensis</i> (Murrill) J.E. Wright & Moncalvo
<b>Synonyms</b>	-
<b>Common name(s)</b>	Grapevine trunk disease – ‘Hoja de malvon’
<b>Main hosts</b>	<i>Vitis vinifera</i> , <i>Eucalyptus globulus</i> , <i>Diostea</i> spp., <i>Prunus</i> spp, <i>Quercus</i> spp. <i>Taxodium</i> spp. (Lupo <i>et al.</i> 2006; Fischer 2006; Perez <i>et al.</i> 2008)
<b>Distribution</b>	North America, South America (Fischer 2006; Lupo <i>et al.</i> 2006; Perez <i>et al.</i> 2008)
<b>Quarantine pest</b>	<i>Monilinia fructigena</i> (Aderh. & Ruhland) Honey
<b>Synonyms</b>	<i>Monilia fructigena</i> Schumacher, <i>Sclerotinia fructigena</i> (J. Schröt.) Norton, <i>Sclerotinia fructigena</i> Aderh., <i>Stromatinia fructigena</i> (J. Schröt.) Boud (Ma 2006; CABI 2012a).
<b>Common name(s)</b>	Brown rot
<b>Main hosts</b>	<i>Amelanchier canadensis</i> , <i>Berberis</i> , <i>Capsicum</i> , <i>Cornus mas</i> , <i>Corylus avellana</i> ,



	<i>Cotoneaster</i> , <i>Crataegus laevigata</i> , <i>Cydonia oblonga</i> , <i>Diospyros kaki</i> , <i>Eriobotrya japonica</i> , <i>Ficus carica</i> , <i>Fragaria</i> spp., <i>Solanum lycopersicum</i> , <i>Malus domestica</i> , <i>Mespilus germanica</i> , <i>Prunus</i> spp., <i>Psidium guajava</i> , <i>Pyrus</i> spp., <i>Rhododendron</i> , <i>Rosa</i> , <i>Rubus</i> spp., <i>Sorbus</i> , <i>Vaccinium</i> , <i>Vitis vinifera</i> (Sharma and Kaul 1989; Mackie 2005; Ma 2006; CABI 2012a).
<b>Distribution</b>	China, Taiwan, Afghanistan, Armenia, Austria, Azerbaijan, Belarus, Belgium, Brazil, Bulgaria, Chile, Croatia, Cyprus, Czech Republic, Denmark, Egypt, Finland, France, Georgia, Germany, Greece, Hungary, India, Iran, Ireland, Israel, Italy, Japan, Latvia, Lebanon, Lithuania, Luxembourg, Moldova, Montenegro, Morocco, Nepal, North Korea, Norway, Netherlands, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, South Korea, Spain, Sweden, Switzerland, Turkey, Ukraine, United Kingdom, Uruguay, Uzbekistan, Yugoslavia (Mackie 2005; Ma 2006; AQSIQ 2007; CABI 2012a).
<b>Quarantine pest</b>	<i>Phaeoacremonium</i> spp. ( <i>P. alvesii</i> , <i>P. angustus</i> , <i>P. argentinense</i> , <i>P. armeniacum</i> , <i>P. austroafricanum</i> , <i>P. cinereum</i> , <i>P. croatiense</i> , <i>P. globosum</i> , <i>P. griseorubrum</i> , <i>P. hispanicum</i> , <i>P. hungaricum</i> , <i>P. inflatipes</i> , <i>P. iraniamum</i> , <i>P. krajdenii</i> , <i>P. mortoniae</i> , <i>P. occidentale</i> , <i>P. rubrigenum</i> , <i>P. scolyti</i> , <i>P. sicilianum</i> , <i>P. subulatum</i> , <i>P. tuscanum</i> , <i>P. venezuelense</i> , <i>P. viticola</i> )
<b>Synonyms</b>	-
<b>Common name(s)</b>	Petri and esca diseases
<b>Main hosts</b>	<i>Dodoneae viscosa</i> , <i>Fraxinus excelsior</i> , <i>Fraxinus latifolia</i> , <i>Fraxinus pennsylvanica</i> , <i>Nectandra</i> spp., <i>Quercus virginiana</i> , <i>Sorbus intermedia</i> , <i>Vitis vinifera</i> (Mostert <i>et al.</i> 2005; Mostert <i>et al.</i> 2006a; Essakhi <i>et al.</i> 2008).
<b>Distribution</b>	Canada, Czech Republic, Chile, Costa Rica, Croatia, Democratic Republic of Congo, France, Germany, India, Iran, Italy, Japan, Norway, Portugal, South Africa, Spain, Sweden, Turkey, USA, Venezuela, Zaire (Mostert <i>et al.</i> 2005; Mostert <i>et al.</i> 2006a; Essakhi <i>et al.</i> 2008; Gramaje <i>et al.</i> 2009a)
<b>Quarantine pest</b>	<i>Phakopsora</i> spp. ( <i>Phakopsora euvitis</i> , <i>Phakopsora muscadinae</i> , <i>Phakopsora uva</i> )
<b>Synonyms</b>	Synonyms of <i>P. euvitis</i> : <i>Aecidium meliosmae-myrianthae</i> , <i>Phakopsora ampelopsidis</i> , <i>Physopella ampelopsidis</i> , <i>Physopella vialae</i> , <i>Physopella vitis</i> , <i>Uredo vialae</i> , <i>Uredo vitis</i> (Hennen <i>et al.</i> 2005; CABI 2012a). Note: <i>P. muscadinae</i> has been determined to be conspecific with <i>P. uva</i> reported from Mexico (Hennessy <i>et al.</i> 2007). <i>P. uva</i> was reported to occur on unidentified species of <i>Vitis</i> in Colombia and in Mexico (Chalkley 2011).
<b>Common name(s)</b>	Grapevine rust, grapevine leaf rust
<b>Main hosts</b>	<i>Vitis</i> spp. ( <i>V. amurensis</i> , <i>V. coignetiae</i> , <i>V. ficifolia</i> , <i>V. flexuosa</i> , <i>V. labrusca</i> , <i>V. vinifera</i> ), <i>Meliosma</i> spp., <i>Meliosma dilleniifolia</i> subsp. <i>cuneifolia</i> , <i>Meliosma myriantha</i> (Ono 2000; Weinert <i>et al.</i> 2003; Chalkley 2011; CABI 2012a).
<b>Distribution</b>	Bangladesh, Barbados, Brazil, China, Colombia, Costa Rica, Cuba, Democratic People's Republic of Korea, Guatemala, Honduras, India, Indonesia, Jamaica, Japan, Korea, Malaysia, Mexico, Myanmar, Nepal, Philippines, Puerto Rico, Russian Far East, Sri Lanka, Thailand, Trinidad and Tobago, USA, Venezuela, Vietnam, Virgin Islands (Ono 2000; Tessman <i>et al.</i> 2004; Chalkley 2011; CABI 2012a).
<b>PHYTOPLASMAS</b>	
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma asteris [16Srl –Aster yellows group]
<b>Synonyms</b>	
<b>Strains</b>	16Srl-A; 16Srl-B, 16Srl-C
<b>Common name(s)</b>	grapevine yellows, North American grapevine yellows, Virginia grapevine yellows I
<b>Main hosts</b>	Wide host including Grapevines (Firrao <i>et al.</i> 2005)
<b>Distribution</b>	On grapevines reported from Canada (Olivier <i>et al.</i> 2009b), Chile (Gajardo <i>et al.</i> 2009), Germany (Prince <i>et al.</i> 1993), Israel (Tanne and Orenstein 1997), Italy (Alma <i>et al.</i> 1996), South Africa (Engelbrecht <i>et al.</i> 2010), Tunisia (Mhirsi <i>et al.</i> 2004), USA (Davis <i>et al.</i> 1998)

	and Turkey (Canik <i>et al.</i> 2011).
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma fraxini [16SrVII]
<b>Synonyms</b>	
<b>Strains</b>	
<b>Common name(s)</b>	Chile grapevine yellows
<b>Main hosts</b>	<i>Fraxinus</i> spp. and grapevines (Gajardo <i>et al.</i> 2009)
<b>Distribution</b>	In grapes reported from Chile (Gajardo <i>et al.</i> 2009)
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma phoenicium [16SrIX]
<b>Synonyms</b>	Phytoplasma 16SrIX
<b>Strains</b>	
<b>Common name(s)</b>	Grapevine yellows
<b>Main hosts</b>	<i>Vitis vinifera</i> (grapes)
<b>Distribution</b>	Turkey (Canik <i>et al.</i> 2011)
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma pruni [16SrIII – peach X-disease phytoplasmas group]
<b>Synonyms</b>	Western x Virginia grapevine yellows III
<b>Strains</b>	
<b>Common name(s)</b>	Grapevine yellows x disease
<b>Main hosts</b>	<i>Delphinium</i> spp. (Harju <i>et al.</i> 2008), grapevine (Davis and Dally 2001), <i>Prunus</i> spp. (Zhao <i>et al.</i> 2009).
<b>Distribution</b>	In grapevine reported from Israel (Tanne and Orenstein 1997), Italy (Bianco <i>et al.</i> 1996) and the USA (Davis and Dally 2001)
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma solani [16SrXII–A Stolburg group]
<b>Synonyms</b>	Bois noir Phytoplasma
<b>Strains</b>	STOL Type I; STOL Type II; STOL Type III (Langer and Maxiner 2004).
<b>Common name(s)</b>	Bois noir, Legno nero, Vergilbungskrankheit, Schwartzholzkrankheit
<b>Main hosts</b>	<i>Calystegia sepium</i> (Mori <i>et al.</i> 2007); <i>Convolvulus arvensis</i> ; <i>Lavandula</i> spp.; <i>Ranunculus</i> spp.; <i>Solanum</i> spp.; <i>Urtica dioica</i> ; <i>Vitis</i> species (Constable 2010). Type 1: <i>Urtica dioica</i> ; Type II: <i>Calystegia sepium</i> , <i>Convolvulus arvensis</i> ; Type III: <i>Calystegia sepium</i> (Mori <i>et al.</i> 2007).
<b>Distribution</b>	Bois noir Phytoplasma is widespread and occurs from Spain to Ukraine and from Germany and Northern France to Lebanon and Israel (Maixner 2011). It has also been reported from Canada (Rott <i>et al.</i> 2007), Syria (Contaldo <i>et al.</i> 2011), Turkey (Canik <i>et al.</i> 2011) and the USA (Davis <i>et al.</i> 1998). Additionally Stolbur group-related grapevine phytoplasmas have been also recently been reported from Iran (Karimi <i>et al.</i> 2009), Chile (Gajardo <i>et al.</i> 2009) and China (Duduk <i>et al.</i> 2010).
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma ulmi [16SrV–A Elm yellows phytoplasma group]
<b>Synonyms</b>	
<b>Strains</b>	
<b>Common name(s)</b>	Grapevine yellows disease
<b>Main hosts</b>	Wide host range including grapes
<b>Distribution</b>	In grapevine reported from Italy (Botti and Bertaccini 2007)
<b>Quarantine pest</b>	<i>Candidatus</i> Phytoplasma vitis [16SrV–Elm yellows phytoplasma group]
<b>Synonyms</b>	Grapevine Flavescence dorée Phytoplasma
<b>Strains</b>	FD-I, FD-II, FD-III, Phytoplasma strains FD-associated belong to ribosomal subgroups



	16SrV-C, 16SrV-D.
<b>Common name(s)</b>	Flavescence dorée
<b>Main hosts</b>	<i>Vitis vinifera</i> (grapes), but <i>V. riparia</i> can also be infected naturally (Maixner and Pearson 1992).
<b>Distribution</b>	Croatia (Filippin <i>et al.</i> 2009), France (Steffek <i>et al.</i> 2006), Germany (Johannesen <i>et al.</i> 2008), Italy (Barba <i>et al.</i> 2006), Macedonia (Filippin <i>et al.</i> 2009), Portugal (DeSousa <i>et al.</i> 2003), Serbia (Duduk <i>et al.</i> 2003), Slovenia (Filippin <i>et al.</i> 2009), Spain (Battlle <i>et al.</i> 2000), Switzerland (Steffek <i>et al.</i> 2006).
<b>Quarantine pest</b>	European stone fruit yellows Phytoplasma [16SrX –B Apple proliferation group]
<b>Synonyms</b>	Grapevine yellows
<b>Strains</b>	
<b>Common name(s)</b>	Grapevine yellows
<b>Main hosts</b>	<i>Vitis vinifera</i> (grapes)
<b>Distribution</b>	In grape vine reported from Hungary (Varga <i>et al.</i> 2000) and Serbia (Duduk <i>et al.</i> 2003).
<b>VIRUSES</b>	
<b>Quarantine pest</b>	<i>Arabis mosaic virus</i> – grape strains
<b>Synonyms</b>	None
<b>Common name(s)</b>	Arabis mosaic
<b>Main hosts</b>	The strain of ArMV infecting grapevine affects a range of host plants and produces characteristic symptoms (Fortusini <i>et al.</i> 1983; Belli <i>et al.</i> 1982)
<b>Distribution</b>	Balkans, Bulgaria, Canada, Croatia, Central Europe, France, Germany, Hungary, Israel, Italy, Japan, New Zealand, Romania, Switzerland, Ukraine Yugoslavia (Cadman <i>et al.</i> 1960; Kearns and Mossop 1984; MacKenzie <i>et al.</i> 1996; Delibašić <i>et al.</i> 2000), Iran (Pourrahim <i>et al.</i> 2004) and Spain (Abelleira <i>et al.</i> 2010)
<b>Quarantine pest</b>	<i>Artichoke Italian latent virus</i> (AILV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Artichoke patchy chlorotic stunting disease, Yellowing disease of artichoke
<b>Main hosts</b>	<i>Cynara scolymus</i> , <i>Cichorium intybus</i> , <i>Crepis neglecta</i> , <i>Gladiolus</i> spp., <i>Helminthia echinoides</i> , <i>Hypochoeris aetensis</i> , <i>Lactuca virosa</i> , <i>Urospermum dalechampii</i> , <i>Lamium amplexicaule</i> , <i>Pelargonium zonale</i> , <i>Sonchus</i> spp., <i>Vitis vinifera</i> (Brunt <i>et al.</i> 1996)
<b>Distribution</b>	Bulgaria (Savino <i>et al.</i> 1977), Greece (Kyriakopoulou 2008), Italy (Roca <i>et al.</i> 1975) and Russia (Gallitelli <i>et al.</i> 2004).
<b>Quarantine pest</b>	<i>Blueberry leaf mottle virus</i> (BLMoV) New York (NY) strain
<b>Synonyms</b>	
<b>Common name(s)</b>	Fanleaf degeneration/decline disease
<b>Main hosts</b>	Grapevines (Uyemoto <i>et al.</i> 1977)
<b>Distribution</b>	USA (Uyemoto <i>et al.</i> 1977)
<b>Quarantine pest</b>	<i>Cherry leafroll virus</i> – grapevine isolate (CLRV)
<b>Synonyms</b>	
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis vinifera</i>
<b>Distribution</b>	Chile (Herrera and Madariaga 2001) and Germany (Ipach <i>et al.</i> 2003).
<b>Quarantine pest</b>	<i>Grapevine ajinashika virus</i> (GAGV)
<b>Synonyms</b>	

<b>Common name(s)</b>	Grapevine ajinashika disease
<b>Main hosts</b>	<i>Vitis vinifera</i> cv. Koshu (Namba <i>et al.</i> 1991b)
<b>Distribution</b>	Japan (Namba <i>et al.</i> 1991b).
<b>Quarantine pest</b>	<i>Grapevine angular mosaic-associated virus</i> (GAMaV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine angular mosaic
<b>Main hosts</b>	<i>Vitis vinifera</i> (Girgis <i>et al.</i> 2009).
<b>Distribution</b>	Greece (Girgis <i>et al.</i> 2009).
<b>Quarantine pest</b>	<i>Grapevine Anatolian ringspot virus</i> (GARSV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Fanleaf degeneration/decline disease
<b>Main hosts</b>	<i>Vitis vinifera</i> cultivar Kizlar Tahasi (Gokalp <i>et al.</i> 2003).
<b>Distribution</b>	Turkey (Cigsar <i>et al.</i> 2002; Gokalp <i>et al.</i> 2003; Laimer <i>et al.</i> 2009).
<b>Quarantine pest</b>	<i>Grapevine asteroid mosaic associated virus</i> (GAMV)
<b>Synonyms</b>	
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis vinifera</i> (Martelli and Boudon-Padieu 2006).
<b>Distribution</b>	California, USA (Martelli and Boudon-Padieu 2006). Records from Italy and South Africa have not been confirmed experimentally and a record from Greece was proven to refer to <i>Grapevine rupestris vein feathering virus</i> (Martelli and Boudon-Padieu 2006).
<b>Quarantine pest</b>	<i>Grapevine berry inner necrosis virus</i> (GINV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine berry inner necrosis disease
<b>Main hosts</b>	<i>Vitis vinifera</i> (Yoshikawa <i>et al.</i> 1997).
<b>Distribution</b>	Japan (Yoshikawa <i>et al.</i> 1997).
<b>Quarantine pest</b>	<i>Grapevine Bulgarian latent virus</i> (GBLV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Fanleaf degeneration/decline disease
<b>Main hosts</b>	<i>Vitis vinifera</i> (Gokalp <i>et al.</i> 2003).
<b>Distribution</b>	Bulgaria (Martelli <i>et al.</i> 1977, 1978), Portugal (Sequeira and Mendonça, 1992) Yugoslavia, Czechoslovakia, former USSR, Hungary (Martelli 1993).
<b>Quarantine pest</b>	<i>Grapevine chrome mosaic virus</i>
<b>Synonyms</b>	<i>Bratislava mosaic virus</i> , <i>Hungarian yellow mosaic</i> , <i>Hungarian chrome mosaic virus</i>
<b>Common name(s)</b>	Fanleaf degeneration/decline disease
<b>Main hosts</b>	<i>Vitis vinifera</i> (Dimou <i>et al.</i> 1994).
<b>Distribution</b>	Austria, Croatia, the former Czechoslovakia, and Hungary (Uyemoto <i>et al.</i> 2009).
<b>Quarantine pest</b>	<i>Grapevine deformation virus</i>
<b>Synonyms</b>	
<b>Common name(s)</b>	Fanleaf degeneration/decline disease
<b>Main hosts</b>	<i>Vitis vinifera</i> (Cigsar <i>et al.</i> 2003).
<b>Distribution</b>	Turkey (Cigsar <i>et al.</i> 2003; Digiario <i>et al.</i> 2003).

<b>Quarantine pest</b>	<i>Grapevine fanleaf virus</i> (GFLV)
<b>Synonyms</b>	<i>Grapevine arrciamento virus</i> , <i>Grapevine court-noué virus</i> , <i>Grapevine infectious degeneration virus</i> , <i>Grapevine Reisigkrankheit Virus</i> , <i>Grapevine roncet virus</i> <i>Grapevine urticado virus</i>
<b>Common name(s)</b>	Fanleaf disease
<b>Main hosts</b>	<i>Vitis</i> species (Andret-Link <i>et al.</i> 2004)
<b>Distribution</b>	Asia, Africa, Europe, New Zealand, North America and South America (Andret-Link <i>et al.</i> 2004).
<b>Quarantine pest</b>	<i>Grapevine leafroll associated virus 6</i> (GLRaV-6)
<b>Synonyms</b>	
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis</i> species
<b>Distribution</b>	Brazil (Kuniyuki <i>et al.</i> 2008), Italy (Boscia <i>et al.</i> 2000), North Africa (Eddin <i>et al.</i> 2008, Mahfoudhi <i>et al.</i> 2009), Switzerland (Gugerli and Ramel 1993) Turkey (Cigsar <i>et al.</i> 2002), USA (Martinson <i>et al.</i> 2008, Fuchs 2007).
<b>Quarantine pest</b>	<i>Grapevine leafroll associated virus 7</i> (GLRaV-7)
<b>Synonyms</b>	
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis</i> species
<b>Distribution</b>	Albania, Armenia, Greece, Italy, Jordan (Digiario <i>et al.</i> 2000 ) and Portugal (Santos <i>et al.</i> 2000)
<b>Quarantine pest</b>	<i>Grapevine leafroll associated virus 10</i> (GLRaV-10)
<b>Synonyms</b>	<i>Grapevine leafroll associated virus De</i> (GLRaV-De)
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis</i> species
<b>Distribution</b>	Greece (Maliogka <i>et al.</i> 2008b)
<b>Quarantine pest</b>	<i>Grapevine leafroll associated virus 11</i> (GLRaV-11)
<b>Synonyms</b>	<i>Grapevine leafroll associated virus Pr</i> (GLRaV-Pr)
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis</i> species
<b>Distribution</b>	Greece (Maliogka <i>et al.</i> 2008b)
<b>Quarantine pest</b>	<i>Grapevine line pattern virus</i> (GLPV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine line pattern
<b>Main hosts</b>	<i>Vitis vinifera</i> (Martelli and Boudon-Padieu 2006).
<b>Distribution</b>	Hungary (Martelli and Boudon-Padieu 2006).
<b>Quarantine pest</b>	<i>Grapevine Pinot gris virus</i> (GPGV)
<b>Synonyms</b>	
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis vinifera</i> (Giampetruzzi <i>et al.</i> 2012).
<b>Distribution</b>	Italy (Giampetruzzi <i>et al.</i> 2012), Korea (Cho <i>et al.</i> 2013).

<b>Quarantine pest</b>	<i>Grapevine red blotch-associated virus</i> (GRBaV)
<b>Synonyms</b>	<i>Grapevine cabernet franc-associated virus</i> (GCFaV)
<b>Common name(s)</b>	Red blotch disease
<b>Main hosts</b>	<i>Vitis vinifera</i> (Stamp and Wei 2013)
<b>Distribution</b>	United States (Stamp and Wei 2013). A virus genetically identical to GRBaV has also been detected in Canada (Sudarshana 2012).
<b>Quarantine pest</b>	<i>Grapevine red globe virus</i>
<b>Synonyms</b>	
<b>Common name(s)</b>	None
<b>Main hosts</b>	<i>Vitis vinifera</i> (Martelli and Boudon-Padieu 2006).
<b>Distribution</b>	Albania, Italy (Sabanadzovic <i>et al.</i> 2000) and California (Martelli and Boudon-Padieu 2006).
<b>Quarantine pest</b>	<i>Grapevine rupestris vein feathering virus</i> (GRVFV)
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine fleck complex, Syrah Decline
<b>Main hosts</b>	<i>Vitis vinifera</i> (Uyemoto <i>et al.</i> 2009).
<b>Distribution</b>	California, USA (Al Rwahnih <i>et al.</i> 2009), Greece, Italy (Uyemoto <i>et al.</i> 2009)
<b>Quarantine pest</b>	<i>Grapevine syrah virus I</i> (GSyV-I)
<b>Synonyms</b>	
<b>Common name(s)</b>	Syrah decline
<b>Main hosts</b>	<i>Vitis vinifera</i> (Uyemoto <i>et al.</i> 2009).
<b>Distribution</b>	Chile (Engel <i>et al.</i> 2010) and the US (Al Rwahnih <i>et al.</i> 2009)
<b>Quarantine pest</b>	<i>Grapevine Tunisian ringspot virus</i> (GTRSV)
<b>Synonyms</b>	
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis</i> species (Ouertani <i>et al.</i> 1992)
<b>Distribution</b>	Tunisia (Ouertani <i>et al.</i> 1992)
<b>Quarantine pest</b>	<i>Grapevine virus B</i> (GVB) (strains associated with grapevine corky bark)
<b>Synonyms</b>	
<b>Common name(s)</b>	Corky bark disease
<b>Main hosts</b>	<i>Vitis vinifera</i>
<b>Distribution</b>	Brazil, Bulgaria, France, Italy, Japan, Mexico, South Africa, Spain, Switzerland, USA (California), Yugoslavia (Namba <i>et al.</i> 1991a), and Tunisia (Abdallah <i>et al.</i> 2003).
<b>Quarantine pest</b>	<i>Grapevine virus E</i> (GVE)
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine rugose wood complex.
<b>Main hosts</b>	<i>Vitis vinifera</i>
<b>Distribution</b>	Japan (Nakaune <i>et al.</i> 2008) and South Africa (Coetzee <i>et al.</i> 2010)
<b>Quarantine pest</b>	<i>Grapevine virus F</i> (GVF)
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine rugose wood complex
<b>Main hosts</b>	<i>Vitis vinifera</i>

<b>Distribution</b>	USA (California) (Martelli 2012)
<b>Quarantine pest</b>	<i>Peach rosette mosaic virus</i> (PeRMV)
<b>Synonyms</b>	<i>Rosette mosaic virus</i> , <i>Grape decline virus</i> , <i>Grapevine degeneration virus</i>
<b>Common name(s)</b>	
<b>Main hosts</b>	Blueberry, grapevine and peach (Ramsdell and Gillet 1998)
<b>Distribution</b>	Egypt (Fayek <i>et al.</i> 2009), Canada (Ontario) and USA (Michigan) (Ramsdell and Gillet 1998).
<b>Quarantine pest</b>	<i>Petunia asteroid mosaic virus</i> (PeAMV)
<b>Synonyms</b>	<i>Tomato bushy stunt virus – petunia strain</i>
<b>Common name(s)</b>	
<b>Main hosts</b>	Woody hosts (cherries, plums, grapes, privet and dogwood), hops and spinach. PeAMV has also been reported from the roots of <i>Chenopodium album</i> , <i>Cucumis melo</i> , <i>Plantago major</i> and <i>Stellaria media</i> (Lovisol 1990).
<b>Distribution</b>	PeAMV is widely distributed in Asia, Europe and North America; however, on grapes it is reported only from Czechoslovakia, Italy and West Germany (Bercks 1967; Novák and Lanzová 1976; Koenig <i>et al.</i> 1989; Martelli 1993; Constable <i>et al.</i> 2010).
<b>Quarantine pest</b>	<i>Raspberry ringspot virus</i> (RpRSV) – Grapevine strain
<b>Synonyms</b>	
<b>Common name(s)</b>	Grapevine fanleaf disease
<b>Main hosts</b>	<i>Vitis vinifera</i>
<b>Distribution</b>	Germany (Martelli and Boudon-Padieu 2006; Wetzel <i>et al.</i> 2006)
<b>Quarantine pest</b>	<i>Sowbane mosaic virus</i> (SoMV) – grape strains
<b>Synonyms</b>	<i>Chenopodium mosaic virus</i> , <i>Apple latent virus 2</i> , <i>Chenopodium star mottle virus</i>
<b>Common name(s)</b>	
<b>Main hosts</b>	<i>Vitis vinifera</i> (grapevine) (Bercks and Querfurth 1969).
<b>Distribution</b>	Bulgaria, Czechoslovakia and Germany (Bercks and Querfurth 1969; Jankulova 1972; Pozdena <i>et al.</i> 1977)
<b>Quarantine pest</b>	<i>Strawberry latent ringspot virus</i>
<b>Synonyms</b>	<i>Aesculus line pattern virus</i> , <i>Rhubarb virus 5</i>
<b>Common name(s)</b>	
<b>Main hosts</b>	Wide host range 126 species belonging to 27 families (Tzanetakis <i>et al.</i> 2006) including asparagus, blackberries, black currants, celery, cherries, <i>Gladiolus</i> , <i>Narcissus</i> , grapes, plums, peaches, raspberries, red currants, roses, rhubarb, <i>Sambucus nigra</i> and strawberries.
<b>Distribution</b>	SLRSV has been reported from Europe and Israel, New Zealand, North America and Turkey (EPPO 2010a). However, in grapevines, SLRSV infections were reported in Czech Republic (Komínek 2008), France (Walter 1997), Germany (Bercks <i>et al.</i> 1977), Italy (Babini and Bertaccini 1982), Romania (Eppler <i>et al.</i> 1989) and Turkey (Savino <i>et al.</i> 1987; Akbas and Erdiller 1993).
<b>Quarantine pest</b>	<i>Tobacco necrosis virus</i> – grape strain
<b>Synonyms</b>	
<b>Common name(s)</b>	<i>Tobacco necrosis virus</i>
<b>Main hosts</b>	Grapevine (Cesati and Van Regenmortel 1969). NTVs hosts include: <i>Brassica oleracea</i> (cabbage), <i>Chenopodium quinoa</i> (quinoa), <i>Cucumis sativus</i> (cucumber), <i>Cucurbita pepo</i> (zucchini), <i>Daucus carota</i> (carrot), <i>Fragaria x</i>

	<i>ananassa</i> (strawberry), <i>Glycine max</i> (soybean), <i>Malus pumila</i> (apple), <i>Nicotiana tabacum</i> (tobacco), <i>Lactuca sativa</i> (lettuce), <i>Olea europaea</i> (olive), <i>Phaseolus vulgaris</i> (common bean), <i>Solanum tuberosum</i> (potato), <i>Tulipa</i> spp. (tulip) (other hosts are infected but remain symptomless) (Kassanis 1970; Brunt and Teakle 1996; CABI 2012a; Zitikaite and Staniulis 2009).
<b>Distribution</b>	South Africa (Cesati and Van Regenmortel 1969). TNV probably worldwide but species and strain distributions are largely unknown) Belgium, Brazil, Canada, China, Czechoslovakia (former), Denmark, Finland, France, Germany, Hungary, India, Italy, Japan, Latvia, Netherlands, New Zealand, Norway, Romania, Russia, South Africa, Spain, Sweden, Switzerland, Turkey, United Kingdom (CABI 2012a).
<b>Quarantine pest</b>	<i>Tomato black ring virus</i> (TBRV)
<b>Synonyms</b>	<i>Grapevine Joannes-Seyve virus</i> (GJSV), <i>Potato bouquet virus</i> , <i>Potato pseudo-aucuba virus</i> , <i>Tomato blackring virus</i> .
<b>Common name(s)</b>	Ring spot of beet
<b>Main hosts</b>	Wide host range, including carrot, celery, cucumber, <i>Fragaria</i> species, <i>Prunus</i> spp., <i>Ribes</i> spp., <i>Rubus</i> spp., solanaceous species, <i>Vitis vinifera</i> and a number of weed and ornamental species (Harrison 1957, 1958; Pospieszny <i>et al.</i> 2004, Jonczyk <i>et al.</i> 2004).
<b>Distribution</b>	Europe, India, Japan, North and South America (Harper <i>et al.</i> 2010), Israel and Turkey (Uyemoto <i>et al.</i> 2009).
<b>Quarantine pest</b>	<i>Tomato ringspot virus</i> (ToRSV)
<b>Synonyms</b>	ToRSV
<b>Common name(s)</b>	Ringspot virus decline
<b>Main hosts</b>	ToRSV infects a wide range including black currants, cherries and other <i>Prunus</i> spp., <i>Fraxinus americana</i> , <i>Gladiolus</i> , gooseberries, grapes, <i>Hydrangea</i> , peaches, <i>Pelargonium</i> , raspberries, <i>Rubus laciniatus</i> , strawberries. ToRSV also infects many common weeds in vineyards including common chickweed, dandelions, red clover and sheep sorrel (Schilder 2011).
<b>Distribution</b>	China, Canada, Egypt, Japan, Korea, USA (Fayek <i>et al.</i> 2009; EPPO 2010b).

## Glossary

Term or abbreviation	Definition
<b>Additional declaration</b>	A statement that is required by an importing country to be entered on a Phytosanitary Certificate and which provides specific additional information on a consignment in relation to regulated pests (FAO 2009).
<b>Appropriate level of protection</b>	The level of protection deemed appropriate by the Member establishing a sanitary or phytosanitary measure to protect human, animal or plant life or health within its territory (WTO 1995).
<b>Area</b>	An officially defined country, part of a country or all or parts of several countries (FAO 2009).
<b>Plant Biosecurity</b>	A branch within the Australian Government Department of Agriculture, Fisheries and Forestry, responsible for recommendations for the development of Australia's plant biosecurity policy.
<b>Certificate</b>	An official document which attests to the phytosanitary status of any consignment affected by phytosanitary regulations (FAO 2009).
<b>Consignment</b>	A quantity of plants, plant products and/or other articles being moved from one country to another and covered, when required, by a single Phytosanitary Certificate (a consignment may be composed of one or more commodities or lots) (FAO 2009).
<b>Control (of a pest)</b>	Suppression, containment or eradication of a pest population (FAO 2009).
<b>Endangered area</b>	An area where ecological factors favour the establishment of a pest whose presence in the area will result in economically important loss (FAO 2009).
<b>Entry (of a pest)</b>	Movement of a pest into an area where it is not yet present, or present but not widely distributed and being officially controlled (FAO 2009).
<b>Establishment</b>	Perpetuation, for the foreseeable future, of a pest within an area after entry (FAO 2009).
<b>Fruits and vegetables</b>	A commodity class for fresh parts of plants intended for consumption or processing and not for planting (FAO 2009).
<b>Host range</b>	Species capable, under natural conditions, of sustaining a specific pest or other organism (FAO 2009).
<b>Import Permit</b>	Official document authorising importation of a commodity in accordance with specified phytosanitary import requirements (FAO 2009).
<b>Import Risk Analysis</b>	An administrative process through which quarantine policy is developed or reviewed, incorporating risk assessment, risk management and risk communication.
<b>Infestation (of a commodity)</b>	Presence in a commodity of a living pest of the plant or plant product concerned. Infestation includes infection (FAO 2009).
<b>Inspection</b>	Official visual examination of plants, plant products or other regulated articles to determine if pests are present and/or to determine compliance with phytosanitary regulations (FAO 2009).
<b>Intended use</b>	Declared purpose for which plants, plant products, or other regulated articles are imported, produced, or used (FAO 2009).
<b>Interception (of a pest)</b>	The detection of a pest during inspection or testing of an imported consignment (FAO 2009).
<b>International Standard for Phytosanitary Measures</b>	An international standard adopted by the Conference of FAO [Food and Agriculture Organization], the Interim Commission on phytosanitary measures or the Commission on phytosanitary measures, established under the IPPC (FAO 2009).
<b>Introduction</b>	The entry of a pest resulting in its establishment (FAO 2009).
<b>National Plant Protection Organisation</b>	Official service established by a government to discharge the functions specified by the IPPC (FAO 2009).
<b>Official control</b>	The active enforcement of mandatory phytosanitary regulations and the application of mandatory phytosanitary procedures with the objective of eradication or containment of quarantine pests or for the management of regulated non-quarantine pests (FAO 2006).



Term or abbreviation	Definition
<b>Pathway</b>	Any means that allows the entry or spread of a pest (FAO 2009).
<b>Pest</b>	Any species, strain or biotype of plant, animal, or pathogenic agent injurious to plants or plant products (FAO 2009).
<b>Pest categorisation</b>	The process for determining whether a pest has or has not the characteristics of a quarantine pest or those of a regulated non-quarantine pest (FAO 2009).
<b>Pest Free Area</b>	An area in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this condition is being officially maintained (FAO 2009).
<b>Pest free place of production</b>	Place of production in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this condition is being officially maintained for a defined period (FAO 2009).
<b>Pest free production site</b>	A defined portion of a place of production in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this conditions is begin officially maintained for a defined period and that is managed as a separate unit in the same way as a pest free place of production (FAO 2009).
<b>Pest Risk Analysis</b>	The process of evaluating biological or other scientific and economic evidence to determine whether an organism is a pest, whether it should be regulated, and the strength of any phytosanitary measures to be taken against it (FAO 2009).
<b>Pest risk assessment (for quarantine pests)</b>	Evaluation of the probability of the introduction and spread of a pest and the magnitude of the associated potential economic consequences (FAO 2009).
<b>Pest risk management (for quarantine pests)</b>	Evaluation and selection of options to reduce the risk of introduction and spread of a pest (FAO 2009).
<b>Phytosanitary Certificate</b>	Certificate patterned after the model certificates of the IPPC (FAO 2009).
<b>Phytosanitary measure</b>	Any legislation, regulation or official procedure having the purpose to prevent the introduction and/or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests (FAO 2009).
<b>Phytosanitary regulation</b>	Official rule to prevent the introduction and/or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests, including establishment of procedures for phytosanitary certification (FAO 2009).
<b>Polyphagous</b>	Feeding on a relatively large number of host plants from different plant families.
<b>PRA area</b>	Area in relation to which a Pest Risk Analysis is conducted (FAO 2009).
<b>Quarantine pest</b>	A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled (FAO 2009).
<b>Regulated article</b>	Any plant, plant product, storage place, packaging, conveyance, container, soil and any other organism, object or material capable of harbouring or spreading pests, deemed to require phytosanitary measures, particularly where international transportation is involved (FAO 2009).
<b>Restricted risk</b>	Risk estimate with phytosanitary measure(s) applied.
<b>Rhizomes</b>	A horizontal plant stem with shoots above and roots below serving as a reproductive structure. Rhizomes may also be referred to as creeping rootstalks, or rootstocks
<b>Spread</b>	Expansion of the geographical distribution of a pest within an area (FAO 2009).
<b>Stakeholders</b>	Government agencies, individuals, community or industry groups or organizations, whether in Australia or overseas, including the proponent/applicant for a specific proposal, who have an interest in the policy issues.
<b>Unrestricted risk</b>	Unrestricted risk estimates apply in the absence of risk management measures.



## References

---

- Abdallah FB, Chebil S, Ghorbel A (2003) More about in vitro grape virus symptomatology. *Phytopathologia Mediterranea* 42: 35–40.
- Abelleira A, Mansilla JP, Padilla V, Hita I, Cabaleiro C, Bertolini E, Olmos A, Legorburu FJ (2010) First report of *Arabis mosaic virus* on grapevine in Spain. *Plant Disease* 94: 635–635.
- Abou Ghanem-Sabanadzovic N, Sabanadzovic S, Martelli GP (2003) Sequence analysis of the 3' end of three *Grapevine fleck virus*-like viruses from grapevine. *Virus Genes* 27: 11–16.
- Abou-Mansour E, Polier J, Pezet R, Tabacchi R (2009) Purification and partial characterisation of a 60 KDa laccase from *Fomitiporia mediterranea*. *Phytopathologia Mediterranea* 48: 447–453.
- Abu Yaman IK (1967) Population study of the grape leafhopper in Iraq. *Zeitschrift für Angewandte Entomologie* 60: 182–187.
- AFD (2008) Australian Faunal Directory. Australian Biological Resources Study. <http://www.environment.gov.au/biodiversity/abrs/online-resources/fauna/afd/> Accessed February 2008.
- Afonin AN, Greene SL, Dzyubenko NI, Frolov AN (2008) Interactive agricultural ecological atlas of Russia and neighboring countries. Economic plants and their diseases, pests and weeds. <http://www.agroatlas.ru> Accessed February 2010.
- AgroAtlas (2009a) Interactive agricultural ecological atlas of Russia and neighboring countries. Economic plants and their diseases, pests and weeds. Pests – *Agrotis segetum* Denis & Schiffermuller 1775. [http://www.agroatlas.ru/en/content/pests/Agrotis\\_segetum/](http://www.agroatlas.ru/en/content/pests/Agrotis_segetum/) Accessed July 2009.
- AgroAtlas (2009b) Interactive agricultural ecological atlas of Russia and neighboring countries. Economic plants and their diseases, pests and weeds. Pests – *Autographa gamma* L. [http://www.agroatlas.ru/en/content/pests/Autographa\\_gamma/](http://www.agroatlas.ru/en/content/pests/Autographa_gamma/) Accessed March 2009.
- AgroAtlas (2009c) Interactive agricultural ecological atlas of Russia and neighboring countries. Pests – *Melolontha melolontha* L. – Common Cockchafer [http://www.agroatlas.ru/en/content/pests/Melolontha\\_melolontha/](http://www.agroatlas.ru/en/content/pests/Melolontha_melolontha/) Accessed July 2009.
- AgroAtlas (2011a) Interactive agricultural ecological atlas of Russia and neighboring countries. Economic plants and their diseases, pests and weeds. Pests – *Eupoecilia ambiguella* (Hubner). [http://www.agroatlas.spb.ru/en/content/pests/Eupoecilia\\_ambiguella](http://www.agroatlas.spb.ru/en/content/pests/Eupoecilia_ambiguella) Accessed July 2011.
- AgroAtlas (2011b) Interactive agricultural ecological atlas of Russia and neighboring countries. Economic plants and their diseases, pests and weeds. Pests – *Sparganothis pilleriana* Den. et Schiff. – Grape leafroller, Vine tortrix Moth, Long-palpi tortrix, Leaf-rolling tortrix. [http://www.agroatlas.ru/en/content/pests/Sparganothis\\_pilleriana/](http://www.agroatlas.ru/en/content/pests/Sparganothis_pilleriana/) Accessed July 2011.
- Ahadiyat A, Zangeneh AM (2007) First Report of *Ectopsocus briggsi* and *Trichopsocus dalii* (Psocoptera: Psocomorpha: Ectopsocidae and Trichopsocidae) from Iran. *Florida Entomologist* 90: 790–791.

- Akbas B, Erdiller G (1993) Research on grapevine virus diseases and determination of their incidences in Ankara, Turkey. *Journal of Turkish Phytopathology* 22: 55–63.
- Al Rwahnih M, Daubert S, Golino D, Rowhani A (2009) Deep sequencing analysis of RNAs from a grapevine showing Syrah decline symptoms reveals a multiple virus infection that includes a novel virus. *Virology* 387: 395–401.
- Al Rwahnih M, Dave A, Anderson M, Uyemoto JK, Sudarshana MR (2012a) Association of a circular DNA virus in grapevines affected by red blotch disease in California. *Proceedings of the 17<sup>th</sup> Congress of ICVG, Davis, California, USA, October 7–14 2012* 104–105. Foundation Plant Services, University of California, Davis.
- Al Rwahnih M, Sudarshana MR, Uyemoto JK, Rowhani A (2012b) Complete genome sequence of a novel vitivirus isolated from grapevine. *Journal of Virology* 86(17): 9545.
- Alaniz S, León M, Vicent A, García-Jiménez J, Abad-Campos P, Armengol J (2007) Characterization of *Cylindrocarpon* species associated with black foot disease of grapevines in Spain. *Plant Disease* 91: 1187–1193.
- Alford DV (2007) *Pests of fruit crops: a colour handbook*. Manson Publishing, London
- AliNiazee MT (1974) Contribution to the bionomics of the grape leaffolder, *Desmia funeralis* (Hübner): A laboratory study with field observations. *Pacific Coast Entomological Society Memorandum* 50: 269–278.
- AliNiazee MT, Stafford EM (1972) Notes on the biology, ecology, and damage of *Platynota stultana* on grapes. *Journal of Economic Entomology* 65: 1042–1044.
- Allsopp PG (1980) The biology of false wireworms and their adults (soil-inhabiting Tenebrionidae) (Coleoptera): a review. *Bulletin of Entomological Research* 70: 343–379.
- Alma A, Davis RE, Vibio M, Danielli A, Bosco D, Arzone A, Bertaccini A (1996) Mixed infection of grapevines in northern Italy by phytoplasmas including 16S rRNA RFLP subgroup 16SrI-B strains previously unreported in this host. *Plant Disease* 80: 418–421.
- Amalfi M, Yombiyeni P, Decock C (2010) *Fomitiporia* in sub-Saharan Africa: morphology and multigene phylogenetic analysis support three new species from the Guineo-Congolian rainforest. *Mycologia* 102: 1303–1317.
- Andolfi A, Mugnai L, Luque J, Surico G, Cimmino A, Evidente A (2011) Phytotoxins produced by fungi associated with grapevine trunk diseases. *Toxins* 3: 1569–1605.
- Andret-Link P, Laporte C, Valat L, Ritzenthaler C, Demangeat G, Vigne E, Laval V, Pfeiffer P, Stussi-Garaud C, Fuchs M (2004) *Grapevine fanleaf virus*: still a major threat to the grapevine industry. *Journal of Plant Pathology* 86: 183–195.
- Angelotti F, Scapin CR, Tessmann DJ, Vida JB, Oliveira RR, Canteri MG (2008) Diagrammatic scale for assessment of grapevine rust. *Tropical Plant Pathology* 33: 439–443.
- Anon (2005) Pest management plans for major crops of Shaanxi project area, World Bank Agricultural Technology Project. [http://www-wds.worldbank.org/external/default/WDSContentServer/WDSP/IB/2005/02/25/000012009\\_20050225142337/Rendered/INDEX/E10100v20PMP0for0Shaanxi.txt](http://www-wds.worldbank.org/external/default/WDSContentServer/WDSP/IB/2005/02/25/000012009_20050225142337/Rendered/INDEX/E10100v20PMP0for0Shaanxi.txt) Accessed December 2011.
- Anon (2008) *Eumorpha achemon*. Spingidae of the United States.

<http://www.silkmoths.bizland.com/eachemon.htm> Accessed April 2008.

Anon (2011) *Thyridium vitis*. Bulletin of the Torrey Botanical Club  
<http://www.ebooksread.com/authors-eng/torrey-botanical-club/bulletin-of-the-torrey-botanical-club-goo/page-5-bulletin-of-the-torrey-botanical-club-goo.shtml> Accessed December 2011.

APHIS-USDA (2002) Importation of grapes (*Vitis* spp.) from Korea into the United States. A qualitative, pathway-initiated pest risk assessment. APHIS-USDA, Riverdale, MD, USA.  
[https://web01.aphis.usda.gov/oxygen\\_fod/fb\\_md\\_ppq.nsf/0/6a4c6eb951006d8685256e2a005ecdc0/\\$FILE/0058.pdf](https://web01.aphis.usda.gov/oxygen_fod/fb_md_ppq.nsf/0/6a4c6eb951006d8685256e2a005ecdc0/$FILE/0058.pdf) Accessed April 2011.

Aptroot A (1995) A monograph of Didymosphaeria. *Studies in Mycology* 37: 1–160.

AQSIQ (2006) Export technical reference information for Chinese grape, received on 20 July 2006. General Administration of Quality Supervision, Inspection and Quarantine of the People's Republic of China, Beijing.

AQSIQ (2007) Reply on supplementary information on pests in table grapes exported from China to Australia received on 19 March 2007. General Administration of Quality Supervision, Inspection and Quarantine of the People's Republic of China, Beijing.

Arnaud G, Malembic-Maher S, Salar P, Bonnet P, Maixner M, Marcone C, Boudon-Padieu E, Foissac X (2007) Multilocus sequence typing confirms the close genetic interrelatedness of three distinct Flavescence dorée Phytoplasma strain clusters and group 16SrV phytoplasmas infecting grapevine and alder in Europe. *Applied and Environmental Microbiology* 73: 4001–4010.

Arnold JW (1982) Larval hemocytes in Noctuidae (Insecta: Lepidoptera). *International Journal of Insect Morphology and Embryology* 11: 173–188.

Aroca A, Gramaje D, Armengol J, García-Jiménez J, Raposo R (2010) Evaluation of the grapevine nursery propagation process as a source of *Phaeoacremonium* spp. and *Phaeomoniella chlamydospora* and occurrence of trunk disease pathogens in rootstock mother vines in Spain. *European Journal of Plant Pathology* 126: 165–174.

Aroca A, Raposo R (2007) PCR-based strategy to detect and identify species of *Phaeoacremonium* causing grapevine diseases. *Applied and Environmental Microbiology* 73: 2911–2918.

Aroca A, Raposo R (2009) Pathogenicity of *Phaeoacremonium* species on grapevine. *Journal of Phytopathology* 157: 413–419.

Arpaia ML, Morse JG (1991) Citrus thrips *Scirtothrips citri* (Moulton) (Thysanoptera: Thripidae) scarring and navel orange fruit quality in California. *Journal of Applied Entomology* 111: 28–32.

Auger J, Aballay EE, Pinto CM, Pastenes VC (1992) Effect of *Grape fanleaf virus* (GFV) on growth and productivity of grapevine plants cv. Thompson Seedless. Original language title: Efecto del virus de la hoja en abanico (VHA) en el desarrollo y productividad de plantas de vid cv. Thompson Seedless. *Fitopatologia* 27: 85–89.

Babini AR, Bertaccini A (1982) Viral aggregates induced by a distinctive strain of *Strawberry latent ringspot virus* from grapevine. *Phytopathologische Zeitschrift* 104: 304–308.

- Backus EA, Serrano M, Ranger CM (2005) Mechanisms of hopperburn: an overview of insect taxonomy, behavior, and physiology. *Annual Review of Entomology* 50: 125–151.
- Bahcecioglu Z, Braun U, Kabaktepe S (2006) *Neoerysiphe rubiae* – a new powdery mildew species on *Rubia* cf. *tinctoria* from Turkey. *Nova Hedwigia* 83: 489–492.
- Balikai RA, Bagali AN, Ryagi YH (1999) Incidence of *Spodoptera litura* Fab. on grapevine, *Vitis vinifera* L. *Insect Environment* 5: 32.
- Barba M, Ferretti L, Pasquini G (2006) I giallumi della vite: un problema fitosanitario di rilevanza nazionale. *Informatore Fitopatologico* 56: 4–8.
- Barr ME (1990) Some dictyosporus genera and species of Pleosporales in North America. *Memoire of the New York Botanical Garden* 62: 1–92.
- Batlle A, Angeles-Martinez M, Lavina A (2000) Occurrence, distribution and epidemiology of Grapevine yellows in Spain. *European Journal of Plant Pathology* 106: 811–816.
- BCCM (2012) BCCM//MUCL fungi & yeasts catalogue – species details. [http://bccm.belspo.be/db/mucl\\_species\\_details.php?NUM=12093&FIRSTITEM=1&RANGE=100](http://bccm.belspo.be/db/mucl_species_details.php?NUM=12093&FIRSTITEM=1&RANGE=100) Accessed March 2012.
- BCMAL (2010) Western grape rootworm, *Bromius obscurus* – British Columbia Ministry of Agriculture and Lands Pest Alert. [http://www.agf.gov.bc.ca/cropprot/bromius\\_obscurus.pdf](http://www.agf.gov.bc.ca/cropprot/bromius_obscurus.pdf). Accessed March 2012.
- Beardsley JW, Gonzalez RH (1975) The biology and ecology of armored scales. *Annual Review of Entomology* 20: 47–73.
- Belli G, Fortusini A, Vegetti G (1982) *Arabis mosaic virus* isolated from grapevine in Italy. *Rivista di Patologia Vegetale* 18: 175–177.
- Ben-Dov Y (1994) *A systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with data on geographical distribution, host plants, biology and economic importance*. Intercept Limited, UK.
- Ben-Dov Y, Miller DR, Gibson GAP (2012) ScaleNet. <http://www.sel.barc.usda.gov/scalenet>. Accessed February 2012.
- Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Wheeler DL (2008) GenBank. *Nucleic Acids Research* 36: D25–D35.
- Bentley WJ, Varela LG, Zalom FG, Smith RJ, Purcell AH, Phillips PA, Haviland DR, Daane KM, Battany MC, Granett J (2008) UC IPM Pest Management Guidelines: Grape: UC ANR Publication 3448 <http://www.ipm.ucdavis.edu/PMG/r302301411.html> Accessed February 2011.
- Benvenuti A, Lucchi A (2005) Corso di Laurea in Viticoltura ed Enologia. Il crisomelide alticino della vite *Altica ampelophaga* Guérin-Meneville: cronologia, biologia, dannosità e controllo' University of Pisa, Italy (Italian). [http://www.agr.unipi.it/tesi05/luglio/ve\\_benvenuti.html](http://www.agr.unipi.it/tesi05/luglio/ve_benvenuti.html) Accessed February 2010.
- Bercks R (1967) Ueber den Nachweis des Tomatenzwergbusch-Virus (*Tomato bushy stunt virus*) in Reben. *Phytopathologische Zeitschrift* 60: 273–277.
- Bercks R, Brückbauer Querfurth G, Rüdel M (1977) Untersuchungen über die viruskrankheit

de rebe unter besonderer Berücksichtigung ‘atypischer formen’ der Reigigkrankheit. *Weinbau und Kell.* 24: 133–180.

Bercks R, Querfurth G (1969) On the detection of *Sowbane mosaic virus* in grapevines. *Phytopathologische Zeitschrift* 66: 365–373.

Berger J, Dalla VJ, Baric S (2009) Development of a TaqMan allelic discrimination assay for the distinction of different subtypes of the grapevine yellow phytoplasma Bois noir. *European Journal of Plant Pathology* 124: 521–526.

Bernard MB, Horne PA, Hoffmann AA (2005) Eriophyoid mite damage in *Vitis vinifera* (grapevine) in Australia: *Calepitrimerus vitis* and *Colomerus vitis* (Acari: Eriophyidae) as the common cause of the widespread ‘Restricted Spring Growth’ syndrome. *Experimental and Applied Acarology* 35: 83–109.

Berry RE (1998) *Wireworms: Insects and Mites of Economic Importance in the Northwest*. 2<sup>nd</sup> Edn. Oregon State University, Oregon.

Bertaccini A, Vibio M, Schaff DA, Murari E, Martini M, Danielli A (1997) Geographical distribution of Elm yellows-related phytoplasmas in grapevine Flavescence dorée outbreaks in Veneto (Italy). In *Extended Abstracts of the 12th Meeting of the International Council for the Study of Virus and Virus-like Diseases of the Grapevine, Lisbon, 28 September–2 October 1997*. Departamento de Fitopatologia, Estação Agronómica Nacional, Oeiras, Portugal.

Bertaccini A, Vibio M, Stefani E (1995) Detection and molecular characterization of phytoplasmas infecting grapevine in Liguria (Italy). *Phytopathologia Mediterranea* 34: 137–141.

Bhuiyan MK, Nishigaki J (1997) Oviposition of the adult cupreous chafer, *Anomala cuprea* hope (Coleoptera: Scarabaeidae), at different water contents of the ovipositing medium under laboratory conditions. *Applied Entomology and Zoology* 32: 431–436.

Bianco PA, Davis RE, Casati P, Fortusini A (1996) Prevalence of Aster yellow (AY) and Elm yellows (EY) group phytoplasmas in symptomatic grapevines in three areas of northern Italy. *Vitis* 35: 195–199.

Bianco PA, Davis RE, Prince JP, Fortusini A, Casati P, Belli G (1994) Elm yellows and Aster yellows MLOs associated with a grapevine yellows disease very similar to Flavescence dorée in Northern Italy. *International Organization for Mycoplasma Letters* 3: 251–252.

Blodgett S (1992) Ground Mealybug. In: *Grape Pest Management*, 2nd edition. Publication 3343: 247–248. University of California Division of Agriculture and Natural Resources, Oakland, CA, USA.

Boerema GH (1976) The *Phoma* species studied in culture by Dr R WG Dennis. *Transactions of the British Mycological Society* 67: 289–319.

Boll S, Herrmann JV (2001) A new method to monitor eggs of the Grape leafhopper (*Empoasca vitis*) in grapevine leaves. *Journal of Plant Diseases and Protection* 108: 77–81.

Booth RG, Cox ML, Madge RB (1990) *IIE Guides to insects of importance to man: 3. Coleoptera*. Cambridge University Press, Cambridge, UK.

Borbon CM, de Becerra V, Bonomo V, Mazzitelli E, Calvo M (2008) Thrips (Insecta: Thysanoptera) in cherry orchards in Mendoza, Argentina. *Revista de la Facultad de Ciencias*

Agrarias, Universidad Nacional de Cuyo 40: 1–10.

Borden AD, Madsen HF (1951) Stink bug on pears: Habits of pest studied to find a control program which may include sprays, clean culture, host plant eradication. *Californian Agriculture* 5: 7.

Bos L (1999) *Plant viruses, unique and intriguing pathogens*. Backhuys Publishers, Leiden.

Boscia D, Digiario M, Savino V, Martelli GP (2000) A survey of *Grapevine leafroll-associated virus 6* distribution in Apulia and Abruzzo (southern Italy) and its association with cv. Cardinal. *Vignevini* 27: 92–95.

Bostanian NJ, Vincent C, Goulet H, Lesage L, Lasnier J, Bellemare J, Mauffette Y (2003) The arthropod fauna of Quebec vineyards with particular reference to phytophagous arthropods. *Journal of Economic Entomology* 96: 1221–1229.

Botero-Garces N, Isaacs R (2003) Distribution of Grape berry moth, *Endopiza viteana* (Lepidoptera: Tortricidae) in natural and cultivated habitats. *Environmental Entomology* 32: 1187–1195.

Botha J, Hardie D, Power G (2000) *Grape phylloxera: exotic threat to Western Australia*. Factsheet 0002–2000. Department of Agriculture, Western Australia.

Botha WJ, Serfontein S, Greyling MM, Berger DK (2001) Detection of *Xylophilus ampelinus* in grapevine cuttings using a nested polymerase chain reaction. *Plant Pathology* 50: 515–526.

Botti S, Bertaccini A (2007) Grapevine yellows in Northern Italy: molecular identification of Flavescence dorée Phytoplasma strains and of Bois noir Phytoplasmas. *Journal of Applied Microbiology* 103: 2325–2330.

Boudon-Padieu E, Larrue J, Caudwell A (1989) ELISA and dot-blot detection of Flavescence dorée MLO in individual leafhopper vectors during latency and inoculative state. *Current Microbiology* 19: 357–364.

Boudon-Padieu E, Maixner M (2007) Potential effects of climate change on distribution and activity of insect vectors of grapevine pathogens. Global warming, which potential impacts on the vineyards?

[http://www.prodinra.inra.fr/prodinra/pinra/data/2008/06/PROD2008d0be8eb\\_20080625041803219.pdf](http://www.prodinra.inra.fr/prodinra/pinra/data/2008/06/PROD2008d0be8eb_20080625041803219.pdf) Accessed January 2012.

Boudon-Padieu E. (2005) Phytoplasmas associated to Grapevine yellows and potential vectors, *Bulletin O.I.V.* 79: 311–320.

Bournier A (1976) Grape insects. *Annual Review of Entomology* 22: 355–376.

Bovey R, Gärtel W, Hewitt WB, Martelli GP, Vuittenez A (1990) *Viruses and virus-like diseases of grapevines* 46–50. Editions Payot, Lausanne, Switzerland.

Brayford D (1987) CMI Descriptions of Pathogenic Fungi and Bacteria no. 926. *Cylindrocarpon lichenicola*. *Mycopathologia* 100: 125–126.

Briceño EX, Latorre BA (2008) Characterization of Cladosporium rot in grapevines, a problem of growing importance in Chile. *Plant Disease* 92: 1635–1642.

Brooks FE (2002) Brown root rot disease in American Samoa's tropical rain forests. *Pacific Science* 56: 377–387.

- Brown DJF, Kyriakopoulou PE, Roberston WM (1997) Frequency of transmission of *Artichoke Italian latent nepovirus* by *Longidorus fasciatus* (Nematoda: Longidoridae) from artichoke fields in Iria and Kandia areas of Argolis in northeast Peloponnesus, Greece. *European Journal of Plant Pathology* 103: 501–506.
- Brown JW (1999) A new genus of tortricid moths (Tortricidae: Euliini) injurious to grapes and stone fruits in Chile. *Journal of the Lepidopterists Society* 53: 60–64.
- Brown JW, Passoa S (1998) Larval food plants of Euliini (Lepidoptera: Tortricidae): from *Abies* to *Vitis*. *Pan Pacific Entomologist* 74: 1–11.
- Brown SL, Hudson W (2005) Green June Beetle – *Cotinis nitida* (Linnaeus). The Bugwood Network. <http://www.gaipm.org/top50/gjb.html> Accessed May 2008.
- Brunt AA, Teakle DS (1996) *Tobacco necrosis necrovirus*. In Brunt AA, Crabtree K, Dallwitz MJ, Gibbs A, Watson L (eds) *Viruses of plants: descriptions and lists from the VIDE database* 1256–1259. CAB International, Wallingford.
- Brunt AA, Crabtree K, Dallwitz MJ, Gibbs AJ, Watson L, Zurcher EJ (1996) `Plant Viruses Online: Descriptions and Lists from the VIDE Database. <http://biology.anu.edu.au/Groups/MES/vid/>. Accessed June 2012.
- Buchanan GA, Furness GO, Charles JG (1994) Grape Phylloxera. In Nicholas P, Margery P, Wachtel M (eds) *Diseases and pests*. Winetitles, Adelaide.
- Buchanan GA, Whiting JR (1991) Phylloxera management: prevention is better than cure. *Australian and New Zealand Wine Industry Journal* 6: 223–230.
- Büchen-Osmond (ed) (2006) *ICTVdB – The Universal Virus Database*. Version 4. <http://www.ncbi.nlm.nih.gov/ICTVdb/ICTVdB> Accessed April 2013.
- Büchen-Osmond C, Crabtree K, Gibbs A, McLean G (1988) *Viruses of plants in Australia: descriptions and lists from the VIDE database*. Australian National University, Canberra.
- Burdsall HH, Jr. (1985) A contribution to the taxonomy of the genus *Phanerochaete* (Corticaceae, Aphyllophorales). *Mycological Memories* 10: 1–165.
- Burgess TI, Barber PA, Hardy GE St J (2005) *Botryosphaeria* spp. associated with eucalypts in Western Australia, including the description of *Fusicoccum macroclavatum* sp. nov. *Australasian Plant Pathology* 34: 557–567.
- Burgess TI, Barber PA, Mohali S, Pegg G, de Beer W, Wingfield MJ (2006) Three new *Lasiodiplodia* spp. from the tropics, recognized based on DNA sequence comparisons and morphology. *Mycologia* 98: 423–435.
- Burts EC, Dunley J (1993) Grape mealybug. Orchard Pest Management Online. Washington State University, Washington. <http://jenny.tfrec.wsu.edu/opm/displayspecies.php?pn=130> Accessed April 2013.
- CABI (2012a) Crop Protection Compendium, CABI International, Wallingford, UK. <http://www.cabi.org/cpc/> Accessed 12 March 2012.
- CABI (2012b) Species Fungorum. CABI International, Wallingford, UK. <http://www.speciesfungorum.org/> Accessed 1 August 2012.
- CABI/EPPO (1990) *Xylella fastidiosa*. Data sheets on quarantine pests. Wallingford, UK,

CAB International.

CABI/EPPO (1999) *Xylophilus ampelinus*. Distribution Maps of Plant Diseases, Map No. 531. CAB International, Wallingford, UK.

Cabrera BJ, Scheffrahn RH (2005) Western drywood termite – *Incisitermes minor* (Hagen) (Insecta: Isoptera: Kalotermitidae) EENY-248 Featured Creatures, Department of Entomology and Nematology, Division of Plant Industry University of Florida.

[http://entnemdept.ufl.edu/creatures/urban/termites/western\\_drywood\\_termite.htm](http://entnemdept.ufl.edu/creatures/urban/termites/western_drywood_termite.htm) Accessed August 2011.

Cadman CH, Dias HF, Harrison BD (1960) Sap transmissible viruses associated with diseases of grapevine in Europe and North America. *Nature* 187: 577–579.

CAES (2008) *Pestalotia*. College of Agricultural and Environmental Sciences, University of Georgia. <http://plantpath.caes.uga.edu/extension/Fungi/pestalotia.html> Accessed May 2011.

Cahill DM, Rookes JE, Wilson BA, Gibson L, McDougall KL (2008) *Phytophthora cinnamomi* and Australia's biodiversity: impacts, predictions and progress towards control. *Australian Journal of Botany* 56: 279–310.

Cammell ME (1981) The black bean aphid, *Aphis fabae*. *Biologist* 28: 247–258.

Campos SL, Faccin MA, Echeverría MN, Sazo RL (1981) Distribución y ciclo evolutivo del tortricido enrollador de la vid *Proeulia auraria* (Clarke). *Agricultura Técnica* 41: 246–256.

Canik D, Ertunc F, Paltrinieri S, Contaldo N, Bertaccini A (2011) Identification of different phytoplasmas infecting grapevine in Turkey. *Bulletin of Insectology* 64: S225–S226.

Cannizzaro G, Rosciglione B, Castellano MA (1990) The presence of phytopathogenic viruses in waterways of western Sicily. *Informatore Fitopatologico*. 40: 55–56.

Capinera JL (2008) Yellowstriped armyworm, *Spodoptera ornithogalli* (Guenee) (Insecta: Lepidoptera: Noctuidae). IFAS Extension, University of Florida, EENY 216.

<http://edis.ifas.ufl.edu/pdf/IN/IN37300.pdf> Accessed April 2010.

Capinera JL, Fasulo TR (2006) *Spodoptera exigua* (Hübner) (Insecta: Lepidoptera: Noctuidae). IFAS Extension, University of Florida, EENY 105

[http://entnemdept.ufl.edu/creatures/veg/leaf/beet\\_armyworm.htm](http://entnemdept.ufl.edu/creatures/veg/leaf/beet_armyworm.htm) Accessed March 2012.

Cardoso JMS, Félix MR, Clara MIE, Oliveira S (2005) The complete genome sequence of a new necrovirus isolated from *Olea europaea* L. *Archives of Virology* 150: 815–823.

Carew ME, Goodisman MAD, Hoffmann AA (2004) Species status and population structure of grapevine eriophyoid mites (Acari: Eriophyidae). *Entomologia experimentalis et applicata* 111: 87–96.

Carlucci A, Frisullo S (2009) First report of *Diplodia corticola* on grapevine in Italy. *Journal of Plant Pathology* 91: 233.

Carpenter D, Luckett D (2003) *Cucumber mosaic virus* in lupins.

[http://www.dpi.nsw.gov.au/\\_data/assets/pdf\\_file/0005/157433/pulse-point-18.pdf](http://www.dpi.nsw.gov.au/_data/assets/pdf_file/0005/157433/pulse-point-18.pdf) Accessed September 2011.

Carter DJ (1984) *Pest Lepidoptera of Europe: with special references to the British Isles*. Dr W Junk Publishers, London.



- Cartwright D (2009) *Tomato ring spot virus*, 29th September 2009 (personal communication). Primary Industries and Resources, South Australia.
- Castillo-Pando M, Somers A, Green CD, Priest M, Sriskanthades M (2001) Fungi associated with dieback of Semillon grapevines in the Hunter Valley of New South Wales. *Australasian Plant Pathology* 30: 59–63.
- Castillo-Pando M, Steel C, Somers T (1999) Fungal bunch rots of grapes in the Hunter and Hastings Valleys. *The Australian Grapegrower and Winemaker* September: 33–36.
- Catal M, Jordan SA, Butterworth SC, Schilder AMC (2007) Detection of *Eutypa lata* and *Eutypella vitis* in grapevine by nested multiplex polymerase chain reaction. *Phytopathology* 97: 737–747.
- Caudwell A (1961) Les phénomènes de rétablissement chez la Flavescence dorée de la vigne. *Annales des Epiphyties* 12: 347–354.
- Caudwell A, Larrue J, Boudon-Padiou E, McLean GD (1997) Flavescence dorée elimination from dormant wood of grapevines by hot-water treatment. *Australian Journal of Grape and Wine Research* 3: 21–25.
- Caudwell A, Larrue J, Tassart V, Boidron R, Grenan S, Leguay M, Bernard P (1994) Ability of grapevine rootstock varieties to transmit flavescence dorée. Study of the case of 3309 C and Fercal. *Agronomie* 14: 83–94.
- Cesati RR, Van Regenmortel MHV (1969) Serological detection of a strain of *Tobacco necrosis virus* in grapevine leaves. *Journal of Phytopathology* 64: 362–366.
- Chabbouh N, Martelli GP, Savino V, Greco N, Laforteza R (1993) Potato virus X in Tunisian grapevines. *Vitis* 32: 165–169.
- Chalkley D (2011) Grape leaf rust: *Phakopsora euvitidis*. Invasive Fungi. Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.ars-grin.gov/taxadescriptions/factsheets/pdfPrintFile.cfm?thisApp=Phakopsoraeuvitis> Accessed January 2012.
- Chang CP (1988) The ecology and control of insects and other animal pests on grapevine. *Chinese Journal of Entomology* 2: 11–31.
- Chatasiri S, Ono Y (2008) Phylogeny and taxonomy of the Asian grapevine leaf rust fungus, *Phakopsora euvitidis*, and its allies (Uredinales). *Mycoscience* 49: 66–74.
- Chee KH (1976) Assessing susceptibility of *Hevea* clones to *Microcyclus ulei*. *Annals of Applied Biology* 84: 135–145.
- Cheng CP (1984) The morphology and life history of the Coffee borer (*Zeuzera coffeae* Nietner) on grapevine in Taiwan. *Plant Protection Bulletin, Taiwan* 26: 145–153.
- Cherepanov AI (1991) *Cerambycidae of northern Asia: Laminae*. Nauka Publishers, New Delhi.
- Chicau G, Aboim-Inglez M, Cabral S, Cabral JPS (2000) *Phaeoacremonium chlamydosporum* and *Phaeoacremonium angustius* associated with esca and grapevine decline in Vinho Verde grapevines in northwest Portugal. *Phytopathologia Mediterranea* 39: 80–86.
- Cho IS, Jung SM, Cho JD, Choi GS, Lim HS (2013) First report of *Grapevine pinot gris virus* infecting grapevine in Korea. *New Disease Reports* 27: 10.

- Chu PWG, Francki RIB, Hatta T (1983) Some properties of *Tomato ringspot virus* isolated from *Pentas lanceolata* in South Australia. *Plant Pathology* 32: 353–356.
- Ciccarone C, Graniti A, Schiaffino A, Marras F (2004) Molecular analysis of *Fomitiporia mediterranea* isolates from esca-affected grapevines in southern Italy. *Phytopathologia Mediterranea* 43: 268–272.
- Cigsar I, Digiario M, Gokalp K, Ghanem-Sabanadzovic NA, de Strandis A, Boscia D, Martelli GP (2003) *Grapevine deformation virus*, a novel Nepovirus from Turkey. *Journal of Plant Pathology* 85: 183–191.
- Cigsar I, Digiario M, Martelli GP (2002) Sanitary status of grapevines in South-Eastern and Central Anatolia (Turkey). *Bulletin OEPP/EPPO Bulletin* 32: 471–475.
- Cirami RM, van Velsen RJ, Niejalke J (1988) Grapevine virus indexing in the South Australian Vine Improvement Scheme, 1974–1987. *Australian Journal of Experimental Agriculture* 28: 645–649.
- Cleveland TC (1982) Hibernation and host plant sequence studies of Tarnished plant bugs, *Lygus lineolaris*, in the Mississippi Delta. *Environmental Entomology* 11: 1049–1052.
- Cline ET, Farr DF (2006) Synopsis of fungi listed as regulated plant pests by the USDA Animal and Plant Health Inspection Service: notes on the nomenclature, disease, plant hosts, and geographic distribution. *Plant Health Progress*.  
<http://ddr.nal.usda.gov/bitstream/10113/11908/1/IND43838170.pdf> Accessed May 2012.
- Coetzee B, Maree HJ, Stephan D, Freeborough M-J, Burger JT (2010) The first complete nucleotide sequence of a *Grapevine virus E* variant. *Archives of Virology* 155: 1357–1360.
- Coller GJ Van, Denman S, Groenewald JZ, Lamprecht SC, Crous PW (2005) Characterisation and pathogenicity of *Cylindrocladiella* spp. associated with root and cutting rot symptoms of grapevines in nurseries. *Australasian Plant Pathology* 34: 489–498.
- Common IFB (1990) *Moths of Australia*. Melbourne University Press, Melbourne.
- Constable F, Rodoni B (2011a) Fact Sheet: *Grapevine fleck* and associated viruses. Grape and Wine Research and Development Corporation.  
[http://www.gwrdc.com.au/webdata/resources/files/GWR\\_081\\_Fleck\\_and\\_Associated\\_Viruses\\_Fact\\_Sheet\\_FINAL.pdf](http://www.gwrdc.com.au/webdata/resources/files/GWR_081_Fleck_and_Associated_Viruses_Fact_Sheet_FINAL.pdf) Accessed November 2011.
- Constable F, Rodoni B (2011b) Rugose wood – associated viruses.  
[http://www.gwrdc.com.au/webdata/resources/files/GWR\\_080\\_Rugose\\_Wood\\_Disease\\_Fact\\_Sheet\\_PAGES\\_FINAL.pdf](http://www.gwrdc.com.au/webdata/resources/files/GWR_080_Rugose_Wood_Disease_Fact_Sheet_PAGES_FINAL.pdf) Accessed September 2011.
- Constable FE (2010) Phytoplasma Epidemiology: Grapevines as a model. In Weintraub PG, Jones P (eds) *Phytoplasmas: Genomes, Plant Hosts and vectors*. CAB International, Wallingford, UK.
- Constable FE, Drew C (2004) Review of vine health parameters, implementation priorities and capabilities for vine improvement groups and accredited nurseries. Grape and Wine Research and Development Corporation, Australian Government.  
<http://www.gwrdc.com.au/webdata/resources/project/NVH0301.pdf> Accessed March 2011.
- Constable FE, Nicholas P, Rodoni BC (2010) *Development and validation of diagnostic protocols for the detection of endemic and exotic pathogens of grapevines*. Department of

Primary Industries, Victoria.

Contaldo N, Soufi Z, Bertaccini A (2011) Preliminary identification of phytoplasmas associated with grapevine yellows in Syria. *Bulletin of Insectology* 64: 217–218.

Cook RP, Dubé AJ (1989) *Host-pathogen index of plant disease in South Australia*. Field Crops Pathology Group, South Australian Department of Agriculture, South Australia.

Coombe BG, Dry PR (1992) *Viticulture* Volume 2: Practices. Winetitles, Adelaide.

Coombs M, Khan SA (1998) Population levels and natural enemies of *Plautia affinis* Dallas (Hemiptera: Pentatomidae) on raspberry, *Rubus idaeus* L., in south-eastern Queensland. *Australian Journal of Entomology* 37: 125–129.

Cortesi P, Fischer M, Milgroom MG (2000) Identification and spread of *Fomitiporia punctata* associated with wood decay of grapevine showing symptoms of esca. *Phytopathology* 90: 967–972.

Coutts RH, Rigden JE, Slabas AR, Lomonosoff GP and Wise PJ (1991) The complete nucleotide sequence of *Tobacco necrosis virus strain D*. *Journal of General Virology* 72: 1521–1529.

Cox JM (1989) The mealybug genus *Planococcus* (Homoptera: Pseudococcidae). *Bulletin of the British Museum Natural History Entomology* 58: 1–78.

CPPDR (1994) *Black vine thrips, Retithrips syriacus*. The California Plant Pest and Disease Report (CPPDR) Volume 13 (3–4). CFDA, California.

Crous P, Gams W, Wingfield MJ, van Wyk PS (1996) *Phaeoacremonium* gen. nov. associated with wilt and decline diseases of woody hosts and human infections. *Mycologia* 88: 786–796.

Crous PW, Swart L, Coertze S (2001) The effect of hot water treatment on fungi occurring in apparently healthy grapevine cuttings. *Phytopathologia Mediterranea* 40: S464–S466.

CSIRO (1991) *Insects of Australia* Volume 1, 2<sup>nd</sup> edn. Melbourne University Press, Carlton, Australia.

CSIRO (2005) Australian Insect Common Name. Commonwealth Scientific and Industrial Research Organisation. <http://www.ento.csiro.au/aicn/index.htm> Accessed April 2011.

CSIRO (2013) Australian moths online: a photo gallery. Commonwealth Scientific and Industrial Research Organisation. <http://www1.ala.org.au/gallery2/main.php> Accessed April 2013.

Dahiya KK, Lakra RK (2001) Seasonal occurrence and succession of thrips, *Rhipiphorothrips cruentatus* Hood in important horticultural crops of Haryana. *Crop Research Hisar* 21: 112–114.

Daire X, Clair D, Reinert W, Boudon-Padieu E (1997) Detection and differentiation of grapevine yellows phytoplasmas belonging to Elm yellows group and to the Stolbur subgroup by PCR amplification of non-ribosomal DNA. *European Journal of Plant Pathology* 103: 507–514.

Damm U, Fourie PH, Crous PW (2007) *Aplosporella prunicola*, a novel species of anamorphic Botryosphaeriaceae. *Fungal Diversity* 27: 35–43.

- Davis RE, Dally EL (2001) Revised subgroup classification of group 16SrV phytoplasmas and placement of Flavescence dorée-associated phytoplasmas in two distinct subgroups. *Plant Disease* 85:790–797.
- Davis RE, Jomantiene R, Dally EL, Wolf TK (1998) Phytoplasmas associated with Grapevine yellows in Virginia belong to group 16SrI, subgroup A (Tomato big bud phytoplasma subgroup), and group 16SrIII, new subgroup I. *Vitis* 37: 131–137.
- De Andrade ER, Dalbó MA, Schuck E, Gallotti GJM (1995) Evaluation of grapevine (*Vitis* spp.) resistance to *Fusarium oxysporum* f.sp. *herbemontisin* rio do peixe valley, Santa Catarina state, Brazil. *Acta Horticulturae* 388: 65–70.
- De Klerk CA (1987) Chemical control of *Margarodes prieskaensis* (Jakubski) (Coccoidae: Margarodidae) on grape vines. *South African Journal for Enology and Viticulture* 8: 11–15.
- De Klerk CA (2010) Chemical control of male pre-pupae and adult females of *Margarodes prieskaensis* (Jakubski) (Coccoidae: Margarodidae) on grape vines. *South African Journal for Enology and Viticulture* 31: 160–164.
- Deighton FC (1976) Three fungi on leaves of *Vitis*. *Transactions British Mycological Society* 67: 223–232.
- Delibašić G, Babović M, Jakovljević D (2000) The appearance and distribution of *Grapevine fanleaf virus* and *Arabis mosaic virus* in Yugoslavia. *Extended abstracts International Council for the study of viruses and virus-like diseases of the grapevine* 72–73 Adelaide, Australia.
- Deng S and Hiruki D (1991) Amplification of 16S rRNA genes from culturable and nonculturable mollicutes. *Journal of Microbiological Methods* 14: 53–61.
- Dennehy TJ, Clark LG (2008) Integrated Pest Management: Grape rootworm. Cornell University, New York State Agricultural Experiment Station, USA.  
<http://www.nysipm.cornell.edu/factsheets/grapes/pests/grw/grw.asp> Accessed February 2010.
- Dennill GB (1991) A pruning technique for saving vineyards severely infested by the Grape vine bud mite *Colomerus vitis* (Pagenstecher) (Eriophyidae). *Crop Protection* 10: 310–314.
- DeSousa E, Cardoso F, Casati P, Bianco PA, Guimaraes M and Pereira V (2003) Detection and identification of phytoplasmas belonging to 16SrV-D in *Scaphoideus titanus* adults. In *Extended Abstracts of the 14th ICVG Conference Locorotondo, 12–17 September 2003, Portugal* 78. ICVG, Switzerland.
- Diaz GA, Prehn D, Latorre BA (2011) First report of *Cryptovalsa ampelina* and *Eutypella leprosa* associated with grapevine trunk diseases in Chile. *Plant Disease* 95: 490–491.
- Dibble JE, Joose J, LaVine P, Haire S and Bearden BE (1979) Climbing cutworms; early-season pests of grapes. *California Agriculture* 33: 14–15.
- Digiario M, Elbeaino T, Martelli GP (2007) Development of degenerate and species-specific primers for the differential and simultaneous RT-PCR detection of grapevine-infecting nepoviruses of subgroups A, B and C. *Journal of Virological Methods* 141: 34–40.
- Digiario M, Ghanem-Sabanadzovic NA, Cigsar I, Gokalp K, de Strandis A, Boscia D, Martelli GP (2003) Two hitherto undescribed nepoviruses from Turkish grapevines. In *Extended Abstracts of the 14th ICVG Conference, Locorotondo, 12–17th September, 2003*, North

American Plant Protection Organization (NAPPO), Ottawa.

Digiario M, Martelli GP, Savino V (2000) Phloem-limited viruses in the Mediterranean and Near East. In *Extended Abstracts of the 13th Meeting of ICVG, March 12–18, 2000 Adelaide, Australia*. ICVG, Switzerland.

Dimou D, D'Onghia AM, da Camara Machado ML, Savino V (1994) Occurrence of *Grapevine chrome mosaic nepovirus* in Austria. *Journal of Phytopathology* 142: 258–262.

Doganlar M, Yigit A (2002) A new potential pest for orchards and vineyards: Black vine thrips, *Retithrips syriacus* (Mayet) (Thysanoptera: Thripidae) in Hatay. *Turkiye Entomoloji Dergisi* 26: 283–94.

Dovas CI, Katis NI (2003) A spot nested RT-PCR method for the simultaneous detection of members of the *Vitivirus* and *Foveavirus* genera in grapevine. *Journal of Virological Methods* 107: 99–106.

Dreaden TJ, Shin K, Smith JA (2011) First report of *Diplodia corticola* causing branch cankers on live oak (*Quercus virginiana*) in Florida. *Plant Disease* 95: 1027.

Dreo T, Seljak G, Janse JD, Beld Ivan der, Tjou-Tam-Sin L, Gorkink-Smits P, Ravnikaar M, (2005) First laboratory confirmation of *Xylophilus ampelinus* in Slovenia. *Bulletin OEPP* 35: 149–155.

Duduk B, Ivanovic M, Dukic N, Botti S, Bertaccini A (2003) First report of an Elm yellows subgroup 16SrV-C Phytoplasma infecting grapevine in Serbia. *Plant Disease* 87: 599.

Duduk B, Tian JB, Contaldo N, Fan XP, Paltrinieri S, Chen QF, Zhao QF, Bertaccini A (2010) Occurrence of phytoplasmas related to Stolbur and to ‘*Candidatus* Phytoplasma japonicum’ in woody host plants in China. *Journal of Phytopathology* 158: 100–104.

Dugan FM, Schubert K, Braun U (2004) Check-list of *Cladosporium* names. *Schelechtendalia* 11: 1-119.

Dupont J, Laloui W, Magnin S, Larignon P, Roquebert MF (2000) *Phaeoacremonium viticola*, a new species associated with esca disease of grapevine in France. *Mycologia* 92: 499–504.

Eaton AT (1978) Studies on distribution patterns, ovipositional preference, and egg and larval survival of *Colaspis brunnea* (Fab.) in North Carolina Coastal Plain soybean fields. Thesis, North Carolina State University. [http://www.ent.iastate.edu/soybeaninsects/Eaton\\_1978](http://www.ent.iastate.edu/soybeaninsects/Eaton_1978) Accessed March 2011.

Ebel R, Schnabel A, Reustle GM, Krczal G, Wetzel T (2003) Molecular characterization of two German *Raspberry ringspot virus* isolates infecting grapevines and construction of full length infectious clones. In *Extended Abstracts of the 13th Meeting of ICVG, March 12–18, 2000 Adelaide, Australia* 16. ICVG, Switzerland.

Ebeling W (2002) Urban Entomology: Wood-destroying insects and fungi. Department of Entomology. University of California, Riverside.  
<http://entomology.ucr.edu/ebeling/ebel51.html#termite%20pest%20species%20list> Accessed April 2010.

Eddin G, Al-Chaabi MS, Khadam A (2008) Investigation on some *Grapevine leafroll-associated viruses* (GLRaVs) in south of Syria. *Arab Journal of Plant Protection* 26: 102–

109.

Edwards J, Pascoe IG (2004) Occurrence of *Phaeomoniella chlamydospora* and *Phaeoacremonium aleophilum* associated with Petri disease and esca in Australian grapevines. *Australasian Plant Pathology* 33: 273–279.

Eicker A (1973) The mycoflora of *Eucalyptus maculata* leaf litter. *Soil Biology and Biochemistry* 5: 441–448.

Ellis JB, Everhart BM (1897) New species of fungi from various localities. *Bulletin of the Torrey Botanical Club* 24: 457–477.

Ellis MA (2008) Grape black rot. The Ohio State University Extension Factsheet. [ohioline.osu.edu/hyg-fact/3000/pdf/HYG\\_3004\\_08.pdf](http://ohioline.osu.edu/hyg-fact/3000/pdf/HYG_3004_08.pdf) Accessed November 2010.

Ellis MB, Ellis JP (1997) *Microfungi on land plants: an identification handbook* 2<sup>nd</sup> edn. The Richmond Publishing Co. Ltd., UK.

Engel EA, Escobar P, Montt C, Gomez-Talquenca S, Valenzuela PDT (2008) First report on the occurrence of *Grapevine leafroll-associated virus 7* and *9* in Chilean grapevines. *Plant Disease* 92: 1252.

Engel EA, Rivera PA, Valenzuela PDT (2010) First report of *Grapevine Syrah virus-1* in Chilean grapevines. *Plant Disease* 94: 63–633.

Engelbrecht M, Joubert J, Burger JT (2010) First report of Aster yellows Phytoplasma in grapevines in South Africa. *Plant Disease* 94: 373.

Eppler A, Lesan V, Lazar A (1989) Viruses and virus diseases in some vineyards in Romania. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent*. 54: 491–497.

EPPO (2002a) Alert list: *Phakopsora euvitis*. European and Mediterranean Plant Protection Organisation. [http://www.eppo.org/QUARANTINE/Alert\\_List/fungi/PHLLAM.htm](http://www.eppo.org/QUARANTINE/Alert_List/fungi/PHLLAM.htm) Accessed December 2010.

EPPO (2002b) Guideline on good plant protection practice for grapevine. *OEPP/EPPO Bulletin* 32: 371–392.

EPPO (2005) *Scirtothrips aurantii*, *Scirtothrips citri*, *Scirtothrips dorsalis*. *OEPP/EPPO Bulletin* 35: 353–356.

EPPO (2006) *Popillia japonica*. *OEPP/EPPO Bulletin* 36: 447–450.

EPPO (2009) PQR database. Paris, France: European and Mediterranean Plant Protection Organization. <http://www.eppo.org> Accessed March 2011.

EPPO (2010a) *Strawberry latent ringspot nepovirus*. EPPO A2 List of pests recommended for regulation as quarantine pests. <http://www.eppo.org/QUARANTINE/listA2.htm> Accessed September 2010.

EPPO (2010b) Data Sheets on Quarantine Pests: *Tomato ringspot nepovirus*. [http://www.eppo.org/QUARANTINE/virus/Tomato\\_ringspot\\_virus/TORSV0\\_ds.pdf](http://www.eppo.org/QUARANTINE/virus/Tomato_ringspot_virus/TORSV0_ds.pdf) Accessed January 2010.

Eskalen A, Rooney-Latham S, Gubler WD (2005) First report of perithecia of

*Phaeoacremonium viticola* on grapevine (*Vitis vinifera*) and ash tree (*Fraxinus latifolia*) in California. *Plant Disease* 89:686–686.

Essakhi S, Mugnai L, Crous PW, Groenewald JZ, Surico G (2008) Molecular and phenotypic characterization of novel *Phaeoacremonium* species associated with Petri disease and esca of grapevine. *Persoonia* 21: 119–134.

Evans AV, Hogue JN (2006) Field guide to beetles of California. California Natural History Guides, 88. University of California Press. Los Angeles, USA.

Eyres N, Wood C, Taylor A (2006) Black rot *Guignardia bidwellii*. Western Australia Department of Agriculture Factsheet Note 167.

[http://www.agric.wa.gov.au/objtwr/imported\\_assets/content/pw/ph/dis/vit/fs2006\\_blackrot\\_nyres.pdf](http://www.agric.wa.gov.au/objtwr/imported_assets/content/pw/ph/dis/vit/fs2006_blackrot_nyres.pdf) Accessed March 2012.

Ezzat YM, McConnell HS (1956) The mealybug tribe *Planococcini* (Pseudococcidae: Homoptera). *Bulletin of the Maryland Agriculture Experiment Station* A84: 1–108.

Faggioli F, Ferretti L, Pasquini G, Barba M (2002) Detection of *Strawberry latent ring spot virus* in leaves of olive trees in Italy using a one-step RT-PCR. *Journal of Phytopathology* 150: 636–639.

Faggioli F, Riccioni L, Mazzei M, Barba M (1992) Purification and characterization of a new virus found in grapevine. *Phytopathologia Mediterranea* 31: 37–40.

FAO (Food and Agricultural Organization of the United Nations) (2004) International Standards for Phytosanitary Measures (ISPM) No. 11: *Pest risk analysis for quarantine pests including analysis of environmental risks and living modified organisms*. Secretariat of the International Plant Protection Convention, Rome, Italy.

FAO (Food and Agricultural Organization of the United Nations) (2006) International Standards for Phytosanitary Measures (ISPM) No. 1: *Phytosanitary principles for the protection of plants and the application of phytosanitary measures in international trade*. Secretariat of the International Plant Protection Convention, Rome, Italy.

FAO (Food and Agricultural Organization of the United Nations) (2007) International Standards for Phytosanitary Measures (ISPM) No. 2: *Framework for pest risk analysis*. Secretariat of the International Plant Protection Convention, Rome, Italy.

FAO (Food and Agricultural Organization of the United Nations) (2009) International Standards for Phytosanitary Measures (ISPM) No. 5: *Glossary of phytosanitary terms*. Secretariat of the International Plant Protection Convention, Rome, Italy.

Farquhar D, Williams M (2000) Borers and weevils. Department of Primary Industries, Parks, Water and Environment, Tasmania, Australia.

<http://www.dpiw.tas.gov.au/inter.nsf/webpages/ljem-5se3es?open#BorersinAustralianVi> Accessed October 2011.

Farr DF, Bills GF, Chamuris GP, Rossman AY (1989) *Fungi on plants and plant products in the United States*. APS Press, St. Paul, Minnesota USA.

Farr DF, Rossman AY (2011) Fungal Database, Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.ars-grin.gov/fungalatabases/fungushost/fungushost.cfm> Accessed June 2010.

- Fasulo TR, Brooks RF (2010) Scale pests of Florida citrus. Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida ENY-814 <http://edis.ifas.ufl.edu> Accessed July 2011.
- Fayek MA, Jomaa AH, Shalaby ABA, Al-Dhaher MMA (2009) Meristem tip culture for in vitro eradication of *Grapevine leaf roll associated virus-1* (GLRaV-1) and *Grapevine fan leaf virus* (GFLV) from infected flame seedless grapevine plantlets. *Iniciacion A La Investigacion, Revista Electronica, Universidad de Jaen* 4: 1–11.
- Feil H, Feil WS, Purcell AH (2000) Effects of temperature on the flight activity of *Graphocephala atropunctata* (Hemiptera: Cicadellidae). *Journal of Economic Entomology* 93: 88–92.
- Ferreira FA, Barrreto RW, Almada RW (2005) Spots on eucalyptus wood planks caused by *Arxiomyces vitis*. *Fitopatologia Brasileira* 30: 92
- Fettig CJ (2005) *Sinoxylon perforans*. North American Forest Commission Exotic Forest Pest Information System. <http://spfnic.fs.fed.us/exfor/data/pestreports.cfm?pestidval=96&langdisplay=english> Accessed February 2008.
- Filip I (1986) The vine borer (*Sinoxylon perforans* Schrank) a new pest of grapevine plantations in Dobrudja. *Probleme de Protectia Plantelor* 14: 205–208.
- Filippin L, Jovic J, Cvrkovic T, Forte V, Clair D, Tosevski I, Boudon-Padieu E, Borgo M, Angelini E (2009) Molecular characteristics of phytoplasmas associated with Flavescence dorée in clematis and grapevine and preliminary results on the role of *Dictyophara europaea* as a vector. *Plant Pathology* 58: 826–837.
- Finlay JR, Teakle DS (1969) The effect of pH on the particle stability of a Phosphotungstate stained *Tobacco necrosis virus*. *Journal of General Virology* 5: 93–96.
- Firrao G (2004) *Candidatus* Phytoplasma, a taxon for the wall-less, non-helical prokaryotes that colonize plant phloem and insects. *International Journal of Systematic and Evolutionary Microbiology* 54: 1243–1255.
- Firrao G, Gibbs K, Streten C (2005) Short taxonomic guide to the genus ‘*Candidatus* Phytoplasma’. *Journal of Plant Pathology* 87: 249–264.
- Fischer M (2002) A new wood-decaying basidiomycete species associated with esca of grapevine: *Fomitiporia mediterranea* (Hymenochaetales). *Mycological Progress* 1: 315–324.
- Fischer M (2006) Biodiversity and geographic distribution of basidiomycetes causing esca-associated white rot in grapevine: a worldwide perspective. *Phytopathologia Mediterranea* 45: S30–S42.
- Fisher M, Binder M (2004) Species recognition, geographic distribution and host-pathogen relationships: a case study in a group of lignicolous basidiomycetes, *Phellinus*. *Mycologia* 96: 799–811.
- Flaherty DL, Wilson LT, Welter S, Lynn CD, Hanna R (1992) Spider mites. In Flaherty DL, Christensen PL, Lannini T, Marois J, Wilson LT (eds) *Grape Pest Management*. Publication 3343: 181–192. University of California Division of Agriculture and Natural Resources Publication, California.



- Fleury D, Paré J, Vincent C, Maufette Y (2006) Feeding impact of *Lygus lineolaris* (Heteroptera: Miridae) on *Vitis vinifera*: a behavioural and histological study. *Canadian Journal of Botany* 84: 493–500.
- Flint MR (2006) The UC IPM pest management guidelines: Grapevines. (Agriculture and Natural Resources. University of California. Davis).  
<http://www.ipm.ucdavis.edu/PMG/selectnewpest.grapes.html> Accessed March 2010.
- Forneck A, Huber L (2009) (A)sexual reproduction—a review of life cycles of grape phylloxera, *Daktulosphaira vitifoliae*. *Entomologia Experimentalis et Applicata* 131: 1–10.
- Forte V, Angelini E, Maixner M, Borgo M (2010) Preliminary results on population dynamics and host plants of *Hyalesthes obsoletus* in North-Eastern Italy. *Vitis* 49: 39–42.
- Fortusini A, Tolentino D, Belli G (1983) Symptomatology of herbaceous test plants of a strain of *Arabis mosaic virus* isolated from grapevine. *Informatore Fitopatologico* 33: 43–46.
- Frazier NW (1970) *Virus Diseases of Small Fruits and Grapevines*. University of California, Division of Agricultural Science, Berkeley.
- Frison EA, Ikin R (1991) *FAO/IBPGR Technical guidelines for the safe movement of grapevine germplasm*. Food and Agriculture Organisation, Rome, Italy.
- Frolov AN (2009) *Eupoecilia ambiguella* (Hubner): European grape berry moth, Grape berry moth, Grape bud moth, Vine moth, Grape moth. Interactive agricultural ecological atlas of Russia and neighbouring countries: Economic plants and their diseases, pests and weeds.  
[http://www.agroatlas.ru/pests/Eupoecilia\\_ambiguella\\_en.htm](http://www.agroatlas.ru/pests/Eupoecilia_ambiguella_en.htm) Accessed September 2010.
- Fuchs E, Gruntzig M, Auerbach I, Einecke I, Muller C, Kragenow M (1994) On the occurrence of plant pathogenic viruses in waters in the region of Halle/Saale (German Federal State of Saxony-Anhalt). *Archives of Phytopathology and Plant Protection* 29: 133–141.
- Fuchs M (2007) Grapevine Leafroll Disease. NY IPM Program factsheet, Cornell University.  
[www.nysipm.cornell.edu/factsheets/grapes/diseases/grape\\_leafroll.pdf](http://www.nysipm.cornell.edu/factsheets/grapes/diseases/grape_leafroll.pdf) Accessed September 2011.
- Fujinaga M, Ogiso H, Wakabayashi H, Morikawa T, Natsuaki T (2009) First report of a *Grapevine Algerian latent virus* disease on statice plants (*Limonium sinuatum*) in Japan. *Journal of General Plant Pathology* 75: 157–159.
- Gajardo A, Fiore N, Prodan S, Paltrinieri S, Botti S, Pino AM, Zamorano A, Montealegre J, Bertaccini A (2009) Phytoplasmas associated with Grapevine yellows disease in Chile. *Plant Disease* 93:789–796.
- Gallitelli D, Martelli GP, Di Franco A (1989) *Grapevine Algerian latent virus*, a newly recognized tombusvirus. *Phytoparasitica* 17: 61–62.
- Gallitelli D, Rana GL, Volvas C, Martelli GP (2004) Viruses of globe artichoke: An overview. *Journal of Plant Pathology* 86: 267–281.
- Gallotti GJM (1991). Resistance of *Vitis* spp. to *Fusarium oxysporum* f. sp. *herbemontis*. *Fitopatologia Brasileira* 16: 74–77.
- Galvan TL, Burkness EC, Hutchison WD (2007) Grape flea beetle.  
<http://www.vegedge.umn.edu/vegpest/grapes/FleaBeetle.htm> Accessed March 2011.

- Garran J, Gibbs A (1982) Studies on *Alfalfa mosaic virus* and alfalfa aphids. *Australian Journal of Agricultural Research* 33: 657–64.
- GBIF (2012) *Trametes ochracea*. Global Biodiversity Information Facility. <http://data.gbif.org/species/Trametes%20ochracea> Accessed March 2012.
- Geiger CA, Daane KM (2001) Seasonal movement and distribution of the grape mealybug (Homoptera: Pseudococcidae): developing a sampling program for San Joaquin Valley vineyards. *Journal of Economic Entomology* 94: 291–301.
- Giampetruzzi A, Roumi V, Roberto R, Malossini U, Yoshiwaka N, La Notte P, Terlizzi F, Credi R, Saldarelli P (2012) A new grapevine virus discovered by deep sequencing of virus- and viroid-derived small RNAs in Cv *Pinot gris* 163: 262–268.
- Gilbertson RL, Bigelow DM (1998) Annotated checklist of wood rotting Basidiomycetes of the Sky Islands in southeastern Arizona. *Journal of the Arizona–Nevada Academy of Science* 31: 13–36.
- Gillings M, Ophel-Keller K (1995) Comparison of strains of *Agrobacterium vitis* from grapevine source areas in Australia. *Australasian Plant Pathology* 24: 29–37
- Giménez-Jaime A, Aroca A, Raposo R, García-Jiménez J, Armengol J (2006) Occurrence of fungal pathogens associated with grapevine nurseries and the decline of young vines in Spain. *Journal of Phytopathology* 154: 598–602.
- Girgis SM, Bem FP, Dovas Ch, Sclavounos A, Avgelis A, Tsagris M, Katis N, Kyriakopoulou P (2009) Characterization of a novel ilarvirus causing Grapevine angular mosaic disease. *European Journal of Plant Pathology* 125: 203–211.
- Girgis SM, Bem F, Kyriakopoulou PE, Dovas CI, Sklavounos AP, Avgelis A, Katis N, Tzortzakaki S, Tsagris M (2000) A new ilarvirus isolated from grapevine in Greece. *Plant Disease* 84: 1345.
- Giri RY, Chaudhary V, Bhanja M R, Reddy SM (1996) Fungal diseases of *Eucalyptus* from Warangal-II. *Indian Forester* 122: 817–822.
- Glawe DA, Rogers JD (1984) Diatrypaceae in the Pacific Northwest. *Mycotaxon* 20: 401–460.
- Gokalp K, Digiario M, Cigsar I, Ghanem-Sabanadzovic NA, de Strandis A, Boscia D, Martelli GP (2003) Properties of a previously undescribed nepovirus from South-East Anatolia. *Journal of Plant Pathology* 85: 35–41.
- Golino DA (1993) Potential interactions between rootstocks and grapevine latent viruses. *American Journal of Enol. Viticulture* 44: 148–152.
- Golino DA, Sim S, Rowhani A (2003) The role of rootstock genotype in the effects of single and mixed infections of grapevine viruses. *Extended abstracts of the 14th Meeting of the ICVG, Locorotondo, Italy, 2003* 136–137. ICVG, Switzerland.
- Golzar H, Phillips D, Mack S (2007) Occurrence of Strawberry root and Crown rot in Western Australia. *Australasian Plant Disease Notes* 2: 145–147.
- Gonzalez RH (1968) *Biología y control de la falsa araña de la vid Brevipalpus chilensis Baker (Acaria: Phytoptipapidae)*. Boletín Técnico No 1 Experimental Agronómica,

Universidad de Chile, Santiago, Chile.

Gonzalez RH (1983) *Manejo de plagas de la vid*. Publicacions en Ciencias Agrícolas No. 13. Facultad de Ciencias Agrarias, Veterinarias y Forestales, Universidad de Chile, Santiago, Chile.

Gonzalez-Andujar JL, Ramirez-Davila JF, Lopez MA, Ocete R (2006) R Spatial Distribution of *Jacobiasca lybica* (Bergenin & Zanon) (Homoptera: Cicadellidae) Egg populations in an irrigated sherry vineyard. *Journal of Agricultural and Urban Entomology* 23: 51–55.

Gooding GV Jr, Téliv D (1970) *Grapevine yellow vein*. In Frazier NW, Fulton JP, Thresh JM, Converse RH, Varney EH, Hewitt WB (eds) *Virus diseases of small fruits and grapevines* 238–241. University of California, Berkeley.

Goodwin S (2005) Chemical control of fig longicorn, *Acalolepta vastator* (Newman) (Coleoptera: Cerambycidae), infesting grapevines. *Australian Journal of Entomology* 44: 71–76.

Graham R (2007) Black Bean Aphid (*Aphis fabae*). Pest and Diseases Image Library. <http://www.padil.gov.au> Accessed March 2010.

Grall S, Manceau C (2003) Colonization of *Vitis vinifera* by a green fluorescence protein-labelled, gfp-marked strain of *Xylophilus ampelinus*, the causal agent of bacterial necrosis of grapevine. *Applied and Environmental Microbiology* 69: 1904–1912.

Gramaje D, Agustí-Brisach C, Pérez-Sierra A, Moralejo E, Olmo D, Mostert L, Damm U, Armengol J (2012) Fungal trunk pathogens associated with wood decay of almond trees on Mallorca (Spain). *Persoonia* 28: 1–13.

Gramaje D, Alaniz S, Pérez-Sierra A, Abad-Campos P, García-Jiménez J, Armengol J (2007) First report of *Phaeoacremonium mortioniae* causing Petri disease of grapevine in Spain. *Plant Disease* 91: 1206.

Gramaje D, Armengol J (2011) Fungal trunk pathogens in the grapevine propagation process: potential inoculum sources, detection, identification, and management systems. *Plant Disease* 95: 1040–1055.

Gramaje D, Armengol J, Mohammadi H, Banihashemi Z, Mostert L (2009a) Novel *Phaeoacremonium* species associated with Petri disease and esca of grapevine in Iran and Spain. *Mycologia* 101: 920–929.

Gramaje D, Mostert L, Amengol J (2010) Characterization of *Cadophora luteo-olivacea* and *Cadophora melinii* isolates obtained from grapevine nurseries and plants in Spain. In Abstracts of oral and poster presentations given at the 7<sup>th</sup> International Workshop on Grapevine Trunk Diseases, Santa Cruz, Chile, 17–21 January 2010. *Phytopathologia Mediterranea* 49: 104.

Gramaje D, Mostert L, Armengol J (2011) Characterization of *Cadophora luteo-olivacea* and *C. melinii* isolates obtained from grapevines and environmental samples from grapevine nurseries in Spain. *Phytopathologia Mediterranea* 50 (Supplement): S112–S126.

Gramaje D, Muñoz RM, Lerma ML, García-Jiménez J, Armengol J (2009b) Fungal grapevine trunk pathogens associated with Syrah decline in Spain. *Phytopathologia Meditteranea* 48: 396–402.

- Grand LF, Vernia CS (2004) Fungi on plants in North Carolina. North Carolina State University. <http://www.cals.ncsu.edu/course/pp318/host.pdf> Accessed June 2010.
- Granett J, Walker MA, Kocsis L, Omer AD (2001) Biology and management of grape phylloxera. *Annual Review of Entomology* 46: 387–412.
- Graniti A, Surico G, Mugnai L (1994) Considerazioni sul mal dell'esca e sulle venature brune del legno della vite. *Informatore Fitopatologico* 49: 6–12.
- Grasswitz TR, James DG (2008) Movement of grape mealybug, *Pseudococcus maritimus*, on and between host plants. *Entomologia Experimentalis et Applicata* 129: 268–275.
- Grebennikov VV, Gill BD, Vigneault R (2010) *Trichoferus campestris* (Faldermann) (Coleoptera: Cerambycidae), an Asia wood-boring beetle recorded in North America. *The Coleopterists Bulletin* 64: 13–20.
- Greenwood SR (1988) Habitat stability and wing length in two species of arboreal Psocoptera. *Oikos* 52: 235–238.
- Grichanov IY, Ovsyannikova EI (2009) *Aporia crataegi* Linnaeus: Black-veined white. Interactive agricultural ecological atlas of Russia and neighbouring countries: Economic plants and their diseases, pests and weeds. [http://www.agroatlas.ru/en/content/pests/Aporia\\_crataegi/](http://www.agroatlas.ru/en/content/pests/Aporia_crataegi/) Accessed December 2010.
- Griesbach JA (1995) Detection of *Tomato ringspot virus* by polymerase chain reaction. *Plant Disease* 79: 1054–1056.
- Griffiths HM, Sinclair WA, Smart CD, Davis RE (1999) The phytoplasma associated with Ash yellows and Lilac witches'-broom: '*Candidatus* Phytoplasma fraxini'. *International Journal of Systematic Bacteriology* 49: 1605–1614.
- Griffiths M, Wylie R, Lawson S, Pegg G, McDonald J (2004) Known or potential threats from pests and diseases to prospective tree species for high value timber in northern Australia. [http://www.plantations2020.com.au/reports/pfnq/acrobat/2-4\\_griffiths\\_et\\_al.pdf](http://www.plantations2020.com.au/reports/pfnq/acrobat/2-4_griffiths_et_al.pdf) Accessed March 2012.
- Groenewald M, Kang JC, Crous PW, Gams W (2001) ITS and beta-tubulin phylogeny of *Phaeoacremonium* and *Phaeomoniella* species. *Mycological Research* 105: 651–657.
- Grové T, Steyn WP, De Beer MS (1999) The False codling moth, *Cryptophlebia leucotreta* (Meyrick) (Lepidoptera: Tortricidae) on avocado—literature review. *South African Avocado Growers' Association Yearbook* 22: 31–33.
- Guario A, Laccone G (1996) The defence of table grapes from pests. (La difesa dell'uva da tavola dai fitofagi). *Informatore Agrario Suppl.* 52: 31–40.
- Gubler D, Urbez-Torres JR, Trouillas FP, Herche R, Striegler RK, Cartright RD, Kreidty J, Rupe JC (2010) Grapevine trunk diseases: etiology, epidemiology and control. In *Proceedings of the symposium on advances in vineyard pest management, 6–8 February 2010, Missouri 7–21*. University of Missouri Extension, Missouri.
- Gugerli P, Ramel MR (1993) *Grapevine leafroll associated virus II* analyzed by monoclonal antibodies. Extended Abstracts *11th Meeting of ICVG, Montreux 1993* 23–24. ICVG, Switzerland.

- Guidoni S, Mannini F, Ferrandino A, Argamante N, di Stefano R (2000) Effect of virus status on leaf and berry phenolic compounds in two wine grapevine *Vitis vinifera* cultivars. *Acta Horticulturae* 526: 445–452.
- Gullan PJ (2000) Identification of the immature instars of mealybugs (Hemiptera: Pseudococcidae) found on citrus in Australia. *Australian Journal of Entomology* 39: 160–166.
- Gundersen DE, Lee IM, Schaff DA, Harrison NA, Chang CJ, Davis RE, Kingsbury DT (1996) Genomic diversity and differentiation among phytoplasma strains in 16S rRNA groups I (aster yellows and related phytoplasmas) and III (X-disease and related phytoplasmas). *International Journal of Systematic Bacteriology* 46: 64–75.
- Guo JY, Zhou HX, Wan FH, Liu XJ, Han ZJ (2005) Population dynamics and damage of *Lygus lucorum* Mayr in Bt cotton fields under two control measures. *Chinese Bulletin of Entomologie* 42: 4.
- Gutierrez J, Schicha E (1983) The spider mite family Tetranychidae (Acari) in New South Wales. *Acarology* 9: 107–108.
- Guy P (1982) A disease of *Atriplex* caused by *Sowbane mosaic virus*. *Australasian Plant Pathology* 11: 51–52.
- Habili N (2009) *Grapevine viruses in Australian states, 16th June, 2009 (personal communication)*. Commonwealth Scientific and Industrial Research Organisation, Adelaide.
- Habili N, Afsharifar A, Symons RH (2003) First detection of an Ampelovirus, a Maculavirus and two Vitiviruses in Iranian table grapes. *Extended abstracts 14th ICVG meeting September 12–17, 2003, Locorotondo (Bari), Italy* 162–163. ICVG, Switzerland.
- Habili N, Komínek P, Little A (2007) *Grapevine leafroll-associated virus 1* as a common grapevine pathogen. *Plant Viruses* 1: 63–68.
- Habili N, Symons RH (2000) Grapevine viruses detected by Waite diagnostics in Australia. In *13th ICVG Conference* 124–126, University of Adelaide, Adelaide.
- Halleen F, Crous PW, Petrini O (2003) Fungi associated with healthy grapevine cuttings in nurseries, with special reference to pathogens involved in the decline of young vines. *Australasian Plant Pathology* 32: 47–52.
- Halleen F, Fourie PH, Crous PW (2006a) A review of Black foot disease of grapevine. *Phytopathologia Mediterranea* 45: 55–67.
- Halleen F, Mostert L, Crous PW (2005) Pathogenicity testing of *Phialophora*, *Phialophora*-like, *Phaeoacremonium* and *Acremonium* species isolated from vascular tissues of grapevines. *Phytopathologia Mediterranea* 44: 103.
- Halleen F, Mostert L, Crous PW (2007) Pathogenicity testing of lesser-known vascular fungi of grapevines. *Australasian Plant Pathology* 36: 277–285.
- Halleen F, Schroers H-J, Groenewald JZ, Rego C, Oliveira H, Crous PW (2006b) *Neonectria liriodendri* sp. nov., the main causal agent of black foot disease of grapevines. *Studies in Mycology* 55: 227–234.
- Hamon AB (1977) Oleander Pit Scale, *Asterolecanium pustulans* (Cockerell) (Homoptera: Coccoidea: Asterolecantidae). Entomology Circular No. 184. Florida Department of

Agriculture and Consumer Service. Division of Plant Industry.

Hanken DA (2002) Importation of Grapes (*Vitis* spp.) from Korea into the United States. A qualitative, pathway-initiated pest risk assessment. Animal and Plant Health Inspection Service, United States Department of Agriculture; Riverdale, MD, USA.

Harding RB, Wicks TJ (2007) *Verticillium dahliae* and *Pratylenchus* spp: populations in potato soils and plants in Australia. *Australasian Plant Pathology* 36: 62–67.

Hardy S, Donovan N, Barkley PB (2008) *Citrus exocortis*. Primefact 772. NSW Department of Primary Industries, NSW.

Harju VA, Skelton AL, Monger WA, Jarvis B, Mumford RA (2008) Identification of an X-disease (16SrIII) group phytoplasma (*'Candidatus Phytoplasma pruni'*) infecting delphiniums in the UK. *Plant Pathology* 57: 769.

Harman JA, Mao CX, Morse JG (2007) Selection of colour of sticky trap for monitoring adult bean thrips, *Caliothrips fasciatus* (Thysanoptera: Thripidae). *Pest Management Science* 63: 210–216.

Harper SJ, Delmiglio C, Ward LI, Clover GRG (2010) Detection of *Tomato black ring virus* by real-time one-step RT-PCR. *Journal of Virological Methods* 171: 190–194.

Harris R, Ochoa-Corona R, Lebas B, Timudo O, Stewart F, Alexander B (2006) Broad detection and diagnosis of viruses of the genus Tombusvirus by RT-PCR coupled to single strand conformation polymorphism analysis. *Phytopathology* 97: S45.

Harrison BD (1957) Studies of the host range, properties and mode of transmission of *Beet ringspot virus*. *Annals of Applied Biology* 45: 462–472.

Harrison BD (1958) Relationship between *Beet ringspot*, *Potato bouquet* and *Tomato black ring viruses*. *Journal of General Microbiology* 18: 450–460.

Haviland DR, Bentley WJ, Daane KM (2005) Hot-water treatments for control of *Planococcus ficus* (Homoptera: Pseudococcidae) on dormant grape cuttings. *Journal of Economic Entomology* 98: 1109–1115

Hayova VP, Minter DW (2009) *Amerosporium concinnum*. IMI Descriptions of Fungi and Bacteria No. 1791. CAB International, Surrey, UK.

Hennen JF, Figueiredo MB, de Carvalho Jr., Hennen PG (2005) Catalogue of the species of plant rust fungi (Uredinales) in Brazil. [http://200.20.168.5/publica/livros\\_pdf/catalogue.pdf](http://200.20.168.5/publica/livros_pdf/catalogue.pdf) Accessed March 2012.

Hennessy CR, Daly AM, Hearnden MN (2007) Assessment of grapevine cultivars for resistance to *Phakopsora euvitis*. *Australasian Plant Pathology* 36: 313–317.

Herrera GM, Madariaga MV (2001) Presence and incidence of grapevine viruses in the central zone of Chile. *Agric. Téc. Chile* 61: 393–400.

Hewitt WB, Martelli G, Dias HF, Taylor RH (1970) *Grapevine fanleaf virus*. <http://www.dpvweb.net/dpv/showdpv.php?dpvno=28> Accessed March 2011.

Hill BL, Purcell AH (1997) Populations of *Xylella fastidiosa* in plants required for transmission by efficient vector. *Ecology and Population Biology* 87: 1197–1201.

- Hiratsuka N (1935) *Phakopsora* of Japan II. *Botanical Magazine, Tokyo* 49: 853–860.
- Hoddle MS (2004) The potential adventive geographic range of glassy-winged sharpshooter, *Homalodisca coagulata* and the grape pathogen *Xylella fastidiosa*: implications for California and other grape growing regions of the world. *Crop Protection* 23: 691–699.
- Hoegger PJ, Rigling D, Holdenrieder O, Heiniger U (2002) *Crytonectria radicalis*: rediscovery of a lost fungus. *Mycologia* 94: 105–115.
- Holleinová V, Bláhová L, Baránková (2009) The occurrence of viruses in the clonal selection vineyards in the Czech Republic. *Extended abstracts of the 16th meeting of ICVG in Dijon, 31 August–4 September 2009*. ICVG, Switzerland.
- Hosagoudar VB, Archana GR (2009) Host range of meliolaceous fungi in India. *Journal of Threatened Taxa* 1: 269–282.
- Hosagoudar VB, Riju MC, Agarwal DK (2010) Three new Meliolaceae members from Silent Valley National Park. *Indian Phytopathology* 63: 76–78.
- Hosagoudar VB (2003) Asteraceae of India. *Zoos Print Journal* 18: 1280–1285.
- Hossain MS, Williams DG (2003) Phenology of carpophilus beetle populations (Coleoptera: Nitidulidae, *Carpophilus spp.*) in a fruit dump in northern Victoria. *Australian Journal of Experimental Agriculture* 43: 1275–1279.
- Hovore FT (1983) Taxonomic and biological observations on southwestern Cerambycidae (Coleoptera). *The Coleopterists Bulletin* 37: 379–387.
- Hren M, Nikolić P, Rotter A, Blejec A, Terrier N, Ravnikar M, Dermastia M, Gruden K (2009) 'Bois noir' Phytoplasma induces significant reprogramming of the leaf transcriptome in the field grown grapevine. *BMC Genomics* 10: 460.
- Hu JS, Gonsalves D, Teliz (1990) Characterization of closterovirus-like particles associated with Grapevine leafroll disease. *Journal of Phytopathology* 128: 1–14.
- Hudson R, Adams D (2008) Three cornered Alfalfa hopper – *Spissistilus festinus* (Say). The Bug Wood Network. <http://www.gaipm.org/top50/3cornered.html> Accessed August 2008.
- HYPPZ (1998) Grapevine. HYPPZ Pest Encyclopaedia. French National Institute for Agricultural Research. <http://www.inra.fr/hyppz/CULTURES/6c---094.htm> Accessed May 2010.
- HYPPZ (2008) *Sparganothis pilleriana* (Denis & Schiffermüller) – Long palpi tortrix, leaf rolling tortrix. <http://www.inra.fr/hyppz/RAVAGEUR/6eupamb.htm> Accessed July 2011.
- Ipach U, Kling L, Lesemann D (2003) First record of *Cherry leaf roll virus* on grapevine in Germany. In *14th ICVG Conference Locorotondo, 12–17 September, 2003* 162–163. ICVG, Switzerland.
- IRPCM (2004) '*Candidatus Phytoplasma*', a taxon for the wall-less, non-helical prokaryotes that colonize plant phloem and insects. *International Journal of Systemic and Evolutionary Microbiology* 54: 1243–1255.
- Irvin NA, Hoddle MS (2005) Determination of *Homalodisca coagulata* (Hemiptera: Cicadellidae) egg ages suitable for oviposition by *Gonatocerus ashmeadi*, *Gonatocerus triguttatus*, and *Gonatocerus fasciatus* (Hymenoptera: Mymaridae). *Biological Control* 32:

391–400.

Irwin JAG, Jones RM (1977) The role of fungi and nematodes associated with death of white clover (*Trifolium repens*) stolons over summer in south-eastern Queensland. *Australian Journal of Experimental Research and Animal Husbandry* 17: 784–794.

Iwata R, Yamada F (1990) Notes on the biology of *Hesperophanes campestris* (Faldermann) (Col., Cerambycidae), a drywood borer in Japan. *Material und Organismen* 25: 305–313.

James DG, Whitney J (1993) Mite populations on grapevines in south-eastern Australia: implications for biological control of grapevine mites (Acarina: Tenuipalpidae, Eriophyidae). *Experimental and Applied Acarology* 17: 259–270.

Jankulova M (1972) Detection and identification of *Chenopodium mosaic virus* (Sowbane mosaic) in Bulgaria. *Phytopathologische Zeitschrift* 74: 314–317.

Jankulova M, Savino A, Gallitelli D, Quacquarelli A, Martelli GP (1978) Isolation of *Artichoke Italian latent virus* from the grapevine in Bulgaria. *Proceedings of the 6th meeting ICVG* 143–147. Cordoba, Monografias INIA No 18, Ministerio de Agricult, Madrid.

Janse JD, Obradovic A (2010) *Xylella fastidiosa*: its biology, diagnosis, control and risks. *Journal of Plant Pathology* 92:S1.35–S1.48.

Jeppson LR, Keifer HH, Baker EW (1975) *Mites injurious to economic plants*. University of California Press. Berkeley, USA.

Johannesen J, Lux B, Michel K, Seitz A, Maixner M (2008) Invasion biology and host-specificity of the grapevine yellows disease vector *Hyalesthes obsoletus* in Europe. *Entomologia Experimentalis et Applicata* 126: 217–227.

Johnson JA (2007) Survival of Indian meal moth and Navel orangeworm (Lepidoptera: Pyralidae) at low temperatures. *Journal of Economic Entomology* 100: 1482–1488.

Johnson WT, Lyon HH (1988) *Insects that feed on trees and shrubs*. Comstock Publishing Associates, Ithaca, USA.

Jonczyk M, Borodynko N, Pospieszny H (2004) Restriction analysis of genetic variability of Polish isolates of *Tomato black ring virus*. *Acta Biochimica Polonica* 51: 673–681.

Jones AL, Aldwinkle HS (1991) *Compendium of apple and pear diseases* 54–55. APS Press, USA.

Jones AT (1973) A comparison of some properties of four strains of *Cherry leaf roll virus*. *Annals of Applied Biology* 74: 211–217.

Jones AT (1985) *Cherry leaf roll virus*. CMI/ABB Descriptions of Plant Viruses No. 306. Association of Applied Biologists, Wellesbourne, UK.  
<http://www.dpvweb.net/dpv/showdpv.php?dpvno=306> Accessed September 2011.

Jones AT, Brown DJF, McGavin WJ, Rudel M, Altmayer B (1994) Properties of an unusual isolate of *German raspberry ringspot virus* from grapevine in Germany and evidence of its possible transmission by *Paralongidorus maximus*. *Annals of Applied Biology* 124: 283–300.

Jones AT, Koenig R, Lesemann DE, Hamacher J, Nienhaus F and Winter S (1990) Serological comparison of isolates of *Cherry leaf roll virus* from diseased beech and birch trees in a forest decline area in Germany with other isolates of the virus. *Journal of*



*Phytopathology* 129: 339–344.

Jones AT, Mitchell MJ, Brown DJF (1989) Infectibility of some new raspberry cultivars with *Arabis mosaic* and *Raspberry ringspot viruses* and further evidence for variation in British isolates of these two nepoviruses. *Annals of Applied Biology* 115: 57–69.

Joshi R, Gupta BP (1988) Biology of the Galerucid, *Oides scutellata* Hope (Chrysomelidae: Coleoptera) a pest of grapevine at Ghaubattia Kumaon Hills. *Prog. Hort.* 20: 144–151.

Kahrer A (2005) Introduction and possible spread of the planthopper *Metcalfa pruinosa* in Austria. Symposium Proceedings No.81. *Plant protection and plant health in Europe: introduction and spread of invasive species, held at Humboldt University, Berlin, Germany, 9–11 June 2005* 133–134. British Crop Protection Council, Alton, UK.

Kajitani Y, Kanematsu S (2000) *Diaporthe kyushuensis* sp. nov., the teleomorph of the causal fungus of grapevine swelling arm in Japan, and its anamorph *Phomopsis vitimegaspora*. *Mycoscience* 41: 111–114.

Kamp BJ Van der, Hood IA (2002) Armillaria root disease of *Pinus radiata* in New Zealand. 2: Invasion and host reaction. *New Zealand Journal of Forestry Science* 32: 103–115.

Kanzawa T (1936) Studies on *Drosophila suzukii* Mats. *Journal of Plant Protection (Tokyo)* 23: 66–70. In *Review of Applied Entomology* 24: 315 (Abstract only).

Karban R, English-Loeb, GM, Verdigaal P (1991) Vaccinating grapevines against spider mites. *California Agriculture* 45: 19–21.

Karimi M, Contaldo N, Mahmoudi B, Duduk B, Bertaccini A (2009) Identification of stolbur-related phytoplasmas in grapevine showing decline symptoms in Iran. *Extended abstracts of the 16th meeting of ICVG in Dijon, 31 August –4 September 2009*. ICVG, Switzerland.

Karimi MR, Mahmoodi B, Kazemiyan M (2001) First report of esca of grapevine in Iran. *Phytopathologia Mediterranea* 40: S481.

Kassanis B (1970) *Tobacco necrosis virus*. Descriptions of Plant Viruses. <http://www.dpvweb.net> Accessed September 2010.

Kay MK (1979) *Ctenopseustis obliquana* (Walker) (Lepidoptera: Tortricidae), Brownheaded leafroller. *Forest and Timber Insects in New Zealand No. 40*. Forest Research Institute, Rotorua, New Zealand.

Kearns CG, Mossop DW (1984) Detection of nepoviruses of *Vitis vinifera* in New Zealand using enzyme-linked immunosorbent assay (ELISA). *New Zealand Journal of Agricultural Research* 27: 431–435.

Kegler H, Kontzog HG (1990) Non-vectored transmission of plant viruses. Deutschen Phytomedizinischen Gesellschaft (DPG), Braunschweig, Germany, *Proceedings of the First Symposium of the International Working Group on Plant Viruses with Fungal Vectors, Braunschweig, Germany 21–24 August 1990* 159–162. Institute of Phytopathology, Aschersleben, Germany.

Kharizanov A (1969) A new pest of grapevine in Bulgaria. *Rastitelna Zashtita* 17: 21–23.

Kheder AA, Ibrahim IAM and Mazyad HM (2004) Isolation and characterization of *Peach Rosette Mosaic Virus*. *Egyptian Journal of Virology* 1: 259–272

- Kiewnick L (1989) A new discovery of *Ascochyta ampelina* Sacc. on leaves of *Vitis vinifera* L. *Nachrichtenblatt des Deutschen Pflanzenschutzdienstes* 41: 17–19.
- Kim KC, Lee TS (1985) Identification, larval host plant range, and damage of the fruit sucking moths to the major fruit in Cheonnam Province. *Korean Journal of Plant Protection* 24: 183–190.
- Kirchmair M, Neuhauser S, Huber L (2005) *Sorosphaera viticola* spec. nov. (Plasmodiophorids), an intracellular parasite in roots of grape vine. *Sydowia* 57: 223–232.
- Kirk P (2012) Index Fungorum. <http://www.indexfungorum.org/Index.htm> Accessed February 2012.
- Kirk P, Cooper J (2009) The Fungal Records Database of Britain and Ireland (FRDBI) <http://www.fieldmycology.net/Index.htm> Accessed January 2011.
- Klein Koch C, Waterhouse DF (2000) Distribution and importance of arthropods associated with agriculture and forestry in Chile (Distribucion e importancia de los artropodos asociados a la agricultura y silvicultura en Chile). ACIAR Monograph No. 68. Canberra. Australia. 231 pp.
- Koenig R, Feudal M, Lesemann DE (1989) Detection of *Petunia asteroid mosaic*, *Carnation ringspot* and *Tobacco necrosis viruses* in ditches and drainage canals in a grapevine-growing area in West Germany. *Journal of Phytopathology* 127: 169–172.
- Koklu G, Digiario M, Sabanadzovic S, Savino V (1999) Natural infections by *Cucumber mosaic virus* in Turkish grapevines. *Phytopathologia Mediterranea* 38: 33–36.
- Koklu G, Digiario M, Savino V (1998) A survey of grapevine viruses in Turkish Thrace. *Phytopathologia Mediterranea* 37: 140–142.
- Koltunow AM, Krake LR, Johnson SD, Rezaian MA (1989) Two related viroids cause Grapevine yellow speckle disease independently. *Journal of General Virology* 70: 3411–3419.
- Koltunow AM, Krake LR, Rezaian MA (1988) Hop stunt viroid in Australian grapevine cultivars: potential for hop infection. *Australasian Plant Pathology* 17: 7–10.
- Komar V, Vigne E, Demangeat G, Fuchs M (2007) Beneficial effect of selective virus elimination on the performance of *Vitis vinifera* cv. Chardonnay. *American Journal of Enology and Viticulture* 58: 202–210.
- Komínek P (2008) Distribution of grapevine viruses in vineyards of the Czech Republic. *Journal of Plant Pathology* 90: 357–358.
- König H, Uden G, Fröhlich J (2009) Biology of microorganisms on grapes, in must and in wine. Springer-Verlag, Berlin. Heidelberg.  
<http://books.google.com.au/books?id=6CR1vQUMw-sC&printsec=frontcover#v=onepage&q&f=false> Accessed November 2011.
- Kovacs LG, Hanami H, Fortenberry M, Kaps ML (2000) Latent infection by phloem-limited viruses is linked to lower fruit quality in French-American hybrid grapevines. In *Extended abstracts of the 13th Meeting of the ICVG, Adelaide, Australia* 158–159. ICVG, Switzerland.
- Kovacs LG, Hanami H, Fortenberry M, Kaps ML (2001) Latent infection by leafroll agent

- GLRaV-3 is linked to lower fruit quality in French-American hybrid grapevines Vidal blanc and St. Vincent. *American Journal of Enology and Viticulture* 52: 254–259.
- Krake LR, Steel-Scott N, Rezaian MA, Taylor RH (1999) *Graft Transmitted Diseases of Grapevines*. CSIRO Publishing, Collingwood, Victoria.
- Kulkarni NS, Mani M, Banerjee K (2007) *Management of thrips on grapes*. Extension Folder No. 13. National Research Centre for Grapes, Pune.
- Kumar K, Lal SN (1983) Studies on the biology, seasonal abundance and host-parasite relationship of fruit sucking moth *Othreis fullonia* (Clerck) in Fiji. *Fiji Agricultural Journal* 45: 71–77.
- Kummuang N, Diehl SV, Smith BJ, Graves CH Jr (1996) Muscadine grape berry rot diseases in Mississippi: disease epidemiology and crop reduction. *Plant Disease* 80: 244–247.
- Kuniyuki H, Gioria R, Rezende JAM, Willink CG de, Novo JPS, Yuki VA (2006) Transmission of the *Grapevine virus* by the mealybug *Pseudococcus longispinus* Targioni-Tozzetti (Hemiptera: Pseudococcidae) in Brazil. *Summa Phytopathol.* 32: 151–155.
- Kuniyuki H, Rezende JAM, Gaspar JO, Yuki, VA (2008) Detection of *Grapevine leafroll-associated virus 5* in the State of Sao Paulo, Brazil. *Summa Phytopathologica* 34: 366–367.
- Kunugi Y, Asari S, Terai Y, Shinkai A (2000) Studies on the Grapevine berry inner necrosis virus disease, 2: transmission of *Grapevine berry inner necrosis virus* by the grape erineum mite *Colomerus vitis* in Yamanashi. *Bulletin of Yamanashi Fruit Tree Experimental Station* 10: 57–63.
- Kuo M (2003) *Hapalopilus nidulans*. MushroomExpert.Com: [http://www.mushroomexpert.com/hapalopilus\\_nidulans.html](http://www.mushroomexpert.com/hapalopilus_nidulans.html) Accessed March 2011.
- Kyriakopoulou PE (2008) *Artichoke Italian latent virus* causes Artichoke patchy chlorotic stunting disease. *Annals of Applied Biology* 127: 487–497.
- Laimer Da Câmara Machado M, Paltrinieri S, Hanzer V, Arthofer W, Strommer S, Martini M, Bertaccini (2001) Presence of European stone fruit yellows (ESFY or 16SrX-B) Phytoplasmas in apricots in Austria. *Plant Pathology* 50: 130–135.
- Laimer M, Lemaire O, Herrbach E, Goldschmidt V, Minafra A, Bianco P, Wetzl T (2009) Resistance to virus, phytoplasmas and their vectors in the grapevine in Europe: a review. *Journal of Plant Pathology* 91: 7–23.
- Langer M, Maixner M (2004) Molecular characterisation of Grapevine yellows associated phytoplasmas of the Stolbur group based on RFLP-analysis of non-ribosomal DNA. *Vitis* 43: 191–200.
- Larignon P, Dubos B (1997) Fungi associated with esca disease in grapevine. *European Journal of Plant Pathology* 103: 147–157.
- Larsson MC, Leal WS, Hansson BS (2001) Olfactory receptor neurons detecting plant odours and male volatiles in *Anomala cuprea* beetles (Coleoptera: Scarabaeidae). *Journal of Insect Physiology* 47: 1065–1076.
- Lauber HP, Schuepp H (1968) Grape diseases. *Schweizerische Zeitschrift für Obst* 104: 550–555.

- Lazar J, Kolber M, Lehoczy J (1990) Detection of some nepoviruses (GFV, GFVYM, GCMV, ArMV) in the seeds and seedlings of grapevine by ELISA. *Kertgasdasag* 22: 58–72.
- Le Gall O, Candresse T, Dunez J (1995) Transfer of the 31 non-translated region of *Grapevine chrome mosaic virus* RNA-1 by recombination to *Tomato black ring virus* RNA-2 in pseudorecombinant isolates. *Journal of General Virology* 76: 1285–1289.
- Lee IM, Bertaccini A, Vibio M and Gundersen DE (1995) Detection of multiple phytoplasmas in perennial fruit trees with decline symptoms in Italy. *Phytopathology* 85: 728–735.
- Lee IM, Davis RE, Gundersen-Rindal DE (2000) Phytoplasma: phytopathogenic mollicutes. *Annual Review of Microbiology* 54: 221–255.
- Lee IM, Gundersen-Rindal DE, Davis RE, Bottner KD, Marcone C, Seemüller E (2004a) ‘*Candidatus* Phytoplasma asteris’, a novel phytoplasma taxon associated with Aster yellows and related diseases. *International Journal of Systemic and Evolutionary Microbiology* 54: 1037–1048.
- Lee IM, Martini M, Marcone C, Zhu SF (2004b) Classification of phytoplasma strains in the Elm yellows group (16SrV) and proposal of ‘*Candidatus* Phytoplasma ulmi’ for the phytoplasma associated with Elm yellows. *International Journal of Systematic and Evolutionary Microbiology* 54: 337–347.
- Lee S, Lee GS, Goh HG (2002) Mirid bugs (Heteroptera: Miridae) on grapevine: their damages and host plants. *Korean Journal of Applied Entomology* 41: 33–41.
- Lee SC, Yoo JK, Yoo CH (1970) Survey on the kinds of the fruit sucking moths and their damages in Korea. *Korean Journal of Plant Pathology* 9: 37–41.
- Lee YJ, Lee YM, Hseuh T (1973) Observations on the behaviour of *Proagopertha lucidula* Fald. (Coleoptera: Scarabaeidae). *Acta Entomologica Sinica* 16: 25–31.
- Legorburu FJ, Recio E, López E, Baigorri J, Larreina M, Remesal A, Cibrián JF, Caminero L, Suberviola J, Aguirrezábal F (2009) Effect of *Grapevine fanleaf virus* (GFLV) and *Grapevine leafroll-associated virus 3* (GLRaV-3) on red wine quality. *Extended abstracts of the 16th meeting of ICVG in Dijon, 31 August–4 September 2009*. ICVG, Switzerland.
- Lehoczy J (1991) Remarkable differences in pathogenicity between *Grapevine chrome mosaic* and *Tomato black ring viruses* in herbaceous test plants. *Kertgasdasag* 23: 49–54.
- Lehoczy J, Martelli GP, Lazar J (1992) Seed transmission of *Grapevine line pattern virus*. *Phytopathologia Mediterranea* 31: 115–116.
- Lehoczy J, Tasnády G (1971) The effects of Fanleaf and Chrome mosaic virus diseases on the yield and fruit sugar content of grapevines. *Kiserletugyi Közlemények* 64: 49–64. (Abstract only).
- Leong SL, Hocking AD, Scott ES (2006) *Aspergillus* species producing ochratoxin A: isolation from vineyard soils and infection of Semillon bunches in Australia. *Journal of Applied Microbiology* 102: 124–133.
- Lessio F, Alma A (2004a) Dispersal patterns and chromatic response of *Scaphoideus titanus* Ball (Homoptera Cicadellidae), vector of the phytoplasma agent of Grapevine flavescence dorée. *Agricultural and Forest Entomology* 6: 121–127.

- Lessio F, Alma A (2004b) Seasonal and Daily Movement of *Scaphoideus titanus* Ball (Homoptera: Cicadellidae). *Environmental Entomology* 33: 1689–1694.
- Leu LS (1988) Rust. In Pearson RC, Goheen AC (eds) *Compendium of grape diseases* 28–30. The American Phytopathological Society Press, St. Paul.
- Lew H, Chelkowski J, Pronczuk P, Edinger W (1996) Occurrence of the mycotoxin moniliformin in maize [*Zea mays* L.] ears infected by *Fusarium subglutinans* (Wollenw. & Reinking) Nelson *et al.* *Food Additives and Contaminants* 13: 321–324.
- Li ZX (2004) *Control of pests and diseases of table grape*. 2<sup>nd</sup> Edition. Jindun Press, Beijing, China.
- Liberato JR, Daly AM, Shivas RG (2007) Grapevine leaf rust (*Phakopsora euvitidis*). Pest and Diseases Image Library. [www.padil.gov.au](http://www.padil.gov.au) Accessed September 2010.
- Liburd O, Seferina G, Weihma S (2004) Insect pests of grapes in Florida. Entomology and Nematology Department, Institute of Food and Agricultural Sciences (IFAS), University of Florida. <http://edis.ifas.ufl.edu/pdf/IN/IN52700.pdf> Accessed February 2010.
- Lingafelter SW, Hoebeke ER (2002) *Revision of Anoplophora*. Entomological Society of Washington, Washington, DC.
- Liu S, Cheng ZJ, Zhang CX (1996) Study on occurrence and integrated control of *Alternaria viticola*. In Qiu, S (ed.) Progress of research on plant protection in China. *Proceedings of the third national conference of integrated pest management, Beijing, China, 12–15 November 1996* 341–345. China Science and Technology Press, Beijing.
- Liu Y, Liu DY, Cao BQ, Zhang ZY (2004) The occurrence of green plant bug in grape yard and its control. *China Fruits* 6: 40.
- Loch A, Slack J (2007) Grape phylloxera: the world's worst grapevine pest. *Primefacts: Profitable and Sustainable Primary Industries* 553: 1–4.
- Lorenzo LE, Messuti MI (2009) Additions and update to the knowledge of the genus *Hysterographium* (Ascomycota, Hysteriaceae) in southern South America. *Darwiniana* 47: 289–296.
- Lovisolo O (1990) Ecological observations on *Petunia asteroid mosaic virus* (PeAMV). Deutschen Phytomedizinischen Gesellschaft (DPG), Braunschweig, Germany. *Proceedings of the First Symposium of the International Working Group on Plant Viruses with Fungal Vectors, Braunschweig, Germany 21–24 August 1990* 155–158. Institute of Phytopathology, Aschersleben, Germany.
- Lucchi A, Santini L (1993) Note morfo-biologiche sugli stadi preimmaginali di *Metcalfa pruinosa* (Say) (Homoptera, Flatidae). *Frustula Entomologica* 16: 175–185.
- Luck J, Mann R, van Rijswijk B, Moran J, Merriman P (2012) National diagnostic protocol for Pierce's disease, *Xylella fastidiosa*. <http://www.padil.gov.au/Media/Sphds/NDP%206%20-%20Pierce's%20disease%20issued%202010%20V1.1.pdf> Accessed May 2012.
- Luo L, Wen Y, Wu Z (2005) *Description and damage review on 17 pests of grape wine borers in South Guizhou*. South Guizhou Province State Plant Protection Station, China.
- Lupo S, Bettucci L, Perez A, Martinez S, Cesari C, Escoriaza G, Gatica M (2006)

- Characterization and identification of the basidiomycetous fungus associated with ‘hoja de malvón’ grapevine disease in Argentina. *Phytopathologia Mediterranea* 45: 110–116.
- Luque J, Pera J, Parladé J (2008) Evaluation of fungicides for the control of *Botryosphaeria corticola* on cork oak in Catalonia (NE Spain). *Forest Pathology* 38: 147–155.
- Ma CS (2006) Pests, diseases and weeds of apples published by Key Laboratory of Agriculture Department, China. Key Laboratory of Agriculture Department.  
[http://www.ecolsafety.org.cn/db/Fruit/login\\_pg.asp](http://www.ecolsafety.org.cn/db/Fruit/login_pg.asp) Accessed September 2011.
- Ma J, Zhu X, Zhao L, Mei Y, Patiguli A, Zhu R, Wang Y (2004) Preliminary study on Spike-stalk brown spot of grape. *Xinjiang Academy of Agricultural Sciences, Urumchi, China* 41: 353–354.
- MacGill EI (1932) The biology of *Erythroneura* (*Zygina*) *pallidifrons*, Edwards. *Bulletin of Entomological Research* 23: 33–43.
- Machowicz-Stefaniak Z, Zalewska E (2002) Hazel diseases and their fight. Plant press.  
<http://www.ho.haslo.pl/article.php?id=1119> Accessed May 2011.
- Mackenzie A (1996) A trade-off for host plant utilization in the black bean aphid, *Aphis fabae*. *Evolution* 50: 155–162.
- Mackenzie A, Dixon AFG (1990) Host alternation in aphids: constraint versus optimization. *The American Naturalist* 136: 132–134.
- MacKenzie DJ (1997) *A standard protocol for the detection of viruses and viroids using a reverse transcription-polymerase chain reaction technique*. Document CPHBT-RTPCR1.00. Canadian Food Inspection Agency, Canada.
- MacKenzie DL, Johnson RC, Warner C (1996) Incidence of four important viral pathogens in Canadian vineyards. *Plant Disease* 80: 955–958.
- Mackie A (2005) Brown rot *Monilinia fructigena*. Western Australia Department of Agriculture and Food Factsheet. Note 181.  
[http://www.agric.wa.gov.au/objtwr/imported\\_assets/content/pw/ph/dis/fn/fs2006\\_brownrot\\_a\\_mackie.pdf](http://www.agric.wa.gov.au/objtwr/imported_assets/content/pw/ph/dis/fn/fs2006_brownrot_a_mackie.pdf) Accessed March 2012.
- MacLeod A (2006) CSL Pest Risk Analysis for *Planococcus lilacinus*.  
[fera.defra.gov.uk/plants/plantHealth/pestsDiseases/documents/plano.pdf](http://fera.defra.gov.uk/plants/plantHealth/pestsDiseases/documents/plano.pdf) Accessed July 2011.
- Madge D (2007) *Organic viticulture: an Australian manual*. Victorian Department of Primary Industries, Victoria.
- MAF (2009) *Importation of Nursery Stock*. MAF Biosecurity New Zealand 155.02.06. Ministry of Agriculture and Forestry. Wellington, New Zealand.
- Magarey PA, Gadoury DM, Emmett RW, Biggins LT, Clarke K, Wachtel MF, Wicks TJ, Seem RC (1997) Cleistothecia of *Uncinula necator* in Australia. *Proceedings 2nd International Workshop on Grapevine Downy and Powdery Mildew Modelling, Freiburg 1997* 122. Fraund, Germany.
- Magarey RD, Ernme RW, Magareye PA, Franz PR (1993) Evaluation of control of grapevine anthracnose caused by *Elsinoe ampelina* by pre-infection fungicides. *Australasian Plant Pathology* 22: 48–52.

- Mahfoudhi N, Digiario M, Dhouibi MH (2009) Transmission of *Grapevine leafroll viruses* by *Planococcus ficus* (Hemiptera: Pseudococcidae) and *Ceroplastes rusci* (Hemiptera: Coccidae). *Plant Disease* 93: 999–1002.
- Mahfoudhi N, Digiario M, Savino V, Di Terlizzi B (1998) Viruses and virus diseases of grapevine in Tunisia. *EPPO Bulletin* 28: 197–204.
- Maixner M (2011) Recent advances in Bois noir research. *2nd European Bois noir Workshop, 27 February–1 March 2011, Castelbrando, Cison di Valmarino (TV), Italy* 17–32. CRA, Padova, Italy.
- Maixner M, Pearson RC (1992) Studies on *Scaphoideus titanus*, a possible vector of grapevine yellows on wild and cultivated grapes in New York. *Proceedings of the 10th Meeting of ICVG*, Volos 1990.
- Male MF, Vawdrey LL (2010) Efficacy of fungicides against damping-off in papaya seedlings caused by *Pythium aphanidermatum*. *Australasian Plant Disease Notes* 5: 103–104.
- Maliogka VI, Dovas CI, Katis NI (2008a) Evolutionary relationships of virus species belonging to a distinct lineage within the Ampelovirus genus. *Virus Research* 135: 125–135.
- Maliogka VI, Dovas CI, Katis NI (2008b) Generic and species-specific detection of viruses belonging to a distinct lineage within the Ampelovirus genus. *Journal of Virological methods* 154: 41–47.
- Malipatil MB, Medhurst AK, Bates VI, Williams DG (1996) *Pests of Pome and Stone Fruit and their Predators and Parasitoids: a pocket guide*. Agriculture Victoria, Australia.
- Manceau C, Grall S, Brin C, Guillaumes J (2005) Bacterial extraction from grapevine and detection of *Xylophilus ampelinus* by a PCR and Microwell plate detection system. *Bulletin OEPP/EPPO Bulletin* 35: 55–60.
- Mannini F, Credi R (2000) Appraisal of agronomic and enological modifications in the performances of grapevine clones after virus eradication. *Extended abstracts of the 13th Meeting of the ICVG, Adelaide* 151–154. ICVG, Switzerland.
- Marais E (1997) Phloem-feeding by the leafhopper *Acia lineatifrons* (Naude) (Homoptera: Cicadellidae) on grapevines (*Vitis vinifera* L.) and wild blackberry, *Rubus chrysocarpus* (Smith). *African Entomology* 5: 177–180.
- Marco S Di, Calzarano F, Osti F, Mazzullo A (2004) Pathogenicity of fungi associated with a decay of kiwifruit. *Australasian Plant Pathology* 33: 337–342.
- Margaria P, Rosa C, Marzachi C, Turina M, Palmano S (2007) Detection of Flavescence dorée Phytoplasma in grapevine by reverse-transcription PCR. *Plant Disease* 91: 1496–1501.
- Martelli GP (1978) Nematode-borne viruses of grapevine, their epidemiology and control. *Nematologia Mediterranea* 6: 127.
- Martelli GP (1993) Graft-transmissible diseases of grapevines. Handbook for detection and diagnosis. Food and Agriculture Organization of the United Nations, Viale delle Terme di Caracalla, 00100 Rome, Italy. <http://www.fao.org/docrep/t0675e/T0675E00.htm#Contents> Accessed September 2011.
- Martelli GP (2009) Grapevine virology highlights 2006–2009. *Extended abstracts of the 16th*

- meeting of ICVG in Dijon, 31 August–4 September 2009 15–24. ICVG, Switzerland.
- Martelli GP (2012) Grapevine virology highlights: 2010–2012. *Proceedings of the 17<sup>th</sup> Congress of ICVG, Davis, California, USA, October 7–14 2012* 13–31. Foundation Plant Services, University of California, Davis.
- Martelli GP, Adams MJ, Kreuze JF, Dolja VV (2007) Family Flexiviridae: A case study in virion and genome plasticity. *Annual Review of Phytopathology* 45: 73–100.
- Martelli GP, Boudon-Padieu E (2006) Directory of infectious diseases of grapevines. *Options Médit. Ser. B, Studies and Research* 55: 7–201.
- Martelli GP, Gallitelli D, Abracheva P, Savino V, Quaquarelli A (1977) Some properties of grapevine *Bulgarian latent virus*. *Annals of Applied Biology* 85: 51–58.
- Martelli GP, Lehoczky J, Quacquarelli A (1970) *Hungarian chrome mosaic*. In Frazier NW, Fulton JP, Thresh JM, Converse RH, Verney EH, Hewitt WB (eds) *Virus diseases of small fruits and grapevines* 236–237. University of California, Berkeley.
- Martelli GP, Quacquarelli A, Rana GL, Gallitelli D (1978) Description of plant viruses: *Grapevine Bulgarian latent virus*. <http://www.dpvweb.net/dpv/showdpv.php?dpvno=186> Accessed September 2011.
- Martelli GP, Walter B (1993) True virus diseases: Grapevine degeneration – European nepoviruses. In *Graft transmissible diseases of grapevine: handbook for detection and diagnosis*. International council for the study of viruses and virus diseases of the grapevine. Food and Agriculture Organisation of the United States, Rome.
- Martin MT, Martin L, Cuesta MJ (2011a) First report of *Cylindrocarpon pauciseptatum* associated with grapevine decline from Castilla y León, Spain. *Plant Disease* 95: 361.
- Martin MT, Martin L, Cuesta MJ (2011b) First Report of *Neofusicoccum mediterraneum* and *N. australe* causing decay in *Vitis vinifera* in Castilla y León Spain. *Plant Disease* 95: 876.
- Martin RR, Tzanetakis IE (2006) Characterization and recent advances in the detection of strawberry viruses. *Plant Disease* 90: 384–396.
- Martini M, Botti S, Marcone C, Marzachi` C, Casati P, Bianco PA, Benedetti R, Bertaccini A (2002) Genetic variability among Flavescence dorée Phytoplasmas from different origins in Italy and France. *Molecular Cell Probes* 16: 197–208.
- Martinson T, Fuchs M, Loeb G, Hoch H (2008) *Grapevine leafroll* – an increasing problem in the Finger Lakes, the US and the world. Cornell University Cooperative Extension [http://winegrape.ag.psu.edu/docs/winegrapeinfo/attach/Grapevine\\_Leafroll\\_Cornell\\_Report.pdf](http://winegrape.ag.psu.edu/docs/winegrapeinfo/attach/Grapevine_Leafroll_Cornell_Report.pdf) Accessed September 2011.
- Martinson TE, Dunst R, Lakso A, English-Loeb G (1997) Impact of feeding injury by Eastern grape leafhopper (Homoptera: Cicadellidae) on yield and juice quality of Concord grapes. *American Journal of Enology and Viticulture* 48: 291–302.
- Marullo R (2009) Host-plant ranges and pest potential: habits of some thrips species in areas of southern Italy. *Bulletin of Insectology* 62: 253–255.
- Masri S, Rast H, Johnson R, Monette P (2006) *Grapevine virus C* and *Grapevine leafroll associated virus 2* are serologically related and appear to be the same virus. *Extended*



- abstracts of the 15th Meeting of the ICVG, Stellenbosch, South Africa, 3–7 April 2006 49. ICVG, Switzerland.
- Mathew G (1987) Cossid pests of plantation crops in India and the prospects of their management. *Journal of Coffee Research* 17: 137–140.
- Matsubara Y-I, Kayukawa Y, Yano M, Fukui H (2000) Tolerance of asparagus seedlings infected with arbuscular mycorrhizal fungus to violet root rot caused by *Helicobasidium mompa*. *Journal of the Japanese Society of Horticultural Science* 69: 552–556.
- Matus JT, Vega A, Loyola R, Serrano C, Cabrera S, Arce-Johnson P (2008) Phytoplasma and virus detection in commercial plantings of *Vitis vinífera* cv. Merlot exhibiting premature berry dehydration. *Electronic Journal of Biotechnology* 11: 1–10.
- Mau RFL, Mitchell WC (1978) Development and reproduction of the oriental stink bug, *Plautia stali* (Hemiptera: Pentatomidae). *Annals of the Entomological Society of America* 71: 756–757.
- May TW, Milne J, Shingles S (2003) *Catalogue and bibliography of Australian Fungi 2. Basidiomycota p.p. and Myxomycota p.p. Fungi of Australia*. Volume 2B. ABRS/CSIRO Publishing, Melbourne.
- Mazzoni V, Anfora G, Ioratti C, Lucchi A (2008) Role of the winter host plants in vineyard colonization and phenology of *Zygina rhamni* (Hemiptera: Cicadellidae: Typhlocibinae). *Annals of Entomological Society of America* 101: 1003–1009.
- Mazzoni V, Cosci F, Lucchi A, Santini L (2001) Leafhoppers and planthoppers vectors in Ligurian and Tuscan vineyards. In Lozzia C (ed) Integrated Control in Viticulture Working Group Proceedings, Ponte de Lima, Portugal, 3–7 March 2000. *IOBC wprs Bulletin* 24: 263–266.
- McGee PA (1989) Vesicular-arbuscular mycorrhizal and saprophytic fungi of the Swain Reefs, Australia. *Mycological Research* 93: 375–378.
- McGregor G, Whattam M, Washington B (1996) Raspberries and cultivated blackberries: pests and diseases. AG0570. <http://www.dpi.vic.gov.au/agriculture/horticulture/fruit-nuts/berries/raspberries-and-cultivated-blackberries-pests-and-diseases> Accessed October 2012.
- McKenzie EHC, Foggo MN (1989) Fungi of New Zealand subantarctic islands. *New Zealand Journal of Botany* 27: 91–100.
- McLaren GF, Grandison G, Wood GA, Tate G, Horner I (1999) *Summerfruit in New Zealand*. University of Otago Press, New Zealand.
- McLeod MJ, Williams RN (1990) Life history and vineyard damage by rose chafer, *Macrodactylus subspinosus* (F.). *Vinifera Wine Growers Journal* 17: 25–27.
- McPherson JE, McPherson RM (2000) *Stink bugs of economic importance in America North of Mexico*. CRC Press, USA.
- Mead FW, Webb SE (2001) Grape Leaf folder. Florida Department of Agriculture and Consumer Services, University of Florida. Publication Number: EENY-192. [http://creatures.ifas.ufl.edu/fruit/grape\\_leaffolder.htm](http://creatures.ifas.ufl.edu/fruit/grape_leaffolder.htm) Accessed April 2010.

- Meagher JW, Brown RH, Taylor RH, Harris AR (1976) The distribution of *Xiphinema index* and other parasitic nematodes associated with grapevines in north-eastern Victoria. *Australian Journal of Experimental Agriculture and Animal Husbandry* 16: 932–936.
- Meena B (2010) Effect of foliar application of *Pseudomonas fluorescence* on the activity of lytic enzymes in response to *Cercosporidium personatum* in groundnut. *International Journal of Plant Protection* 3: 279–281.
- Meulewaeter F, Seurinck J, van Emmelo J (1990) Genome structure of *Tobacco necrosis virus strain A*. *Virology* 177: 699–709.
- Mhirsi S, Achecheb S, Fattoucha S, Boccardo G, Marrakchib M, Marzoukia N (2004) First report of phytoplasma in the Aster yellow group infecting grapevines in Tunisia. *Plant Pathology* 53: 521.
- Miller AN, Huhndorf SM (2009) Pyrenomycetes of the World. University of Illinois. <http://www-s.life.illinois.edu/pyrenos> Accessed November 2011.
- Minafra A, Hadidi A (1994) Sensitive detection of *Grapevine virus A, B, or Leafroll-associated III* from viruliferous mealybugs and infected tissue by cDNA amplification. *Journal of Virological Methods* 47: 175–187.
- Minfra A, Grieco, F, Gallitelli D, Martelli GP (1994) Improved PCR procedures for multiple identification of some artichoke and grapevine viruses. *Bulletin OEPP/EPPO Bulletin* 25: 283–287.
- Mizell RF, Andersen PC, Tipping C, Brodbeck B (2008) *Xylella fastidiosa* diseases and their leafhopper vectors. ENY-683 (IN174), one of a series of the Department of Entomology and Nematology, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. <http://edis.ifas.ufl.edu> Accessed March 2011.
- Mohanan C (2005) Diseases of rattan in nurseries, plantations and natural stands in Kerala, India. *Journal of Bamboo and Rattan* 4: 151–162.
- Moleas T (1988) Epidemiological observations on *Sinoxylon sexdentatum* Oliv. and *Amphicerus* (= *Schistoceros*) *bimaculatus* Oliv. (Coleoptera: Botrychidae) on grape in Apulia (Italy). *Informatore Fitopatologico* 38: 55–58.
- Monthey R, Cross R (2000) Some edible fungi growing on trees in Northeast Woodlots. USDA Forest Service, State and Private Forestry Northeastern Area. <http://na.fs.fed.us/stewardship/pubs/ediblefungi/ediblefungi.htm> Accessed December 2011.
- Morales CF (1991) Margarodidae (Insecta: Hemiptera). *Fauna of New Zealand* 21. Manaaki Whenua Press, New Zealand.
- Moran J (1994) Virus diseases of carnations. Agriculture Note AG 0174. Department of Primary Industries. <http://www.dpi.vic.gov.au/agriculture/pests-diseases-and-weeds/plant-diseases/flowers/virus-diseases-of-carnations> Accessed September 2011.
- Moreau J, Villemant C, Benrey B, Thiéry D (2010) Species diversity of larval parasitoids of the European grapevine moth (*Lobesia botrana*, Lepidoptera: Tortricidae): the influence of region and cultivar. *Biological Control* 54: 300–306.
- Mori N, Pavan F, Bondavalli R, Reggiani N, Paltrinieri S, Bertaccini A (2007) Factors affecting the spread of ‘Bois noir’ disease in north Italy vineyards. *Vitis* 47: 65–72.

- Mostert L, Groenewald JZ, Summerbell RC, Gams W, Crous PW (2006a) Taxonomy and pathology of *Togninia* (Diaporthales) and its *Phaeoacremonium* anamorphs. *Studies in Mycology* 54: 1–115.
- Mostert L, Groenewald JZ, Summerbell RC, Robert V, Sutton DA, Padhye AA, Crous PW (2005) Species of *Phaeoacremonium* associated with infections in humans and environmental reservoirs in infected woody plants. *Journal of Clinical Microbiology* 43: 1752–1767.
- Mostert L, Halleen F, Fourie P, Crous PW (2006b) A review of *Phaeoacremonium* species involved in Petri disease and esca of grapevine. *Phytopathologia Mediterranea* 45: S12–S29.
- Mound L (2008) Bean thrips (*Caliothrips fasciatus*) Pest and Diseases Image Library. <http://www.padil.gov.au> Accessed March 2010.
- Mugnai L, Graniti A, Surico G (1999) Esca (black measles) and brown wood-streaking: two old and elusive diseases of grapevine. *Plant Disease* 83: 404–418.
- Murant AF (1983) Seed and pollen transmission of nematode borne viruses. *Seed Science and Technology* 11: 973–979
- Nakaune R, Toda S, Mochizuki M, Nakano M (2008) Identification and characterization of a new vitivirus from grapevine. *Archives of Virology* 153: 1827–1832.
- Namba S, Boscia D, Azzam O, Maixner M, Hu JS, Golino D, Gonsales D (1991a) Purification and properties of closterovirus like particles associated with grapevine corky disease. *Phytopathology* 81: 964–970.
- Namba S, Boscia D, Yamashita S, Tsuchizaki T, Gonzalves D (1991b) Purification and properties of spherical virus particles associated with grapevine ajinashika disease. *Plant Disease* 75: 1249–1253.
- Namba S, Yamashita S, Doi Y, Yora K (1979) A small spherical virus associated with the ajinashika disease of Koshu grapevine. *Annals of the Phytopathological Society of Japan* 45: 70–73.
- Nan ZB (1995) Fungicide seed treatments of sainfoin control seed-borne and root-invading fungi. *New Zealand Journal of Agricultural Research* 38: 413–420.
- Narai Y, Murai T (2002) Individual rearing of the Japanese mealybug, *Planococcus kraunhiae* (Kuwana) (Homoptera: Pseudococcidae) on germinated broad bean seeds. *Applied Entomology and Zoology* 37: 295–298.
- Nascimento ARP, Mariano RLR (2004) Cancro bacteriano da videira: etiologia, epidemiologia e medidas de controle. *Ciência Rural, Santa Maria* 34: 301–307.
- Nascimento ARP, Michereff SJ, Mariano R de LR, Gomes AMA (2005) Elaboração e validação de escala diagramática para cancro bacteriano da videira. *Summa Phytopathologica* 31: 59–64.
- Naumann ID (1993) *CSIRO Handbook of Australian Insect Names: Common and scientific names for insects and allied organisms of economic and environmental importance*. CSIRO Publications, East Melbourne, Victoria, Australia.
- Navarrete F, Abreo E, Betucci L, Lupo S (2010) Pathogenicity of fungi less frequently associated with trunk disease in grapevines. In Abstracts of oral and poster presentations

- given at the 7<sup>th</sup> International Workshop on Grapevine Trunk Diseases, Santa Cruz, Chile, 17–21 January 2010. *Phytopathologia Mediterranea* 49: 105.
- NCBI (2010) Necrovirus. National Centre for Biotechnology Information.  
<http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=12052> Accessed September 2010.
- Neate SM, Cruickshank RH, Rovira AD (1988) Pectic enzyme patterns of *Rhizoctonia solani* isolates from agricultural soils in South Australia. *Transactions of the British Mycological Society* 90: 37–42.
- NHB (2007) *Grapes – Pests*. National Horticulture Board, India.  
[http://nhb.gov.in/bulletin\\_files/fruits/grape/gra008.pdf](http://nhb.gov.in/bulletin_files/fruits/grape/gra008.pdf) Accessed May 2010.
- Nicholas P, Magarey P, Wachtel M (1994) *Diseases and Pests. Grape Production Series Number 1*. Winetitles, Adelaide.
- Nickel H, Remane R (2002) Check list of the planthoppers and leafhoppers of Germany, with notes on food plants, diet width, life cycles, geographic range and conservation status (Hemiptera, Fulgoromorpha and Cicadomorpha).  
[http://wwwuser.gwdg.de/~hnickel/brdlist\\_eng.pdf](http://wwwuser.gwdg.de/~hnickel/brdlist_eng.pdf) Accessed October 2011.
- Nielsen ES, Edwards ED, Rangsi TV (1996) *Checklist of the Lepidoptera of Australia*. Commonwealth Scientific and Industrial Research Organisation, Collingwood.
- Nietschke BS, Borchert DM, Magarey RD, Ciomperlik MA (2008) Climatological potential for *Scirtothrips dorsalis* (Thysanoptera: Thripidae) establishment in the United States. *Florida Entomologist* 91: 79–86.
- NIIR (2004) *Cultivation of fruits, vegetables and floriculture*. National Institute of Industrial Research. New Delhi, India.
- Nishijima T, Terai Y, Kunugi Y (2000) Studies on the *Grapevine berry necrosis virus* disease. I. Symptoms on vines, varietal susceptibility and natural spread. *Bulletin of the Yamanashi Fruit Tree Experimental Station* 67: 47–56.
- Norton M (1991) Timing *Bacillus thuringiensis* insecticides for omnivorous leafroller control in grapes. *Components* 2: 5–6.
- Novák JB, Lanzová J (1976) Identification of *Alfalfa mosaic virus* and *Tomato bushy stunt virus* in hop (*Humulus lupulus* L.) and grapevine (*Vitis vinifera* subsp. *sativa* (DC/HEGI) plants in Czechoslovakia (Brief Communication). *Biologia Plantarum* 18: 152–154.
- NPQS (2007) *PRA materials of grapes from Korea*. National Plant Quarantine Service, Korea.
- NVHSC (2005) *Pest risk analysis for the establishment of phylloxera in Australia as a result of harvest processes*. National Vine Health Steering Committee, South Australia.
- OARDC (2008) Insect and mite pests of grapes in Ohio and the Midwest. Ohio Agricultural Research and Development Centre. The Ohio State University. Ohio, USA.  
<http://www.oardc.ohio-state.edu/grapeipm/index.htm> Accessed March 2008.
- Ochoa-Corona FM, Lebas BS, Tang JZ, Bootten TJ, Stewart FJ, Harris R, Elliott DR, Alexander BJR (2006) RT-PCR detection and strain typing of *Raspberry ringspot virus*.

- Proceedings of the XXth International Symposium on Virus and Virus-like Diseases of Temperate Fruit Crops & XIth International Symposium on Small Fruit Virus Diseases, Antalya, Turkey.*
- OEPP/EPPO (1982) Data sheets on quarantine organisms No. 102, *Tomato ringspot virus*. *Bulletin OEPP/EPPO Bulletin* 12: 1.
- OEPP/EPPO (1984) Data sheets on quarantine organisms No. 133, *Xanthomonas ampelina*. *Bulletin OEPP/EPPO Bulletin* 14: 39–43.
- OHU (2010) Insects attacking grape foliage. Ohio State University. [http://www.oardc.ohio-state.edu/grapeipm/insects\\_attacking\\_grape\\_foliage.htm](http://www.oardc.ohio-state.edu/grapeipm/insects_attacking_grape_foliage.htm) Accessed May 2010.
- Okafor UA, Emezue TN, Okochi VI, Onyegeme-Okerenta BM, Nwodo-Chinedu S (2007) Xylanase production by *Penicillium chrysogenum* (PCL501) fermented on cellulosic wastes. *African Journal of Biochemistry Research* 1: 048–053.
- Oklahoma State University (2011) *Arabidopsis Mosaic Virus*. <http://www.ento.okstate.edu/ddd/diseases/armv.htm> Accessed September 2011.
- Oliver JE, Fuchs MF (2011) Fanleaf degeneration/decline disease of grapevines. [www.nysipm.cornell.edu/factsheets/grapes/diseases/fanleaf.pdf](http://www.nysipm.cornell.edu/factsheets/grapes/diseases/fanleaf.pdf) Accessed September 2011.
- Olivier CY, Lowery DT, Stobbs LW (2009a) Phytoplasma diseases and their relationships with insect and plant hosts in Canadian horticultural and field crops. *The Canadian Entomologist* 141: 425–462.
- Olivier CY, Lowery DT, Stobbs LW, Vincent C, Galka B, Saez J, Bittner L, Johnson R, Rott M, Masters C, Green M (2009b) First report of Aster yellow phytoplasmas ('*Candidatus Phytoplasma asteris*') in Canadian grapevines. *Plant Disease* 93: 669.
- Ono Y (2000) Taxonomy of the *Phakopsora ampelopsidis* species complex on vitaceous hosts in Asia including a new species, *P. euvitidis*. *Mycologia* 92: 154–173.
- Ono Y, Adhikari MK, Rajbhandari KR (1990) Uredinales of Nepal. *Report of the Tottori Mycological Institute* 28: 57–75.
- OSU (2010) Insects – The tall Fescue of the twenty-first century monograph. Oregon State University (OSU). <http://forages.oregonstate.edu/tallfescuemonograph/preface> Accessed November 2010.
- Ouertani R, Savino V, Minafra A, Boscia D, Castellano MO, Martelli GP, Greco N (1992) Properties of a previously undescribed grapevine nepovirus from Tunisia. *Archives of Virology* 126: 107–117.
- Owen J (1991) *The Ecology of a Garden: The First Fifteen Years*. Cambridge University Press, England.
- Pacifico D, Alma A, Bagnoli B, Foissac X, Pasquini G, Tessitori M, Marzachi C (2009) Characterization of Bois noir isolates by restriction fragment length polymorphism of a Stolbur-specific putative membrane protein gene. *Phytopathology* 99: 711–715.
- Panagopoulos CG (1969) The disease 'Tsilik marasi' of grapevine: its description and identification of the causal agent (*Xanthomonas ampelina* sp. nov.). *Annales de l'Institut Phytopathologique Benaki* 9: 59–81.

- Panagopoulos CG (1987) Recent research progress on *Xanthomonas ampelina*. *Bulletin OEPP/EPPO Bulletin* 17: 225–230.
- Panagopoulos CG (1988) *Xanthomonas ampelina* Panagopoulos. In Smith, I. M. Dunez, J. Lelliot, R. A. Phillips, D. H. Archer, S. A (eds) *European handbook of plant diseases* 157–158. Blackwell Scientific Publications, Oxford.
- Paoletti E, Anselmi N, Franceschini A (2007) Pre-exposure to ozone predisposes oak leaves to attacks by *Diplodia corticola* and *Biscogniauxia mediterranea*. *The Scientific World Journal* 7: 222–230.
- Papademetriou MK, Dent FJ (2001) Grape production in Asia Pacific. FAO Regional Office for Asia and the Pacific. Bangkok, Thailand.  
<http://www.fao.org/docrep/003/x6897e/x6897e00.HTM> Accessed April 2010.
- Paradies F, Finetti Sialer M, Gallitelli D, Castellano MA, Di Franco A, Digiario M, Martelli GP, Yilmaz MA (2000) Partial characterization of *cucumber mosaic virus* isolates from citrus and grapevine. *Journal of Plant Pathology* 82: 133–145.
- Park E, Choi K, Kim J, Cho S, Kim G (2001) Effect of temperature on development and reproduction of the persimmon fruit moth, *Stathmopoda masinissa* (Lepidoptera: Stathmopodidae). *Korean Journal of Applied Entomology* 40: 297–300.
- Parmenter KS, Thomas JE and Parry JN (2009) *Cherry leaf roll virus* isolate 2005 3' UTR. National Centre for Biotechnology Information, GenBank, Accession code GQ370453.  
<http://www.ncbi.nlm.nih.gov/nuccore/255964967> Accessed March 2011.
- Pascoe IG, Cunnington JH, Fischer M, Edwards J (2005) Basidiomycetous pathogens on grapevine: a new species from Australia – *Fomitiporia australiensis*. *Mycotaxon* 91: 85–96.
- Paxton DW, Thorvilson HG (1996) Oviposition of three *Erythroneura* species on grape leaves in western Texas. *Southwestern Entomologist* 21: 141–144.
- Peacock W (1992) Western grape rootworm. In *Grape Pest Management* 2nd edn, Publication 3343: 239–240. University of California Division of Agriculture and Natural Resources, Oakland, CA.
- Pearson PC, Goheen AC (1988) *Compendium of grape diseases*. APS Press St Paul, Minnesota.
- Pegler DN, Laessøe T, Spooner BM (1995) *British puffballs, earthstars and stinkhorns*. Royal Botanic Gardens, Kew.
- Peixoto AR, Mariano RLR, Moreira JOT, Viana IO (2007) Alternative hosts of *Xanthomonas campestris* pv. *viticola*. *Fitopatologia Brasileira* 32: 161–164.
- Pérez G, Lupo S, Bettucci L (2008) Polymorphisms of the ITS region of *Inocutis jamaicensis* associated with *Eucalyptus globulus*, *Vitis vinifera* and native plants in Uruguay. *Sydowia* 60: 267–275.
- Péros JP, Berger G, Jamaux-Despréaux I (2008) Symptoms, wood lesions and fungi associated with esca in organic vineyards in Languedoc-Roussillon (France). *Journal of Phytopathology*: 156: 297–303.
- Perry EJ (2002) Hoplia beetle – Integrated pest management for home gardeners

<http://www.plantmeds.com/pnhopliabeetle.pdf> Accessed March 2011.

Persley D, Gambley C (2010) Viruses in vegetable crops in Australia. Integrated virus disease management.

[http://www.dpi.qld.gov.au/documents/PlantIndustries\\_FruitAndVegetables/Viruses-in-vegies.pdf](http://www.dpi.qld.gov.au/documents/PlantIndustries_FruitAndVegetables/Viruses-in-vegies.pdf) Accessed September 2010.

Persley DM, Thomas JE, Sharman M (2006) Tospoviruses; an Australian perspective. *Australasian Plant Pathology* 35: 161–180.

Petrini LE, Petrini O (2005) Morphological studies in *Rosellinia* (Xylariaceae): the first step towards a polyphasic taxonomy. *Mycological Research* 109: 569–580.

Pfeiffer DG, Schultz PB (1986) *Major Insect and Mite Pests of Grape in Virginia*. Virginia Cooperative Extension Service, USA.

Pfeilstetter E, Zinkernagel V, Kunze L (1992) Occurrence of *Petunia asteroid mosaic* (PAMV) and *Carnation Italian ringspot* (CIRV) viruses in cherry orchards in northern Bavaria. *Acta Horticulturae* 309: 345–352.

PHA (Plant Health Australia) (2001). *Australian Plant Pest Database*, online database. [www.planthealthaustralia.com.au/appd](http://www.planthealthaustralia.com.au/appd) Accessed May 2011.

Phillips AJL (2000) Excoriose, cane blight and related disease of grapevines: a taxonomic review of the pathogens. *Phytopathologia Meditteranea* 39: 341–356.

Phillips AJL, Alves A, Pennycook SR, Johnston PR, Ramaley A, Akulov A, Crous PW (2008) Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the Botryosphaeriaceae. *Persoonia* 21: 29–55.

Phillips C (1994) Leaf diseases

[http://outernode.pir.sa.gov.au/forestry/publications\\_index/forest\\_health\\_information/factSheets/leaf\\_diseases](http://outernode.pir.sa.gov.au/forestry/publications_index/forest_health_information/factSheets/leaf_diseases) Assessed January 2011.

Pilotti M, Tizzani L, Brunetti A, Gervasi F, Di Lernia G, Lumia V (2010) Molecular identification of *Fomitipora mediterranea* on declining and decayed hazelnut. *Journal of Plant Pathology* 92: 115–129.

Pimentel DP (2007) *Encyclopedia of Pest Management*. Volume II. CRC Press, USA.

Pitt WM, Huang R, Steel CC, Savocchia S (2009) Identification, distribution and current taxonomy of Botryosphaeriaceae species associated with grapevine decline in New South Wales and South Australia. *Australian Journal of Grape and Wine Research* 16: 258–271.

Pitta G PB (1994) Diseases of palms in Brazil. *Acta Horticulturae* 360: 231–234.

Pittaway AR, Kitching IJ (2006) Sphingidae of the eastern Palaearctic. <http://tpittaway.tripod.com/china/china.htm> Accessed September 2010.

Plaut HN (1973) On the biology of *Paropta paradoxus* (H.-S.) (Lep., Cossidae) on grapevine in Israel. *Bulletin of Entomological Research* 63: 237–245.

Pospieszny H, Borodynko N, Jonczyk M (2004) First report of *Tomato black ring spot virus* in the natural infection of *Sambucus nigra* in Poland. *Journal of Plant Protection Research* 44: 373–376.

- Pourrahim R, Ahoonmanesh A, Farzadfar SH, Rakhshandehro F (2004) Occurrence of *Arabidopsis mosaic virus* and *Grapevine leaf roll associated virus-3* on Grapevines in Iran. *Plant Disease* 88: 424.
- Pozdena J, Vanek G, Filigarova M (1977) *Sowbane mosaic virus* in grapevine in the CSSR. (In Czech). *Sb.UVTIZ-Ochrana Rostlin* 13: 197–200.
- Pozsár BI, Horváth L, Lehoczky J, Sárospataki G (1969) Effect of the *Grape chrome mosaic* and *Grape fanleaf yellow mosaic virus* infection on the photosynthetic carbon dioxide fixation in vine leaves. *Vitis* 8: 206–210.
- Prince JP, Davis RE, Wolf TK, Lee IM, B. Mogen *et al.* (1993) Molecular detection of diverse Mycoplasma-Like Organisms (MLOs) associated with Grapevine yellows and their classification with Aster yellows, X-disease and Elm yellows MLOs. *Phytopathology* 83: 1130–1137.
- Prodi A, Sandalo S, Tonti S, Nipoti P, Pisi A (2008) *Phialophora*-like fungi associated with kiwifruit elephantiasis. *Journal of Plant Pathology* 90: 487–494.
- Purcell A H (2001) *Xylella fastidiosa* Web Site. University of California at Berkeley. <http://www.cnr.berkeley.edu/xylella/index.html> Accessed March 2011.
- Qui Y, Steel CC, Ash GJ, Savocchia S (2011) Survey of Botryosphaeriaceae associated with grapevine decline in the hunter valley and Mudgee grape growing regions of New South Wales. *Australasian Plant Pathology* 40: 1–11.
- Ragazzini D (1996) *Sinoxylon perforans*: a pest of our grape vineyards. *Informatore Fitopatologico* 27: 3–16.
- Ramsdell DC, Gillet JM (1998) *Peach rosette mosaic virus*. AAB Description of Plant Viruses, No. 364. <http://www.dpvweb.net/dpv/showdpv.php?dpvno=364> Accessed September 2011.
- Ramsdell DC, Milholland RD (1988) Black rot. In Pearson RC, Goheen AC (eds) *Compendium of grape diseases* 15–17. American Phytopathological Society Press, St. Paul.
- Randles JW (1986) Susceptibility of *Echium plantagineum* L. to *Tobacco mosaic*, *Alfalfa mosaic*, *Tobacco ringspot* and *Tobacco necrosis viruses*. *Australasian Plant Pathology* 15: 74–77.
- Ranga Rao PV, Azam KM, Laxminarayana K, Eshbaugh EL (1979) A new record of *Coelosterna scabrator* F. (Cerambycidae: Coleoptera) on grapevines in Andhra Pradesh. *Indian Journal of Entomology* 41: 289–290.
- Rao R (1965) A new species of *Amphisphaeria* from India. *Mycopathologia* 28: 239–240.
- Rao R (1966) A new species of *Teichospora* from India. *Mycopathologia* 28: 63–64.
- Raski DJ, Goheen AC, Lider LA, Meredith CP (1983) Strategies against *Fanleaf virus* and its nematode vector. *Plant Disease* 67: 335–339.
- Rawat RR, Jakhmola SS (1970) Bionomics of the mango-coccid (*Rastrococcus iceryoides* Green; Homoptera: Coccidae). *Indian Journal of Agricultural Sciences* 40: 140–144.
- Rebenstorf K, Candresse T, Marie Josee Dulucq MJ, Büttner C and Obermeier C (2006) Host species-dependent population structure of a pollen-borne plant virus, *Cherry leaf roll virus*.



*Journal of Virology* 80: 2453–2462.

Regniere J, Rabbn RL, Stinner RE (1983) *Popillia japonica* (Coleoptera: Scarabaeidae): Distribution and movement of adults in heterogeneous environments. *Canadian Entomology* 115: 287–294.

Reisenzein H, Berger N, Nieder G (2000) Esca in Austria. *Phytopathologia Mediterranea* 39: 16–20.

Reitz SR (2002) Seasonal and within plant distribution of *Frankliniella* thrips (Thysanoptera: Thripidae) in north Florida tomatoes. *Florida Entomology* 85: 431–439.

Reynolds DR (1971) On the use of hyphal morphology in the taxonomy of sooty molds. *Taxon* 20: 759–768.

Rezaian MA (1990) Australian grapevine viroid-evidence for extensive recombination between viroids. *Nucleic Acids Research* 18: 1813–1818.

Rhoads AS (1918) *Daldinia vernicosa*: a pyroxylophilous fungus. *Mycologia* 10: 277–284.

Riccioni L, Manning M, Valvassori M, Haegi A, Casonato S, Spinelli R (2007) A new disease: leader die-back in *Actinidia chinensis* ‘Hort16A’ in Italy. *Acta Horticulturae* 753: 669–676.

Rice M, Drees BM (1985) Oviposition and girdling habits of the Three-cornered Alfalfa hopper (Homoptera: Membracidae) on prebloom soybeans. *Journal of Economic Entomology* 78: 829–834.

Richardson MJ (1990) *An annotated list of seed-borne disease* 4<sup>th</sup> edn. The Internal Seed Testing Association, Switzerland.

Ridé M, Ridé S, Novoa D (1977) Données nouvelles sur la biologie de *Xanthomonas ampelina* Panagopoulos, agent de la nécrose bactérienne de la vigne. *Annales de Phytopathologie* 9: 87.

Ridé M, Ridé S, Novoa D (1983) Connaissances actuelles sur la nécrose bactérienne de la vigne. *Bulletin Technique des Pyrénées-Orientales* 106: 10–45.

Ridgway H, Whiteman SA, Jaspers MV, Stewart A (2003) Molecular diagnostics for industry: sources of Petri disease in grapevine nurseries. *Phytopathologia Mediterranea* 43: 152.

Riedl H, Taschenberg EF (2008) Integrated pest management – Grape cane girdler, Cornell University, New York State Agricultural Experiment Station, Geneva.  
<http://www.nysipm.cornell.edu/factsheets/grapes/pests/gcg/gcg.asp> Accessed February 2010.

Riedle-Bauer M, Tiefenbrunner W, Ostreba J, Hanak K, Schildberger B, Regner F (2006) Epidemiological observations on Bois noir in Austrian vineyards. *Mitteilungen Klosterneuburg* 56: 166–170.

Rieger M (2005) *Introduction to fruit crops*. Haworth Press. Philadelphia, USA.

Riolo P, Landi L, Nardi S, Isidoro N (2007) Relationships among *Hyalesthes obsoletus*, its herbaceous host plants and ‘Bois noir’ strains in vineyard ecosystems in the Marche region (central-eastern Italy). *Bulletin of Insectology* 60: 353–354.

Robinson GS, Ackery PR, Kitching IJ, Beccaloni GW, Hernández LM (2008) Hosts: a

- database of the host plants of the world's Lepidoptera host plants. The Natural History Museum, London. <http://www.nhm.ac.uk/entomology/hostplants/> Accessed April 2010.
- Roca F, Martelli GP, Lamberti F, Rana GL (1975) Distribution of *Longidorus attenuatus* Hooper in Apulian artichoke fields and its relationship with *Artichoke Italian latent virus*. *Nematologia Mediterranea* 3: 91–101.
- Roditakis E, Roditakis NE (2007) Assessment of the damage potential of three thrips species on white variety table grapes—In vitro experiments. *Crop Protection* 26: 476–483.
- Roehrich R, Boller E (1991) Tortricids in vineyards. In Van der Geest LPS, Evenhuis HH (eds) *Tortricid pests, their biology natural enemies and control* 507–514. Elsevier, Amsterdam.
- Rogiers SY, Whitelaw-Weckert M, Radovanovic-Tesic M, Greer LA, White RG, Steel CC (2005) Effects of spray adjuvants on grape (*Vitis vinifera*) berry microflora, epicuticular wax and susceptibility to infection by *Botrytis cinerea*. *Australasian Plant Pathology* 34: 221–228.
- Rolshausen P (2004) *Biology and control of Eutypa dieback of grapevine in California*. University of California, Davis.
- Romanazzi G, Murolo S, Pizzichini L, Nardi S (2009) Esca in young and mature vineyards, and molecular diagnosis of the associated fungi. *European Journal of Plant Pathology* 125: 277–290.
- Romero AI, Carmaran CC (2003) First contribution to the study of *Cryptosphaeria* from Argentina. *Fungal Diversity* 12: 161–167.
- Romero D, Rivera ME, Cazorla FM, De Vicente A, Perez-Garcia A (2003) Effect of mycoparasitic fungi on the development of *Sphaerotheca fusca* in melon leaves. *Mycological Research* 107: 64–71.
- Ronald FL, Jayma LK (1992) *Listroderes difficilis* (germar). Crop Knowledge Master, Extension Entomology & UH-CTAHR Integrated Pest Management Program. <http://www.extento.hawaii.edu/kbase/crop/Type/listrode.htm> Accessed February 2010.
- Rooney-Latham S, Eskalen A, Gubler WD (2005) Teleomorph formation of *Phaeoacremonium aleophilum*, cause of esca and Grapevine decline in California. *Plant Disease* 89: 177–184.
- Rosenstiel RG, Ferguson GR (1944) Some ecological relationships of *Cnephasia longana*. *Journal of Economic Entomology* 37: 814–817.
- Rott M, Johnson R, Masters C, Green M (2007) First report of Bois noir Phytoplasma in grapevine in Canada. *Plant Disease* 91: 1682.
- Rowhani A, Mircetich SM (1988) Pathogenicity on walnut and serological comparisons of *Cherry leaf roll virus* strains. *Phytopathology* 78: 817–820.
- Rowhani A, Uyemoto JK, Golino DA, Martelli GP (2005) Pathogen testing and certification of *Vitis* and *Prunus* species. *Annual Review of Phytopathology* 43: 261–278.
- Ruan-Soto F, Garibay-Orijel R, Cifuentes J (2006) Process and dynamics of traditional selling wild edible mushrooms in tropical Mexico. *Journal of Ethnobiology and Ethnomedicine* 2: 3.
- Rudel M (1985). Grapevine damage induced by particular virus-vector combinations.

*Phytopathologia Mediterranea* 24: 183–185.

Russo M, Vovlas C, Rubino L, Grieco F, Martelli GP (2002) Molecular characterization of a tombusvirus isolated from diseased pear trees in southern Italy. *Journal of Plant Pathology* 84: 161–166.

Rust MK (1992) Termites. In Flaherty DL, Christensen PL, Lanini TW, Marois JJ, Phillips PA, Wilso LT (eds) *Grape pest management*. Publication No. 3343. University of California, Division of Agriculture and Natural Resources Oakland, California.

Sabanadzovic S, Abou-Ghanem N, Castellano MA, Digiario M, Martelli GP (2000) *Grapevine fleck virus*-like viruses in *Vitis*. *Archives of Virology* 145: 553–565.

Saeki K, Takahashi Y, Oh-Oka H, Umeoka T, Oda Y and Fukuyama K (2001) Primary structure and phylogenetic analysis of the coat protein of a Toyama isolate of *Tobacco necrosis virus*. *Bioscience, Biotechnology and Biochemistry* 65: 719–724.

SAG/USDA (2002) Importation of fresh commercial citrus fruit: clementine (*Citrus reticulata* Blanco var. *clementine*), mandarin (*Citrus reticulata* Blanco) and tangerine (*Citrus reticulata* Blanco) from Chile into the United States. A pathway initiated plant pest risk assessment. Draft March 2002. Departamento Protección Agrícola, Servicio Agrícola Ganadera (Santiago, Chile) and United States Department of Agriculture, Plant Protection and Quarantine (Riverdale, Maryland). <http://www.aphis.usda.gov/ppq/prac/clementine.pdf> Accessed March 2011.

Sambath S, Joshi KC (2004) Bionomics of *Spirama retorta* Cram. (Lepidoptera: Noctuidae). *Indian Journal of Entomology* 66: 102–106.

Santini D, Rolle L, Mannini F (2009) Control of grape berry mechanical properties modifications due to GFLV by mean of texture analysis. *Extended abstracts of the 16th meeting of ICVG in Dijon, 31 August–4 September 2009* 258–259. ICVG, Switzerland.

Santos MT, Mansinho A, Correia PK, Sequeira OA, Nolasco G (2000) Multiple detection of grapevine filamentous viruses in Portugal, by RT-PCR from DS-RNA templates. <http://www.icvg.ch/data/abstrb.pdf> Accessed September 2010.

Savino V, Gallitelli D, Jankulova M, Rana GL (1977) A comparison of four isolates of *Artichoke Italian latent virus* (AILV). *Phytopathologia Mediterranea* 16: 41–50.

Savino V, Martelli GP, D'Onghia AM, Yilmaz MA (1987) Turkey. *Strawberry latent ringspot virus* in grapevine. *FAO Plant Protection Bulletin* 35: 102–104.

Savocchia S, Greer LA, Steel CC (2007) First report of *Phomopsis viticola* causing bunch rot of grapes in Australia. *Plant Pathology* 56: 725.

Schaefer CW, Panizzi AR (2000) *Heteroptera of economic importance*. CRC Press; Boca Raton, Florida.

Schell S, Lockwood J, Schell S, Zimmerman K (2007) Grasshoppers of Wyoming and the West. University of Wyoming. <http://www.wygisc.uwyo.edu/grasshopper/ghwywfrm.htm> Accessed May 2008.

Scheper RWA, Crane DC, Whisson DL, Scott ES (2000) The *Diaporthe* teleomorph of *Phomopsis* taxon 1 on grapevine. *Mycological Research* 104: 226–231.

- Schilder A (2011) Diseases of grapes: Tomato/tobacco ringspot virus decline – *Tomato ringspot virus*, *Tobacco ringspot virus*. <http://grapes.msu.edu/ringspot.htm> Accessed September 2011.
- Schloemann S (2003–2004) New England small fruit pest management guide. United States Department of Agriculture, Cooperative Extension Service, University of Massachusetts, Amherst. <http://www.umass.edu/fruitadvisor/nesfpmg/> Accessed November 2010.
- Schmidt LS, Ghosop JM, Margosan DA, Smilanick JL (2006) Mutation at  $\beta$ -Tubulin codon 200 indicated thiabendazole resistance in *Penicillium digitatum* collected from California citrus packinghouses. *Plant Disease* 90: 765–770.
- Schneider B, Seemüller E, Smart CD and Kirkpatrick BC (1995) Phylogenetic classification of plant pathogenic mycoplasma-like organisms or phytoplasmas. In Razin S, Tully JG (eds) *Molecular and diagnostic procedures in mycoplasmaology* 369–380. Academic Press, San Diego, USA.
- Schoorl JW Jr (1990) A phylogenetic study on Cossidae (Lepidoptera: Ditrysia) based on external adult morphology. *Zoologische Verhandelingen* 263: 295 pp.
- Schroers HJ, Žerjav M, Munda A, Halleen F, Crous PW (2008) *Cylindrocarpon pauciseptatum* sp. nov., with notes on *Cylindrocarpon* species with wide, predominantly 3-septate macroconidia. *Mycological Research* 112: 82–92.
- Schwartz A (1986) Leaf curl mite in vineyards. *Viticulture and Oenology F.* 28. South Africa: South Africa Department of Agriculture and Water Supply, South Africa.
- Schwinghamer M, Schilg M, Moore K, Knights T, Bambach R, Southwell R, Kumari S, Srivastava M, Wratten K (2007) Virus situation in chickpea, faba beans and canola. Grains Research and Development Corporation. [http://www.grdc.com.au/director/events/onfarmtrials?item\\_id=4829BB0CA9AC5A109CF7A53B484E5905&pageNumber=2](http://www.grdc.com.au/director/events/onfarmtrials?item_id=4829BB0CA9AC5A109CF7A53B484E5905&pageNumber=2) Accessed January 2010.
- Seaver FJ (1911) Studies of Colorado fungi: I. Discomycetes. *Mycologia* 3: 57–66.
- Seeman OD, Beard JJ (2011) Identification of exotic pest and Australian native and naturalised species of *Tetranychus* (Acari: Tetranychidae). *Zootaxa* 2961: 1–72.
- Seemuller E, Schneider B, Ma UR (1994) Phylogenetic classification of phytopathogenic mollicutes by sequence analysis of 16S ribosomal DNA. *International Journal of Systematic Bacteriology* 44: 440–446.
- Seljak G (2010) A checklist of scale insects of Slovenia. *Entomologia Hellenica* 19: 99–113.
- Sequeira OA, Mendonça A (1992) Certification of grapevine in Portugal. In *Grapevine viruses and certification in EEC countries: state of the art*. CIHEAM, Bari, Italy.
- Serfontein S, Serfontein TJ, Botha WI, Staphorst JL (1997) The isolation and characterisation of *Xylophilus ampelinus*. *Vitis* 36: 209–210.
- Sergeeva V, Nair T, Spooner-Hart R (2001) *Fungi recorded on grapevines during the course of an industry service on Botrytis monitoring and fungicide resistance*. Annual Technical Issue, Australian and New Zealand Grapegrower and Winemaker, South Australia.
- Sergeeva V, Priest M, Nair NG (2005) Species of *Pestalotiopsis* and related genera occurring

- on grapevines in Australia. *Australasian Plant Pathology* 34: 255–258.
- Serra S, Borgo M, Zanzotto A (2000) Investigation into the presence of fungi associated with esca of young vines. *Phytopathologia Meditteranea* 39: 21–25.
- Seung-Tae K, Jung MP, Kim HS, Shin JH, Lim JH, Kim TW, Lee JH (2006) Insect fauna of adjacent areas of DMZ in Korea. *Journal of Economical and Field Biology* 29: 125–141.
- Sforza R, Clair D, Daire X, Larrue J, Boudon-Padieu E (1998) The role of *Hyalesthes obsoletus* (Hemiptera: Cixiidae) in the occurrence of Bois noir of grapevines in France. *Journal of Phytopathology* 146: 549–556.
- Shao-Hua L (2012) Grape production in China. FAO Corporate Document Repository. <http://www.fao.org/docrep/003/x6897e/x6897e05.htm> Accessed February 2012.
- Sharma RC, Sharma S (2006) Short rotation forestry for industrial and rural development. *Proceedings of the IUFRO-ISTS-UHF International Conference on World Perspective on Short Rotation Forestry for Industrial and Rural Development*, Nauni, Solan, India, 7–13 September 2006 pp. 364–370. Westville Publishing House, New Delhi, India.
- Sharma RL, Kaul JL (1989) Host range of brown rot fungi (*Monilinia* spp.) in India. *Plant Disease Research* 4: 177–178.
- Shim JO, Cho KD, Hahn KD, *et al.* (2002) Blue mold of pear caused by *Penicillium aurantiogriseum* in Korea. *Mycobiology* 30: 105–106.
- Shivas RG (1989) Fungal and bacterial diseases of plants in Western Australia. *Journal of the Royal Society of Western Australia* 72: 1–62.
- Shivas RG, Tan YP (2009) A taxonomic re-assessment of *Colletotrichum acutatum*, introducing *C. fioriniae* comb. et stat. nov. and *C. simmondsii* sp. nov. *Fungi Diversity* 39: 111–122.
- Siegel JP, Lacey LA, Higbee BS, Noble PM, Fritts RJ (2006) Effect of application rates and abiotic factors on *Steinernema carpocapsae* for control of overwintering Navel orangeworm (Lepidoptera: Pyralidae, *Amyelois transitella*) in pistachios. *Biological Control* 36: 324–330.
- Silverside AJ (2006) *Cicadella viridis* (L.) (a leafhopper). [http://bioref.lastdragon.org/Hemiptera/Cicadella\\_viridis.html](http://bioref.lastdragon.org/Hemiptera/Cicadella_viridis.html) Accessed October 2011.
- Simmonds JH (1966) *Host index of plant diseases in Queensland*. Queensland Department of Primary Industries, Brisbane.
- Sinclair WA, Lyon HH, Johnson WT (1987) *Diseases of trees and shrubs*. Cornell University Press, Ithaca, NY USA 575 pp.
- Siswanto, Muhamad R, Omar D, Karmawatic E (2008) Dispersion pattern of *Helopeltis antonii* signoret (Hemiptera: Miridae) on cashew plantation. *Indonesian Journal of Agriculture* 1: 103–108.
- Sivanesan A, Holliday P (1972) *Xylaria-polymorpha*. CMI (Commonwealth Mycological Institute) Descriptions of Pathogenic Fungi and Bacteria 36.
- Sivichai S, Jones EBG, Hywel-Jones NL (2000) Fungal colonisation of wood in a freshwater stream at Khao Yai National Park, Thailand. *Fungal Diversity* 5: 71–88.

- Smith D, Papacek DF, Murray DAH (1988) The use of *Leptomastix dactylopii* Howard (Hymenoptera: Encyrtidae) to control *Planococcus citri* (Risso) (Hemiptera: Pseudococcidae) in Queensland citrus orchards. *Queensland Journal of Agricultural and Animal Sciences* 45: 157–164.
- Smith ESC (1996) Control of longicorn beetles in Mango trees. Northern Territory Department of Primary Industries, Darwin. *Agnote* 685: 1–2
- Smith IM, McNamara DG, Scott PR, Holderness M (1997) *Quarantine pests for Europe. Second Edition. Data sheets on quarantine pests for the European Union and for the European and Mediterranean Plant Protection Organization*. CABI/EPPO, Wallingford, UK.
- Smith LM, Stafford EM (1948) The bud mite and the erineum mite of grapes. *Hilgardia* 18: 317–334.
- Solomon JD (1995) *Guide of insect borers of North American broadleaf trees and shrubs*. Agriculture Handbook 706. US Department of Agriculture, Forest Service, Washington DC.
- Sosnowski MR, Lardner R, Wicks TJ, Scott ES (2007) The influence of grapevine cultivar and isolate of *Eutypa lata* on wood and foliar symptoms. *Plant Disease* 91: 924–931.
- Spadaro D, Pellegrino C, Garibaldi A, Gullino ML (2011) Development of SCAR primers for the detection of *Cadophora luteo-olivacea* on kiwifruit and pome fruit and of *Cadophora malorum* on pome fruit. *Phytopathologia Mediterranea* 50: 430–441.
- Sparapano L, Bruno G, Ciccarone C, Graniti A (2000) Infection of grapevines by some fungi associated with esca. I. *Fomitiporia punctata* as a wood-rot induce. *Phytopathologia Mediterranea* 39: 46–52.
- Srivastava RP (1997) *Mango Insect Pest Management*. First edition). International Book Distributing Co., Lucknow, India.
- Srivastava RP, Verghese A (1985) Record of a new mealy bug, *Perissopneumon ferox* Newstead (Margarodidae: Homoptera) on mango from Uttar Pradesh, India. *Entomon* 10: 184.
- Stamp JA, Wei A (2013) The impact of grapevine red blotch virus. *Wine Business Monthly* March 2013: 56–67.
- Stathas GJ, Kontodimas DC (2001) Ecological data of the scale *Targionia vitis* on grapes in Southern Greece. *Annales de l'Institut phytopathologique Benaki* 19: 134–139.
- Steffek R, Reisenzein H, Zeisner N (2006) Analysis of pest risk from grapevine Flavescence dorée Phytoplasma to Austrian viticulture. *EPPO Bulletin* 37: 191–203.
- Stellmach G, Querfurth G (1978) Untersuchungen zur serologie, pathologie und thermolabilität mehrerer Raben-isolate des Himbeeringflecken-virus (*Raspberry latent virus*). *Weinberg und Keller* 25: 128–136.
- Stern VM, Johnson JA (1984) Biology and control of the Grape bud beetle, *Glyptoscelis squamulata* (Coleoptera: Chrysomelidae), in southern California table grapes. *Journal of Economic Entomology* 77: 1327–1334.
- Stirling GR, Stanton JM, Marshall JM (1992) The importance of plant-parasitic nematodes to Australian and New Zealand agriculture. *Australasian Plant Pathology* 21: 104–115.
- Stobbs LW, van Schagen JG (1984) Occurrence of *Tomato black ring virus* on grapevine in

- Southern Ontario. *Canadian Plant Disease Survey* 64: 3–4.
- Stracener CL (1931) Economic importance of the salt-marsh caterpillar (*Estigmene acraea* Drury) in Louisiana. *Journal of Economic Entomology* 24: 835–838.
- Stukely MJC, Webster JL, Ciampini JA, Kerp NL, Colquhoun IJ, Dunstan WA, Hardy GE St. J (2007) A new homothallic *Phytophthora* from the jarrah forest in Western Australia. *Australasian Plant Disease Notes* 2: 49–51.
- Sudarshana MR (2012) Grapevine red blotch diseases. Agriculture Research Service, United States Department of Agriculture. <http://ucanr.edu/sites/viticulture-fresno/files/157139.pdf> Accessed 15 April 2013.
- Suhag LS, Kaushik JC, Duhan JC (1982) Etiology and epidemiology of fungal foliar diseases on grapevine. *Indian Journal of Mycology and Plant Pathology* 12: 191–197.
- Summerell BA, Leslie JF, Liew ECY, Laurence MH, Bullock S, Petrovic T, Bentley AR, Howard CG, Peterson SA, Walsh JL, Burgess LW (2011) *Fusarium* species associated with plants in Australia. *Fungal Diversity* 46: 1–27.
- Sun YZ, Qi J, Li H (1992) A preliminary report on *Scelodonta lewisii* Baly. *Plant Protection* 18: 1617.
- Sundararaju D1, Babu PCS (2000) Improved mass culture technique for the neem mosquito bug, *Helopeltis antonii* Signoret (Heteroptera: Miridae). *Journal of Entomological Research* 24: 73–82.
- Sweetingham M (1983) Studies on the nature and pathogenicity of soilborne *Cylindrocarpus* spp. Ph.D. Thesis, University of Tasmania.
- Tan SD, Wei JD, Lan RX (1998) Analysis on the similarity of the structure of the litchi and longan pest communities. *Guangxi Science and Technology of Tropical Crops* 69: 410.
- Tandon PL, Verghese A (1987) New insect pests of certain fruit crops. *Indian Journal of Horticulture* 44: 121–122.
- Tanne E, Orenstein S (1997) Identification and typing of grapevine phytoplasma amplified by graft transmission to periwinkle. *Vitis* 36: 35–38.
- Taralashvili LV (1989) *Sinoxylon perforans* on the grapevine. *Zashchita rastenii* (Translated).
- Taylor A, ST Hardy GE, Wood P, Burgess T (2005) Identification and pathogenicity of *Botryosphaeria* species associated with Grapevine decline in Western Australia. *Australasian Plant Pathology* 34: 187–195.
- Taylor CE, Chambers J, Pattullo WI (1965) The effect of *Tomato black ring virus* on the growth and yield of Mallings Exploit raspberry. *Horticultural Research* 5: 19–24.
- Taylor RH (1962) Grapevine viruses in Victoria. *Journal of Agriculture, Victoria* 60: 336.
- Taylor RH, Hewitt WB (1964) Properties and serological relationships of Australian and Californian soil-borne viruses of the grapevine and *Arabis mosaic virus*. *Australian Journal of Agricultural Research* 15: 571.
- Teakle DS (1968) *Sowbane mosaic virus* infecting *Chenopodium trigonon* in Queensland. *Australian Journal of Biological Sciences* 21: 649–53.

- Teakle DS (1988) *Tobacco necrosis necrovirus*. In Buchen-Osmond C, Crabtree K, Gibbs A, McLean G (eds) *Viruses of Plants in Australia* 490–491. Australian National University Printing Service, Canberra.
- Terai Y, Kunihi Y, Yansae H (1993) Back-transmission of a grapevine filamentous virus to grapevine seedlings and induction of foliar and berry symptoms in grapevine. *Annals of the Phytopathological Society of Japan* 53: 423.
- Tessmann DJ, Dianese JC, Genta W, Vida JB, Mio M (2004) Grape rust caused by *Phakopsora euvitis*, a new disease for Brazil. *Fitopatologia Brasileira* 29: 338.
- Thayer MK (2001) An Australian immigrant in southern California: *Paraphloeostiba gayndahensis* (Coleoptera: Staphylinidae). *The ESA 2001 Annual Meeting – 2001: An Entomological Odyssey of ESA*. [http://esa.confex.com/esa/2001/techprogram/paper\\_3670.htm](http://esa.confex.com/esa/2001/techprogram/paper_3670.htm) Accessed May 2010.
- Thomas MC, Heppner JB, Woodruff RE, Weems HV, Steck GJ, Fasulo TR (2001) Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Insecta: Diptera: Tephritidae). EENY-214. Florida Department of Agriculture and Consumer Services, Florida.
- Thompson RS, Nelson B (2003) The butterflies and moths of Northern Ireland. <http://www.ulstermuseum.org.uk/lepidoptera/species.asp?item=6384> Accessed December 2010.
- Thormann MN, Rice AV (2007) Fungi from peatlands. *Fungal Diversity* 24: 241–299.
- Thorn RG, Moncalvo J-M, Redhead SA, Lodge DJ, Martin MP (2005) A new poroid species of *Resupinatus* from Puerto Rico, with a reassessment of the cyphelloid genus *Stigmatolemma*. *Mycologia* 97: 1140–1151.
- Tomazic I, Petrovic N, Korosec-Koruza Z (2005) Effects of rugose wood and GLRaV-1 on yield of cv. 'Refosk' grapevines. *Acta Agriculturae Slovenica*. 85, 91–96.
- Tomazic I, Korošec-Koruza Z, Koruza, B (2000) GLRaV-1 and stem pitting disease – two factors affecting the yield of grapevine cv. Refosk. In *Extended Abstracts of the 13th Meeting of ICVG, March 12–18, 2000 Adelaide, Australia* 159–161. ICVG, Switzerland.
- Tomlinson JA, Faithful EM, Webb MJW, Fraser RSS and Seeley ND (1983) *Chenopodium necrosis*: a distinctive strain of *Tobacco necrosis virus* isolated from river water. *Annals of Applied Biology* 102: 135–147.
- Tovar FJ, Hardy GESTJ, Robinson RM, Burgess T (2008) Testing the wood decay capacity of fungi commonly observed on *Eucalyptus globulus* coppice stumps. In *ICPP 2008 9th International Congress of Plant Pathology, 24–29 August, Torino, Italy*.
- Trdani S, Valič N (2004) Contribution to the knowledge on bionomics of *Byctiscus betulae* L. (Coleoptera, Curculionidae) on grapevine. *Acta Agriculturae Slovenica* 83: 37–43.
- Trindade LC, Lima MF, Ferreira MASV (2005) Molecular characterization of Brazilian strains of *Xanthomonas campestris* pv. *viticola* by rep-PCR fingerprinting. *Fitopatologia Brasileira* 30: 46–54.
- Trouillas FP, Gubler WD (2010) Pathogenicity of Diatrypaceae species in grapevines in California. *Plant Disease* 94: 867–872.



- Trouillas FP, Pitt WM, Sosnowski M, Peduto F, Huang H, Loschiavo A, Savocchia S, Steel C, Scott E, Gubler WD (2010) Diatrypaceae associated with grapevine canker diseases in South Australia and New South Wales. *Phytopathologia Mediterranea* 49: 107–108.
- Trouillas FP, Pitt WM, Sosnowski MR, Huang R, Peduto F, Loschiavo A, Savocchia S, Scott ES (2011) Taxonomy and DNA phylogeny of Diatrypaceae associated with *Vitis vinifera* and other woody plants in Australia. *Fungal Diversity* 49: 203–223.
- Tsui CKM, Hyde KD, Fukushima K (2003) Fungi on submerged wood in the Koito River, Japan. *Mycoscience* 44: 55–59.
- Tzanetakis IE, Postman JD, Gergerich RC and Martin RR (2006) A virus between families: nucleotide sequence and evolution of *Strawberry latent ringspot virus*. *Virus Research* 121: 199–204.
- Uecker FA, Kuo K-C (1992) A new *Phomopsis* with long paraphyses. *Mycotaxon* 44: 425–433.
- Ullrich CI, Kleespies RG, Enders M, Koch E (2009) Biology of the black rot pathogen, *Guignardia bidwellii*, its development in susceptible leaves of grapevine *Vitis vinifera*. *Journal Fur Kulturflanzen* 61: 82–90.
- University of Illinois (2001) Black rot of grape. *Report on Plant Disease* 703: 1–4.
- University of Illinois (2004) Cottony Maple Scale, *Pulvinaria innumerabilis* [http://ipm.illinois.edu/landturf/insects/cottony\\_maple\\_scale/index.html](http://ipm.illinois.edu/landturf/insects/cottony_maple_scale/index.html) Accessed August 2011.
- Untiveros M, Perez-Egusquiza Z, Clover G (2010) PCR assays for the detection of members of the genus Ilarvirus and family Bromoviridae. *Journal of Virological Methods* 165: 97–104.
- Úrbez -Torres JR, Peduto F, Striegler RK, Urrea-Romero KE, Rupe JC, Cartwright RD, Gubler WD (2012) Characterization of fungal pathogens associated with grapevine trunk diseases in Arkansas and Missouri. *Fungal Diversity* 52: 169–189.
- Úrbez-Torres JR (2011) The status of Botryosphaeriaceae species infecting grapevines. *Phytopathologia Mediterranea* 50: (Supplement) S5–S45.
- Úrbez-Torres JR, Gubler WD (2009) Pathogenicity of Botryosphaeriaceae species isolated from grapevine cankers in California. *Plant Disease* 93: 584–592.
- Úrbez-Torres JR, Leavitt GM, Voegel TM, Gubler WD (2006) Identification and distribution of *Botryosphaeria* spp. associated with grapevine cankers in California. *Plant Disease* 90: 1490–1503.
- Úrbez-Torres JR, Peduto F, Gubler WD (2010a) First report of grapevine cankers caused by *Lasiodiplodia crassispora* and *Neofusicoccum mediterraneum* in California. *Plant Disease* 94: 785.
- Úrbez-Torres JR, Peduto F, Rooney-Latham S, Gubler WD (2010b) First report of *Diplodia corticola* causing grapevine (*Vitis vinifera*) cankers and trunk cankers and dieback of canyon live oak (*Quercus chrysolepis*) in California. *Plant Disease* 94: 785.
- USDA (1995) Importation of Japanese Unshu orange fruits (*Citrus reticulata* Blanco var. *unshu* Swingle) into citrus producing states. Pest risk assessment.

<http://naldc.nal.usda.gov/download/CAT11065905/PDF> Accessed February 2012.

USDA (2005) *Qualitative pathway-initiated risk assessment of the importation of fresh table grapes Vitis vinifera L. from Namibia into the United States*. Animal and Plant Health Inspection Service, United States Department of Agriculture, Raleigh.

Uyemoto JK (1975) A severe outbreak of virus-induced grapevine decline in cascade grapes in New York. *Plant Disease Reporter* 59: 98–101.

Uyemoto JK, Martelli GP, Rowhani (2009) Grapevine viruses, virus like diseases and other disorders. In *Virus diseases of plants: Grape, potato, and wheat image collection and teaching resource* CD-Rom. APS Press, St. Paul, MN 55121.

Uyemoto JK, Taschenberg EF, Hummer DK (1977) Isolation and identification of a strain of *Grapevine Bulgarian latent virus* in concord grapevine in New York State. *Plant Disease Reporter* 61: 949–953.

Vaartaja O (1965) New *Pythium* Species from South Australia. *Mycologia* 57: 417–430.

Van Guilder HD, Vrana KE, Freeman WM (2008) Twenty-five years of quantitative PCR for gene expression analysis. *Biotechniques* 44: 619–626.

Van Niekerk JM, Crous PW, Groenewald JZ, Fourie PH, Halleen F (2004) DNA phylogeny, morphology and pathogenicity of *Botryosphaeria* species on grapevines. *Mycologia* 96: 781–798.

Van Niekerk JM, Groenewald JZ, Farr DF, Fourie PH, Halleen F, Crous PW (2005) Reassessment of *Phomopsis* species on grapevines. *Australasian Plant Pathology* 34: 27–39.

Varela CP, Fernández VR, Casal OA, Vázquez JPM (2011) First report of cankers and dieback caused by *Neofusicoccum mediterraneum* and *Diplodia corticola* on grapevine in Spain. *Plant Disease* 95: 1,315.2.

Varela LG, Smith RJ, Cooper ML, Hoenisch RW (2010) European grapevine moth, *Lobesia botrana*, in Practical winery and vineyard. *Napa Valley Vineyards*, Mar/April: 1–5.

Varga K, Kolber M, Martini M (2000) Phytoplasma identification in Hungarian grapevines by two nested-PCR systems. *Extended Abstracts of the 13th Meeting of ICVG, March 12–18, 2000 Adelaide, Australia* 113–115. ICVG, Switzerland.

Vasiliu-Oromulu L, Barbuceanu D, Ion S (2009) The ecological study of thrips populations in a southern Romanian vineyard (Insecta: Thysanoptera). *Acta Entomologica Serbica* 14: 1–11.

Vasilyeva LN, Stephenson SL (2006) Pyrenomycetes of the Great Smoky Mountains National Park. III. *Cryptosphaeria*, *Eutypa* and *Eutypella* (Diatrypaceae). *Fungal Diversity* 22: 243–254.

Voigt E (1974) Damage to vines in Hungary by the caterpillars of the cabbage moth *Mamestra brassicae*. *Weinberg und Keller* 21: 281–288.

Waite H, Beggs S, Dark H, Kwak P, Murrells G (2005) The effects of different post hot water treatment cool down protocols on dormant cuttings of *Vitis vinifera* cultivars. *Phytopathologia Mediterranea* 44: 119

Walker J (1990) *Verticillium albo-atrum* in Australia: a case study of information confusion in plant pathology. *Australasian Plant Pathology* 19: 57–67.

- Walker K (2006) Brownheaded leafroller (*Ctenopseustis obliquana*) Pest and Diseases Image Library. <http://www.padil.gov.au> Accessed April 2011.
- Waller JM, Bigger M, Hillocks RJ (2007) *Coffee pests, diseases and their management*. CABI, Wallingford, UK.
- Walter B (1997) Effects of viruses on grape and its products. II. *Fanleaf virus* and *nepovirus*. *Progres Agricole et Viticole* 114: 54–58.
- Walter B, Martelli G (1998) Consideration on grapevine selection and certification. *Vitis* 37: 87–90.
- Walton VM, Amy J, Dreves AJ, Gent DH, James DG, Martin RR, Chambers, U Skinkis PA (2007) Relationship between rust mites *Calepitrimerus vitis* (Nalepa), bud mites *Colomerus vitis* (Pagenstecher) [Acari: Eriophyidae] and short shoot syndrome in Oregon vineyards. *International Journal of Acarology* 33: 307–318.
- Walton VM, Daane KM, Pringle KL (2004) Monitoring *Planococcus ficus* in South African vineyards with sex pheromone-baited traps. *Crop Protection* 23: 1089–1096.
- Walton VM, Pringle KL (2004) Vine mealybug, *Planococcus ficus* (Signoret) (Hemiptera: Pseudococcidae), a key pest in South African vineyards – A review. *South African Journal of Enology Viticulture* 25: 54–62.
- Warren LO, Tadic M (1970) The Fall webworm, *Hyphantria cunea* (Drury). Agricultural experiment station, Division of Agriculture; University of Arkansas, Fayetteville, USA.
- Waterhouse DF (1993) *The Major Arthropod Pests and Weeds of Agriculture in Southeast Asia*. ACIAR Monograph No. 21. Australian Centre for International Agricultural Research, Canberra, Australia.
- Waterhouse DF, Sands DPA (2001) *Classical biological control of arthropods in Australia*. Australian Centre for International Agricultural Research, Canberra, Australia.
- Watson GW (2005) Arthropods of economic importance – Diaspididae of the world. In Ulenberg SA (ed) *Arthropods of economic importance*. Natural History Museum, London. <http://nlbif.eti.uva.nl/bis/diaspididae.php> Accessed March 2010.
- Weaver RJ (1976) *Grape growing*. John Wiley and Sons, USA.
- Webb SE (2003) Insect management in grapes. ENY-802, one of a series of the Department of Entomology and Nematology, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. <http://edis.ifas.ufl.edu>. Accessed March 2012.
- Weber E, Golino D, Rowhani A (2002) Laboratory testing for grapevine diseases. *Practical Winery and Vineyard Journal*. <http://www.practicalwinery.com/janfeb02p13.htm> Accessed September 2011.
- Weber E, Golino DA, Rowhani A (1993) Leafroll disease of grapevine. *Practical Winery and Vineyard* 13: 21–25.
- Wei T, Clover G (2008) Use of primers with 5' non-complementary sequences in RT-PCR for the detection of nepovirus subgroups A and B. *Journal of Virological Methods* 153: 16–21.
- Weinert MP, Shivas RG, Pitkethley RN, Daly AM (2003) First record of grapevine leaf rust in the Northern Territory, Australia. *Australasian Plant Pathology* 32: 117–118.

- Weintraub PG, Jones P (2010) *Phytoplasmas: genomes, plant hosts and vectors*. CAB International, Wallingford, UK
- Weiss MJ, McDonald G (1998) European earwig, *Forficula auricularia* L. (Dermaptera: Forficulidae), as a predator of the Redlegged earth mite, *Halotydeus destructor* (Tucker) (Acarina: Penthaleidae). *Australian Journal of Entomology* 37: 183–185.
- Werner R, Mühlbach HP, Büttner C (1997) Detection of *Cherry leafroll nepovirus* (CLRV) in birch, beech and petunia by immuno-capture RT-PCR using a conserved primer pair. *European Journal of Forest Pathology* 27: 309–318
- Wetzel T, Ebel R, Moury B, Le Gall O, Endisch S, Reustle GM, Krczal G (2006). Sequence analysis of grapevine isolates of *Raspberry ringspot nepovirus*. *Archives of Virology* 151: 599–606.
- Wetzel T, Jardak R, Meunier L, Ghorbel A, Reustle GM, Krczal G (2002) Simultaneous RT/PCR detection and differentiation of *Arabis mosaic* and *Grapevine fanleaf nepoviruses* in grapevines with a single pair of primers. *Journal of Virological Methods* 101: 63–69.
- Wetzel T, Krczal G (2007) Molecular biology of *Raspberry ringspot nepovirus*. *Plant Viruses* 1: 45–51.
- Whalley AJS, Watling R (1980) *Daldinia concentrica* versus *Daldinia vernicosa*. *Transactions of the British Mycological Society* 74: 399–406.
- Wheeler AG, Stimmel JF (1988) Heteroptera overwintering in magnolia leaf litter in Pennsylvania. *Entomological News* 99: 65–71.
- White CL (2010) *The characterization of the basidiomycetes and other fungi associated with esca of grapevines in South Africa*. MSc Thesis, Stellenbosch University.
- White IM, Elson-Harris MM (1992) *Fruit flies of economic significance: their identification and bionomics*. CAB International, Wallingford, Oxon, UK and the Australian Center for Agricultural Research, Canberra, Australia.
- Whitelaw-Weckert MA (2008) Interactions between *Cylindrocarpon macrodidymum*, *Streptomyces* spp. MW555 and *Vitis vinifera*. International Conference on Biotic Plant Interactions, Brisbane 27–29th March. University of Queensland, Brisbane.
- Whitelaw-Weckert MA, Curtin S, Huang R, Steel CC, Blanchard CL and Roffey P (2007a) Phylogenetic relationships and pathogenicity of grape isolates of *Colletotrichum acutatum* in sub-tropical Australia. *Plant Pathology* 56: 448–463.
- Whitelaw-Weckert MA, Nair NG, Lamont R, Alonso M, Priest MJ, Huang R (2007b) Root infection of *Vitis vinifera* by *Cylindrocarpon liriodendri* in Australia. *Australasian Plant Pathology* 36: 403–406.
- Whitelaw-Weckert MA, Whitelaw ES, Rogiers SY, Quirk L, Clark AC, Huang CX (2011) Bacterial inflorescence rot of grapevine caused by *Pseudomonas syringae* pv. *syringae*. *Plant Pathology* 60: 325–357.
- Wicks T, Hall B (1990) Evaluation of phosphonic (phosphorous) acid to control *Phytophthora cambivora* on almond and cherry in South Australia. *Australasian Plant Pathology* 19: 132–133.

- Wilcox WF (2003) Grapes: Black rot (*Guignardia bidwellii* (Ellis) Viala and Ravaz). [http://www.nysipm.cornell.edu/factsheets/grapes/diseases/grape\\_br.pdf](http://www.nysipm.cornell.edu/factsheets/grapes/diseases/grape_br.pdf) Accessed December 2011.
- Willems A, Gillis M, Kersters K, Broecke L van den, Ley Jde (1987) The taxonomic position of *Xanthomonas ampelina*. *Bulletin OEPP* 17: 237–240.
- Williams DJ (1982) The distribution of the mealybug genus *Planococcus* (Hemiptera: Pseudococcidae) in Melanesia, Polynesia and Kiribati. *Bulletin of Entomological Research* 72: 441–455.
- Williams DJ (1985) *Australian mealybugs*. British Museum (Natural History), London.
- Williams RN, Fickle DS, Welty C, Ellis M (2011) Insects and mites of grapes in Ohio and Midwest. Department of Entomology, Ohio State University. <http://www.oardc.ohio-state.edu/grapeipm/index.htm> Accessed February 2011.
- Winkler AJ, Cook JA, Kliewer WM, Lider AL (1974) *General Viticulture*. Berkely and Los Angeles, California. University of California Press Ltd, London, England.
- Wong DH, Barbetti MJ, Sivasithamparam (1985) Fungi associated with root rot of subterranean clover in Western Australia. *Australian Journal of Experimental Agriculture* 25: 574–579.
- Woodham RC, Krake LR (1984) Grapevine vein necrosis disease detected in rootstocks in Australia. *Journal of the Australian Institute of Agricultural Science* 50: 58–60.
- Woodham RC, Krake LR, Cellier, KM (1983) The effect of Grapevine leafroll plus Yellow speckle disease on annual growth, yield and quality of grapes from Cabernet Franc under two pruning systems. *Vitis* 22: 324–330.
- Wright LC, James DG, Reyna V, Del Conte SC, Gingras S, Landolt P, Brooks T (2010) Species composition of cutworm (Lepidoptera: Noctuidae) larvae in south central Washington vineyards. *Annals of the Entomological Society of America* 103: 592–596.
- WTO (World Trade Organisation) (1995) Agreement on the application of sanitary and phytosanitary measures. World Trade Organisation, Geneva. [http://www.wto.org/english/tratop\\_e/sps\\_e/spsagr\\_e.htm](http://www.wto.org/english/tratop_e/sps_e/spsagr_e.htm) Accessed July 2012.
- Wunderlich N, Steel CC, Ash G, Raman H, Savocchia S (2008) Identification of *Botryosphaeria* spp. and first report of *Dothiorella viticola* (*Botryosphaeria viticola*) associated with bunch rot in Australia. In *Proceedings of the 6th International Workshop on Grapevine Trunk Diseases, Florence, Italy, 1–3 September*. Firenze University Press, Italy.
- Yang X, Zhou J, Wang F, Cui M (1995) A study on the feeding habits of the larvae of two species of longicorn (Anoplophora) to different tree species. *Journal of Northwestern Forestry College* 10: 1–6 (in Chinese) (Abstract only).
- Yoshikawa N, Iida H, Goto S, Magome H, Takahashi T, Terai Y (1997) *Grapevine berry inner necrosis*, a new trichovirus: comparative studies with several known trichoviruses. *Archives of Virology* 142: 1351–1363.
- Yuan ZQ (1996) Fungi and associated tree diseases in Melville Island, Northern Territory, Australia. *Australian Systematic Botany* 9: 337–360.

- Zalom FG, Bolda MP, Phillips PA (2008) Insects and mites (Leaf rollers). In *UC IPM: Pest management guidelines: Strawberry*. University of California.  
<http://www.ipm.ucdavis.edu/PMG/r734500311.html> Accessed March 2012.
- Zhang L, French R and Langenberg WG (1993) Molecular cloning and sequencing of the coat protein gene of a Nebraskan isolate of Tobacco necrosis virus: the deduced coat protein sequence has only moderate homology with those of strain A and strain D. *Archives of Virology* 132: 291–305.
- Zhang Y, Singh K, Kaur R, Qiu W (2011) Association of a novel DNA virus with the grapevine vein-clearing and vine decline syndrome. *Phytopathology* 101: 1081–1090.
- Zhang YP (2005) *Identification and Control of Pests and Diseases of Table Grape*. Jindun Press, Beijing, China.
- Zhao Y, Wei W, Lee IM, Shao J, Suo X, Davis RE (2009) Construction of an interactive online phytoplasma classification tool, iClassifier, and its application in analysis of peach X-disease phytoplasma group (16SrIII). *International Journal of Systemic and Evolutionary Microbiology* 59: 2572–2593.
- Zhou ZL (1991) Studies on the behaviour and control measures of grape clear-wing moth. *Acta Phytomycol Sinica* 18: 45–48.
- Zhu LZ, Li Y, Li JF, Chen ZZ (2006) Prevention and control of grape spike-stalk brown spot (*Alternaria viticola*) in the southeast areas of Shandong province. *Sino-overseas Grapes and Grapevine* 1: 1 (abstract only).
- Zimmermann D, Bass P, Legin R, Walter B (1990) Characterization and serological detection of four closterovirus-like particles associated with grapevine leafroll disease. *Journal of Phytopathology* 130: 205–218.
- Zitikaite I, Staniulis J (2009) Isolation and characterization of *Tobacco necrosis virus* detected on some vegetable species. *Biologia* 55: 35–39.
- Zorloni A, Prati S, Bianco PA and Belli G (2006) Transmission of *Grapevine virus A* and *Grapevine leafroll-associated virus 3* by *Heliococcus bohemicus*. *Journal of Plant Pathology* 88: 325–328.
- Zorloni A, Quaglino F, Mori N, Zanini G, Bianco PA (2011) ‘Bois noir’ Phytoplasma can be transmitted to healthy *Vitis vinifera* L. plants by rootstocks. *Bulletin of Insectology* 64: S185–S186.
- Zunnoon-Khan S, Arocha-Rosete Y, Scott J, Crosby W, Bertaccini A, Michelutti R (2010) First report of ‘*Candidatus* Phytoplasma fraxini’ (group 16SrVII phytoplasma) associated with peach disease in Canada. *Plant Pathology* 59: 1162.
- Zwadlo T (2012) Central Virginia regional report. *Grape Press* 28: 2.