

The 2021 Antarctic Ozone Hole

Report #16, Wednesday 8 December 2021

Paul Krummel and Paul Fraser Climate Science Centre, CSIRO Oceans and Atmosphere Aspendale, Victoria



Australian Government

Department of Agriculture, Water and the Environment

Contents

1	Summary		2
	1.1	August	2
	1.2	September	2
	1.3	October	4
	1.4	November	5
	1.5	December	7
2	The 2021 ozone hole – metrics		8
	2.1	Ozone hole area	8
	2.2	Ozone deficit	9
	2.3	Ozone hole minima	10
	2.4	Average ozone amount in the hole	11
	2.5	Ozone hole metric summary and rankings	13
-	Total column ozone images		
3	l otal co	blumn ozone images	14
3	NASA M	Numn ozone images NERRA heat flux and temperature	
3	NASA M 4.1	May, June, July, August	14 17 17
3	NASA M 4.1 4.2	Mumn ozone images MERRA heat flux and temperature May, June, July, August September	14 17 17 17
3	1 otal co NASA M 4.1 4.2 4.3	Aleran ozone images Aleran heat flux and temperature May, June, July, August September October	
3	1 otal co NASA M 4.1 4.2 4.3 4.4	Mumn ozone images MERRA heat flux and temperature May, June, July, August September October November	
3	1 otal co NASA M 4.1 4.2 4.3 4.4 4.5	May, June, July, August September October November December	
3 4 5	Iotal co NASA M 4.1 4.2 4.3 4.4 4.5 Satellite	Alexandrication	
3 4 5	Total co NASA M 4.1 4.2 4.3 4.4 4.5 Satellite 5.1	Alexandre images May, June, July, August September October November December Instrumentation OMPS	
3 4 5 6	Iotal coNASA M4.14.24.34.44.5Satellite5.1Archive	Alerra heat flux and temperature May, June, July, August September October November December e Instrumentation OMPS of the weekly reports	
3 4 5 6 7	1 otal co NASA M 4.1 4.2 4.3 4.4 4.5 Satellite 5.1 Archive Definitio	Alerra heat flux and temperature May, June, July, August September October November December e Instrumentation OMPS of the weekly reports	

1 Summary

For the 2021 ozone hole we will be reporting images and metrics calculated from the OMPS data products (see the instrumentation section for a description of these). Please note the OMPS data used in this report are the Level 3 data created from Version 2 (V2.1) of the Nadir Mapper (NM) dataset from the Suomi National Polar-orbiting Partnership satellite.

1.1 August

The first excursions below 220 DU of the ozone minima occurred briefly on a couple of occasions in July. Since 5 August, the ozone minima has remained below 220 DU, dropping to about 190 DU on 5 & 6 August before returning to about average levels for this time of year. From 7 to 23 August the ozone minima approximately followed the mean levels for this time of year, ending at 187 DU on 23 August. The third week of August saw a rapid increase in the ozone hole area reaching 6.6 million km² on 15 August, and by 21 August it had reached 10.2 million km² before dropping back to 8.7 million km² on 23 August. By 23 August the estimated daily ozone deficit had reached a modest 2.4 million tonnes. From the total column ozone images, small patches of ozone depletion can be seen forming in the fringes of the Antarctic polar night (which still covers most of Antarctica at this time of year) from 8-14 August. From 15-23 August, the ozone hole can be seen forming in earnest around the polar night terminator. The strong ridge of high ozone that is normally present in the band immediately south of Australia between about 40-60°S is weaker and appears more-patchy in nature in 2021 compared to previous years. From early-June to the end of July, there was a stratospheric cooling event which saw the heat flux at both the 50 & 100 hPa levels move into record high levels for part of Jun and most of July, which resulted in the 60-90°S zonal mean temperatures at both 50 & 100 hPa being at or close to record low levels for much of June and July. August saw the heat flux at both the 50 & 100 hPa levels return to close to the 70th percentile mark of the 1979-2020 range with the 60-90°S zonal mean temperatures at both 50 & 100 hPa moving to be at about the 10th percentile mark, indicating relatively cold temperatures in the mid to lower stratosphere for this time of year. The forecast is suggesting that these conditions will persist for the coming week.

During the last week of August the 60-90°S zonal mean temperatures at both 50 & 100 hPa continued at about the 10th percentile mark, remaining relatively cold for this time of year. The forecast, however, is suggesting that there will be a small warming event in the coming week. The fourth week of August saw the ozone hole area continue to increase, reaching 13.7 million km² by 29 August, the daily ozone deficit increased quite rapidly reaching 7.4 million tonnes, and the daily ozone minima dropped rapidly reaching 146 DU (which is in the lowest 10th percentile of the 1979-2020 range for this time of year). The total column images from 24-29 August clearly show the 2021 ozone hole continuing to form around the polar night with quite a deep pool of ozone depletion to the south of Australia and New Zealand at around -75°S. The coming week should see the ozone hole become fully formed.

1.2 September

During the first week of September the daily ozone hole area increased rapidly reaching 20.4 million km² on 3 September, before dropping back to 16.8 million km² by 5 September. The variability in the ozone hole area at this time of year is due to variations in the ozone hole formation around the polar night and whether the 220 DU contour that defines the ozone hole is fully closed or not. The daily ozone deficit reached 9.3 million tonnes by 5 September while the daily ozone minima showed some variability but ended at 144 DU by 5 September. The total column ozone images from 30 August to 5 September show the ozone hole forming in earnest, with the hole briefly becoming fully formed on 2 & 3 September, with just a small break in the hole

on 4 & 5 September. The coming week will see the ozone hole become fully formed. The small warming event that was forecast did occur during the first week of September whereby the 45-75°S heat flux at both 50 & 100 hPa returned to close to the mean for this time of year. The 60-90°S zonal mean temperatures at both 50 & 100 hPa exhibited a corresponding increase to be in the lower 30th-50th percentile of 1979-2020 range before dropping back down into the 10th-30th percentile range by the end of the first week of September.

The total column ozone images for 6 through to 12 September show that the 2021 Antarctic ozone hole became fully formed on 8 September with the Australian and New Zealand Antarctic stations all within, or on the edge of the ozone hole from 8-12 September. The ozone hole/polar vortex appears to be very stable with the images from 8-12 September showing a very symmetrical ozone hole. The meteorology during the second week of September supports this, with the relatively cold stratospheric conditions seen so far in 2021 continuing, and along with other metrics is indicating potentially another super polar vortex (strong and stable) in 2021 (pers. comm. Harry Hendon and colleagues). The second week of September saw the daily ozone hole area increase from 15.6 million km² on 6 September to 22.7 million km² by 12 September, with the daily ozone deficit increasing rapidly to be at 17.3 million tonnes by 12 September. The daily ozone minima remained relatively constant during this period in the range of 138 – 147 DU, ending at 146 DU on 12 September.

During the third week of September the daily ozone hole area remained relatively constant at approximately 23 million km², reaching 23.4 million km² on 15 September and ending at 23.2 million km² on 19 September. This time of year is when the peak daily ozone hole area is usually seen, so the peak ozone hole area for 2021 will likely be known in the next 1-2 weeks. For the other metrics, the third week of September saw the daily ozone deficit at 23.7 million tonnes by 19 September, and the daily ozone minima drop sharply reaching 112 DU by 18 September (at the lowest 10th percentile mark of the 1979-2020 range), before rising to be at 119 DU on 19 September. The total column ozone images from 13 to 19 September continue to show a very stable polar vortex with the images from 17-19 September showing a small distortion in the ozone hole/polar vortex. The Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 13-19 September, while Casey station was on the edge of, or just outside of the ozone hole during this period. By the end of third week of September the 45-75°S heat flux at both 50 & 100 hPa was at the 90th percentile mark of the 1979-2020 range, with the forecast suggesting that the heat flux at both levels will move into the upper 10th percentile in the coming week. The 60-90°S zonal mean temperature at 50 hPa entered and remained in the lowest 10th percentile band of the 1979-2020 range during the third week of September, with the forecast predicting that in the coming week it will be at record low levels for this time of year. The 100 hPa 60-90°S zonal mean temperature remained in the 10th-30th percentile range during the third week of September.

The meteorological conditions continue to indicate a very strong and stable polar vortex again this year, with the 45-75°S heat flux at both 50 & 100 hPa in the upper 10th percentile of the 1979-2020 range (unusually week) in the fourth week of September, and the 100 hPa trace at, or near, record levels for this time of year. The 60-90°S zonal mean temperature at 50 hPa was at, or near, record low levels during the fourth week of September, while the 100 hPa 60-90°S zonal mean temperature entered the lowest 10th percentile band. The forecast is for these conditions to continue in the coming week. The strong and stable vortex is shown in the total column ozone images as a very symmetrical ozone hole during 20-26 September. Once again, the Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 20-26 September, while Casey station was predominantly on the edge of, or just outside of the ozone hole during this period. The fourth week of September saw the daily ozone hole area peak at 24.5 million km² on 25 September, which is the largest to date this season and close to the peak ozone hole area in 2020, before dropping back to 24.0 million km² on 26 September. The daily ozone deficit increased considerably to be at 30.7 million tonnes by 26 September, while the daily ozone hole minima dropped to 106 DU on 25 September, before increasing marginally to 108 DU on 26 September.

1.3 October

The end of September and beginning of October saw the daily ozone hole area reach a peak of 24.7 million km² on 28 September which is almost identical to the peak value reached in 2020 and may be the peak value for 2021. Similarly, the daily ozone minima dropped to 95 DU on 29 & 30 September, which is very close to the levels seen in 2020 (94 DU). The daily ozone deficit reached 34.4 million tonnes on 2 October which is similar to the peak level reached in 2018 and somewhat less than the levels reached in 2020, at least to date. The total column ozone images from 27 September to 3 October show a very symmetrical ozone hole from 27 September to 1 October, with a small disturbance affecting the ozone hole symmetry on 2 & 3 October. The Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 27 September to 3 October, while Casey station was predominantly outside of the ozone hole during this period. The end of September and beginning of October saw the 45-75°S heat flux at both 50 & 100 hPa remain in the upper 10th percentile of the 1979-2020 range, which is unusually weak and indicates much less heat transport towards the pole than normal. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during this period. These conditions continue to indicate a very strong and stable polar vortex again this year, with the forecast suggesting these conditions will persist in the coming week.

The daily minima reached significantly low levels during 3-10 October, dropping to 92 DU on 7 & 8 October, before rising sharply to end at 113 DU on 10 October. The low of 92 DU is now less than the 2020 level, and is 6th lowest ozone minima on record. This period saw the daily ozone hole area peak again at 24.8 million km² on 7 October, just larger than the 2020 ozone hole, with the trend in this metric for 2021 displaying signs of being persistent, with the value of 24.5 million km² seen on 10 October now being in the upper 10th percentile for this time of year. The daily ozone deficit reached 35.8 million tonnes on 7 October, which is the highest so far in 2021 and just less than the peak value seen in 2020. Overall the 2021 ozone hole is large, and very deep. A small disturbance continued to affect the ozone hole during 4-9 October with the ozone hole displaying an elongation that is seen in the total column ozone images. However, by 10 October the ozone hole was once again quite symmetrical. The Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 4-10 October, while Casey station was predominantly outside of the ozone hole during this period except for 10 October when it was within the ozone hole. During September and October, the ridge of high ozone to the south of Australia, although still somewhat patchy, has strengthened, with Macquarie Island at times under quite high total ozone columns. The second week of October saw similar conditions to the previous week prevail, with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range, continuing the unusually weak heat flux which indicates much less heat transport towards the pole than normal. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during this period. These conditions continue to indicate a strong and stable polar vortex again this year, with the forecast suggesting these conditions will persist in the coming week.

The total column ozone images from 11 to 17 October show a stable polar vortex with the ozone hole being very symmetrical but partially displaced off of the pole towards South America. This partial displacement saw Casey station outside of the ozone hole for the whole 11-17 October period, while Mawson, Davis and Arrival Heights were all within, or on the edge of the hole. By 17 October the daily ozone hole area had declined to 21.5 million km² and the daily ozone deficit declined to be at 28.5 million tonnes. The daily ozone minima dropped again to 102 DU on 12 & 13 October before rising to 108 DU on 17 October. The third week of October saw similar meteorological conditions to the previous week prevail with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range, but with a small dip in this metric a both levels indicating slightly more heat transport towards the pole. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the third week of October. However, the forecast is suggesting that the heat flux at both levels will be at, or above, record

levels in the coming week, while the 50 hPa zonal mean temperature is forecast to be at record low levels again at the end of the coming week, and the 100 hPa trace to be close to record low levels.

Following a similar pattern to previous weeks, the fourth week of October saw similar meteorological conditions prevail with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range and close or at record high levels for this time of year, indicating much less heat transport towards the pole than normal. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the fourth week of October, with the forecast suggesting that the temperatures at both levels will be at or close to record low levels again at the end of the coming week. The 18-24 October period saw the ozone hole metrics remain relatively stable with the daily ozone hole area at about 21.7 million km² and the daily ozone deficit around 28 million tonnes. The daily ozone minima was in the range of 107-111 DU, with the value of 109 DU on 24 October being a record low for that day of year. For the 18-24 October period the total column ozone images continue to show a stable vortex and ozone hole, with a small elongation to it. Mawson, Davis and Arrival Heights were all within or on the edge of the hole during this period while Casey was outside of the hole except for 24 October when it was on the edge of the hole.

The ozone hole metrics are indicating a persistent ozone hole again this year, which is consistent with the meteorological conditions. The last week of October saw the daily ozone minima reach record low levels for this time of year on 25, 28-31 October, and ended the week with a value of 114 DU on 31 October. The daily ozone hole area and daily ozone deficit both declined during the last week of October reaching 20.1 million km² and 24.3 million tonnes respectively on 31 October. However, as of the end of October, the levels for both of these metrics are now in the upper 10th percentile of the 1979-2020 range for this time of year. The polar vortex continued to be stable during this period with the total column ozone images showing a relatively symmetrical ozone hole. The Antarctic stations of Mawson, Davis and Casey all spent time in, on the edge or out of the ozone hole in the last week of October, while Arrival heights was within the hole for the entire week. Continuing a similar pattern to previous weeks, the last week of October saw similar meteorological conditions prevail with much less heat transport towards the pole than normal and the mid to low stratospheric temperatures in the lowest 10th percentile band. The forecast is suggesting similar conditions will prevail in the coming week and the zonal mean temperatures at both 50 & 100 hPa likely to be at record low levels in the coming week.

1.4 November

The similarities in the ozone hole metrics between the 2020 and 2021 ozone holes is quite striking, in terms of the size/depth and persistence, and following similar temporal patterns. The first week of November saw some fluctuations in the daily ozone hole area, with it ending at 20.5 million km² on 7 November, which is still in the upper 10th percentile of the 1979-2020 range for this time of year, and is very similar in magnitude to the 2020 ozone hole. Although showing a gradual decrease, the daily ozone deficit remained in the upper 10th percentile of the 1979-2020 range, ending at 22.6 million tonnes on 7 November. The record low daily ozone minima levels continued into the beginning of November (1-3), with the levels on 4-7 November just below record levels. By 7 November the daily minima had reached 122 DU. The total column ozone images from 1 to 7 November show a relatively symmetrical ozone hole for 1-3 November, however an elongation of the ozone hole began on 4 November and is quite evident by 7 November along an axis of $60^{\circ}E - 120^{\circ}W$. Mawson, Davis, Casey and Arrival Heights were all within or on the edge of the hole during 1-3 November, however during 4-7 November, Mawson, Davis & Casey all spent some time outside of the ozone hole, especially 5 November when all 3 stations were outside of the hole. Continuing a similar pattern to previous weeks, the first week of November saw similar meteorological conditions prevail with much less heat transport towards the pole than normal and the mid to low stratospheric temperatures in the lowest 10th percentile band. The forecast is suggesting similar conditions will prevail in the coming week, and the zonal mean temperatures at both 50 & 100 hPa will increase considerably but will remain in the lowest 10th percentile band.

The second week of November saw some changes, compared with 2020, in the mid to low stratospheric meteorological conditions. The 45-75°S heat flux at both 50 & 100 hPa remained in the upper 10th percentile of the 1979-2020 range, however these are now starting to deviate from the 2020 trajectory with the forecast suggesting that by the end of the coming week both heat fluxes will be at the 90th percentile mark of the 1979-2020 range. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the second week of November, however there were rapid increases in the temperatures at both levels. The forecast is suggesting that by the end of the coming week the temperatures at both levels will be at the 90th percentile mark of the 1979-2020 range, and considerably warmer than in 2020. This has resulted in a noticeable change in the ozone hole minima and deficit metrics during the second week of November, and these are now displaying relatively large deviations compared to the 2020 ozone hole. There was a rapid increase in the daily ozone minima, reaching 143 DU by 14 November, and a rapid decrease in the daily ozone deficit down to 16.9 million tonnes by 14 November, both of which are now around the 90th percentile mark of the 1979-2020 range for this time of year. The ozone hole area averaged around 19.5 million km² during the second week of November, ending the week at 19.2 million km². Since mid-October the ozone hole area has been slowly declining, but is still following a very similar trajectory to the 2020 ozone hole. In the second week of November a noticeable increase in the ozone levels within the hole can be seen in the total ozone column images and by 14 November the ozone hole was displaced off of the pole towards South America. Arrival Heights was within the ozone hole for the whole second week of November, but likely only partially on 14 November. Mawson and Davis spent time on the edge or completely out of the ozone hole during 7-14 November, while Casey was outside of the ozone hole for the entire period and under a ridge of high ozone (~375 DU) by 14 November.

The third week of November saw the 2021 Antarctic ozone hole become smaller and less deep, but generally remains in the upper 10th percentile of the 1979-2020 range for this time of year for the area and deficit metrics, and in the lower 10th percentile for the minima metric. By 21 November the daily ozone hole area had reduced to 16.8 million km² and the daily ozone deficit dropped steadily to 13.4 million. The ozone hole minima remained around 143-145 DU for much of the third week of November before dropping briefly to 139 DU and then rising sharply to 153 DU by 21 November. The total column ozone images show that the ozone hole was elongated on 15-18 November before becoming more symmetrical during 19-21 November, and has been predominantly offset from the pole towards the Atlantic Ocean. This has resulted in the Australian & New Zealand Antarctic stations spending considerable time outside of the ozone hole during the third week of November. During the third week of November the 45-75°S heat flux at both 50 & 100 hPa remained in the upper 10th percentile of the 1979-2020 range, however the forecast is suggesting that in the coming week the heat flux will move into the 70-90th percentile of the 1979-2020 range, indicating slightly more heat transport towards the pole. The 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the second week of November, however both levels ended the week at the 90th percentile mark of the 1979-2020 range. The forecast suggesting that the temperatures at both levels will likely track 90th percentile mark of the 1979-2020 range in the coming week.

During the fourth week of November the daily ozone hole area plummeted from 16.8 million km² on 21 November down to 10.9 million km² by 28 November. Like the ozone hole area, the daily ozone deficit also dropped rapidly during the fourth week of November, down to 6.5 million tonnes by 28 November, while the daily ozone minima continued to steadily increase, following close to the lower 10th percentile line of the 1979-2020 range, and reaching 164 DU by 28 November. The total column ozone images show the ozone hole becoming quite distorted during 22-25 November, with a tongue of low ozone reaching Heard Island in the Southern Indian Ocean on 24 November (almost reaching Kerguelen Island), and Mawson & Davis being within the ozone hole while Casey and Arrival Heights were outside of the hole for this period. During 26-28 November the ozone hole became more symmetrical again but offset from the pole towards the Indian Ocean, resulting in Mawson, Davis & Casey stations being within the ozone hole during this period while Arrival Heights was outside of the hole. The fourth week of November saw the 45-75°S heat flux at both 50 & 100 hPa move into the 70-90th percentile of the 1979-2020 range, however, the forecast is suggesting that the heat flux will again become less negative in the coming week indicating less heat transport towards the pole. The 60-90°S zonal mean temperatures at both 50 & 100 hPa tracked close to the lower 10th percentile of the 1979-2020 range during the fourth week of November, with the forecast suggesting that they will both be close to or at record low levels again be the end of the coming week.

1.5 December

The end of November/beginning of December saw the ozone hole metrics stabilise. The daily ozone hole area increased marginally from a low of 10.7 million km² on 27 November to 12.5 million km² on 29 November before stabilising to be around 11.3 million km² during 2-5 December. The daily estimated ozone deficit increasing slightly to be around 7 million tonnes by the end of the week and the daily ozone minima went from 167 DU on 29 November down to 158 DU by 5 December, which is a record low (by 1 DU!) for that day of year. The total column ozone images from 29 November to 5 December show the ozone hole to still be predominantly offset from the pole towards the Indian Ocean, but by 5 December had become somewhat elongated along and axis of 60°E – 120°W. Mawson and Davis spend the entire week within the ozone hole, while Casey and Arrival heights were often on the edge of or just within/outside of the hole during this period. The first week of December saw the 45-75°S heat flux at 50 hPa remain in the 70-90th percentile of the 1979-2020 range, while at 100 hPa it moved just into the upper 10th percentile range. The forecast is suggesting that the heat flux will become more negative in the coming week indicating more heat transport towards the pole. The 60-90°S zonal mean temperatures at both 50 & 100 hPa moved well and truly into the lower 10th percentile of the 1979-2020 range during the first week of December and were close to record low levels for this time of year. The forecast is suggesting that the 50 hPa trace will be close to record low levels again by the end of the coming week, while the 100 hPa trace is indicating an increase to be in the middle of the lower 10th percentile band.

2 The 2021 ozone hole – metrics

2.1 Ozone hole area

Figure 1 shows that during July there were some minor ozone depletion events, however, the first two weeks of August is when the ozone hole started to form in the fringes around the polar night, reaching relatively small areas of about 2.5 million km². The third week of August saw a rapid increase in the ozone hole area reaching 6.6 million km² on 15 August, and by 21 August it had reached 10.2 million km² before dropping back to 8.7 million km² on 23 August. The fourth week of August saw the ozone hole area continue to increase, reaching 13.7 million km² by 29 August.

During the first week of September the daily ozone hole area increased rapidly, typical for this time of year, reaching 20.4 million km² on 3 September, before dropping back to 16.8 million km² by 5 September. The variability in the ozone hole area at this time of year is due to variations in the ozone hole formation around the polar night and whether the 220 DU contour that defines the ozone hole is fully closed or not. The second week of September saw the daily ozone hole area increase from 15.6 million km² on 6 September to 22.7 million km² by 12 September, close to the 70th percentile mark of the 1979-2020 range for this time of year. During the third week of September the daily ozone hole area remained relatively constant at approximately 23 million km², reaching 23.4 million km² on 15 September and ending at 23.2 million km² on 19 September. This time of year is when the peak daily ozone hole area is usually seen, so the peak ozone hole area for 2021 will likely be known in the next 1-2 weeks. The fourth week of September saw the daily ozone hole area peak at 24.5 million km² on 25 September, which is the largest to date this season and close to the peak ozone hole area in 2020, before dropping back to 24.0 million km² on 26 September.

The end of September and beginning of October saw the daily ozone hole area reach 24.7 million km² on 28 September before reducing marginally to be at 24.0 million km² by 3 October. The 28 September value of 24.7 million km² is almost identical to the peak value reached in 2020, and may be the peak value for 2021. The period of 4-10 October saw the daily ozone hole area peak again at 24.8 million km² on 7 October, which is now just larger than the 2020 ozone hole. The daily ozone hole area for 2021 is displaying signs of being persistent with the value of 24.5 million km² seen on 10 October now being in the upper 10th percentile for this time of year. By 17 October the daily ozone hole area had declined to 21.5 million km², indicating that the peak of 24.8 million km² seen on 7 October will most likely be the highest daily area for the 2021 ozone hole season. During 18-24 October the daily ozone hole area remained relatively stable at about 21.7 million km², with a jump to 22.4 million km² on 22 October and ending at 21.3 million km² on 24 October. The last week of October saw the daily ozone hole area peak at 21.7 million km² on 26 October before reducing to 20.1 million km² on 31 October. The 31 October level is now in the upper 10th percentile of the 1979-2020 range for this time of year.

The first week of November saw some fluctuations in the daily ozone hole area, peaking at 21.7 million km² on 3 November, then down to 19.6 million km² on 5 November and ending at 20.5 million km² on 7 November. The daily ozone hole area remains in the upper 10th percentile of the 1979-2020 range for this time of year, and is very similar in magnitude to the 2020 ozone hole. The second week of November saw the Antarctic ozone hole area average around 19.5 million km², ending the week at 19.2 million km² on 14 November. Since mid-October the ozone hole area has been slowly declining, but following a very similar trajectory to the 2020 ozone hole which set record high area levels from mid-November through to the end of December. During the third week of November the daily ozone hole area declined more rapidly, dropping to 16.8 million km² by 21 November. However, this is still in the upper 10th percentile of the 1979-2020 range for this time of year. During the fourth week of November the daily ozone hole area plummeted from 16.8 million km² on 21 November. However, this is still in the upper 10th percentile of the 1979-2020 range for this time of year. During the fourth week of November the daily ozone hole area plummeted from 16.8 million km² on 21 November down to 10.9 million km² by 28 November.

The end of November/beginning of December saw the daily ozone hole area increase marginally from a low of 10.7 million km² on 27 November to 12.5 million km² on 29 November before stabilising to be around 11.3 million km² during 2-5 December.



Figure 1 Ozone hole area based on OMPS satellite data (data up to 5 December 2021).

2.2 Ozone deficit

Figure 2 shows that it wasn't until the third week of August that there were substantial levels ozone depletion, with the estimated daily ozone deficit reaching 2.4 million tonnes during 21-23 August. During the fourth week of August, the daily ozone deficit increased quite rapidly, reaching 7.4 million tonnes by 29 August.

The first week of September saw a modest increase in the daily ozone deficit, reaching 9.3 million tonnes by 5 September, similar to the long-term average for this time of year. During the second week of September the daily ozone deficit increased rapidly to be at 17.3 million tonnes by 12 September. The third week of September saw a brief plateau in the daily ozone deficit at 21.5 million tonnes before increasing again to be at 23.7 million tonnes by 19 September. During the fourth week of September the daily ozone deficit increased considerably to be at 30.7 million tonnes by 26 September.

The daily ozone deficit reached 34.4 million tonnes on 2 October, before dropping back to 32.8 million tonnes on 3 October. This is similar to the peak level reached in 2018 and slightly less than the levels reached in 2020. Given that the meteorology in 2021 is similar to that seen in 2020 (persistent strong vortex), this could result in quite a large integrated ozone loss again this year. On 7 October the daily ozone deficit reached 35.8 million tonnes, which is the highest so far in 2021 and just less than the peak value seen in 2020. By 10 October it had dropped back to 33.9 million tonnes. The daily ozone deficit remained around 34 million tonnes during 11-13 October before declining again to be at 28.5 million tonnes on 17 October. The period of 18-24 October saw the daily ozone deficit remain around 28 million tonnes, with a small deviation to 29.1 million tonnes on 22 October before dropping back to 27.3 million tonnes by 24 October. During the last week of October the daily ozone deficit declined from 27.2 million tonnes on 25 October to 24.3 million tonnes on 31 October, however, the 31 October level is now in the upper 10th percentile of the 1979-2020 range for this time of year.

Although showing a gradual decrease, the daily ozone deficit remained in the upper 10th percentile of the 1979-2020 range during the first week of November, ending at 22.6 million tonnes on 7 November. During the second week of November the daily ozone deficit dropped considerably, down to 16.9 million tonnes by 14 November, which is now at the 90th percentile mark of the 1979-2020 range for this time of year. The daily ozone deficit continued to drop steadily during the third week of November, down to 13.4 million tonnes by 21 November, but remains in the upper 10th percentile of the 1979-2020 range for this time of year. Like the ozone hole area, the daily ozone deficit also dropped rapidly during the fourth week of November, down to 6.5 million tonnes by 28 November.

The week of 29 November to 5 December saw the daily estimated ozone deficit increasing slightly to be around 7 million tonnes by the end of the week.



Figure 2 Estimated daily ozone deficit based on OMPS satellite data (2021 data date range as in Figure 1).

2.3 Ozone hole minima

The first excursions below 220 DU of the ozone minima (Figure 3) occurred briefly on 4-8 July to about 210-215 DU around the fringes of the polar night, and again on 26 & 28 July. Since 5 August, the ozone minima remained below 220 DU, dropping to about 190 DU on 5 & 6 August before returning to about average levels for this time of year. From 7 to 23 August the ozone minima approximately followed the mean levels for this time of year, ending at 187 DU on 23 August. This metric can be highly variable at this time of year, but the variability is expected to reduce in the next two to three weeks as the polar night reduces, and the ozone hole fully forms. There was a rapid drop in the daily ozone minima during the fourth week of August, reaching 146 DU on 29 August, which is in the lowest 10th percentile of the 1979-2020 range for this time of year.

Similar to the ozone hole area, the daily ozone minima during the first week of September showed some large variability, rising from the low of 146 DU on 29 August, to 170 DU on 3 September, then down to 144 DU by 5 September. During 6-12 September, the daily ozone minima remained relatively constant in the range of 138 – 147 DU, ending at 146 DU on 12 September. The third week of September saw the daily ozone minima drop sharply again reaching 112 DU by 18 September, which is at the lowest 10th percentile mark of the 1979-2020 range, before rising to be at 119 DU on 19 September. The daily ozone hole minima dropped to 106 DU on 25 September, before increasing marginally to 108 DU on 26 September.

The daily ozone minima continued to drop at the end of September, reaching 95 DU on 29 & 30 September before rising then falling again to 96 DU on 3 October. Again, these levels are very close to those seen in 2020. The daily minima reached significantly low levels during 3-10 October, dropping to 92 DU on 7 & 8 October, before rising sharply to end at 113 DU on 10 October. The low of 92 DU is now less than the 2020 level, and is 6th lowest ozone minima on record. The daily ozone minima dropped again to 102 DU on 12 & 13 October before rising again to 108 DU on 17 October. The daily ozone minima remained relatively stable during 18-24 October in the range of 107-111 DU, with the value of 109 DU on 24 October being a record low for that day of year. The last week of October saw the daily ozone minima reach record low levels for this time of year on 25, 28-31 October, and ended the week with a value of 114 DU on 31 October.

The record low daily ozone minima levels continued into the beginning of November (1-3), with the levels on 4-7 November just below record levels. By 7 November the daily minima had reached 122 DU. There was a rapid increase in the daily ozone minima during the second week of November, reaching 143 DU by 14 November, which is around the 90th percentile mark of the 1979-2020 range for this time of year. Up until this point, the 2021 ozone hole had been closely tracking the trajectory of the 2020 ozone hole, but there is now a relatively large deviation between the two years in this metric. For much of the third week of November the daily ozone hole minima remained around 143-145 DU before dropping briefly to 139 DU and then rising sharply to 153 DU by 21 November. The fourth week of November saw the daily ozone minima continue to steadily increase, following close to the lower 10th percentile line of the 1979-2020 range, and reaching 164 DU by 28 November.

The daily ozone minima went from 167 DU on 29 November down to 158 DU by 5 December, which is a record low (by 1 DU!) for that day of year.



Figure 3 Ozone hole depth based on OMPS satellite data (2021 data date range as in Figure 1).

2.4 Average ozone amount in the hole

The average ozone amount in the hole (averaged column ozone amount in the hole weighted by area; Figure 4) shows a similar pattern to that of the ozone hole minima. The average ozone amount dropped to about 200 DU on 5 & 6 August, before returning to near average levels. On 23 August the average ozone amount in the hole was 207 DU. By 29 August, the average amount of ozone in the whole had dropped relatively rapidly to be at 195 DU, which is in the 10th-30th percentile band of the 1979-2020 range.

By 5 September, the average amount of ozone within the hole had dropped to 194 DU. During the second week of September the average ozone amount in the hole continued to decline, reaching 185 DU by 12 September. The steady decline in the average amount of ozone within the hole continued during the third week of September, reaching 172 DU by 19 September. The fourth week of September saw the average amount of ozone within the hole continue to decline, reaching 160 DU by 26 September.

The beginning of October saw the average amount of ozone within the hole drop to 154 DU, before rising to be at 156 DU on 3 October. The average amount of ozone within the hole dropped to a low during 4-10 October of about 152 DU on 6 October, before increasing marginally to 155 DU on 10 October. This may be the lowest level for 2021, and is about 3-4 DU higher than that seen in 2020. The period of 11-17 October saw a general small increase in the average amount of ozone within the hole, end at 158 DU on 17 October. During 18-24 October the daily average amount of ozone within the hole was stable at 159-160 DU, which is close to record low levels for this time of year. By the end of October the daily average amount of ozone within the hole had increased to approximately 164 DU.

The average amount of ozone within the hole continued a gradual increase during the first week of November, ending at approximately 169 DU on 7 November. Similar to the ozone hole minima, the average amount of ozone within the hole increased considerably during the second week of November, reaching 179 DU by 14 November, which is at the 90th percentile mark of the 1979-2020 range for this time of year. The third week of November saw the average amount of ozone within the hole track close to the lowest 10th percentile mark of the 1979-202 range, ending at 183 DU on 21 November. During the fourth week of November the average amount of ozone within the hole increased considerably by about 10 DU, ending at 192 DU by 28 November.

The end of November/beginning fo December saw the average amount of ozone within the hole go from 195 DU on 29 November down to 191 DU on 5 December, which is very similar again to the 2020 ozone hole for this time of year.



Figure 4 Average ozone amount within the ozone hole based on OMPS satellite data (2021 data date range as in Figure 1).

2.5 Ozone hole metric summary and rankings

Of the 42 holes for which we have data since 1979, for some metrics the 2021 hole is one of the deepest/largest on record. In terms of the <u>largest</u> ozone holes, it is provisionally ranked:

- 13th: 15-day average area 5th: 15-day average minimum ozone
 - daily maximum area 6th: daily minimum ozone
- 11th: daily ozone deficit 14th: daily min average ozone amount in the hole
- ??: integrated ozone deficit

17th:

Once the ozone hole has recovered for this year, we will then be able to rank the 2021 ozone hole for the integrated ozone deficit metric. It is currently ranked 10^{th} highest, and potentially could get as high as 6^{th} or 7^{th} highest.

3 Total column ozone images

The most recent 16 days of total column ozone 'images' over Australia and Antarctica from OMPS are shown in Figure 5.

From the total column ozone images, small patches of ozone depletion can be seen forming in the fringes of the Antarctic polar night (which still covers most of Antarctica at this time of year) from 8-14 August. From 15 August until 23 August, the ozone hole can be seen forming in earnest around the polar night terminator. The strong ridge of high ozone that is normally present in the band immediately south of Australia between about 40-60°S is weaker and appears more-patchy in nature in 2021 compared to previous years. The total column images from 24-29 August clearly show the 2021 ozone hole continuing to form around the polar night with quite a deep pool of ozone depletion to the south of Australia and New Zealand at around -75°S. The coming week should see the ozone hole become fully formed.

The total column ozone images from 30 August to 5 September show the ozone hole forming in earnest, with the hole briefly becoming fully formed on 2 & 3 September, with just a small break in the hole on 4 & 5 September. The coming week will see the ozone hole become fully formed. There is now also a ridge of high ozone to the south of Australia around 50-60°S. The images for 6 through to 12 September show that the 2021 Antarctic ozone hole became fully formed on 8 September with the Australian and New Zealand Antarctic stations all within, or on the edge of the ozone hole from 8-12 September. The ozone hole/polar vortex appears to be very stable with the images from 8-12 September showing a very symmetrical ozone hole. The images from 13 to 19 September continue to show a very stable polar vortex with the images from 17-19 September showing a small distortion in the ozone hole/polar vortex. The Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 13-19 September, while Casey station was on the edge of, or just outside of the ozone hole during this period. The ozone hole/polar vortex continued to be very stable during 20-26 September with the total column ozone images showing a very symmetrical ozone hole. Once again, the Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during this period. The ozone hole/polar vortex continued to be very stable during 20-26 September, while Casey station was predominantly on the edge of, or just outside of the ozone for Mawson, Davis and Arrival Heights were all within the ozone hole during this period. The ozone hole/polar vortex continued to be very stable during 20-26 September, while Casey station was predominantly on the edge of, or just outside of the ozone hole during this period.

The total column ozone images from 27 September to 3 October show a very symmetrical ozone hole from 27 September to 1 October, with a small disturbance affecting the ozone hole symmetry on 2 & 3 October. The Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 27 September to 3 October, while Casey station was predominantly outside of the ozone hole during this period. The small disturbance continued to affect the ozone hole during 4-9 October with the ozone hole displaying an elongation that is seen in the total column ozone images. However, by 10 October the ozone hole was once again quite symmetrical. The Antarctic stations of Mawson, Davis and Arrival Heights were all within the ozone hole during 4-10 October, while Casey station was predominantly outside of the ozone hole during this period except for 10 October when it was within the ozone hole. During September and October, the ridge of high ozone to the south of Australia, although still somewhat patchy, has strengthened, with Macquarie Island at times under quite high total ozone columns. The total column ozone images from 11 to 17 October show a stable polar vortex with the ozone hole being very symmetrical but partially displaced off of the pole towards South America. This partial displacement saw Casey station outside of the ozone hole for the whole 11-17 October period, while Mawson, Davis and Arrival Heights were all within or on the edge of the hole. For the 18-24 October period the total column ozone images continue to show a stable vortex and ozone hole, with a small elongation to it. Mawson, Davis and Arrival Heights were all within or on the edge of the hole during this period while Casey was outside of the hole except for 24 October when it was on the edge of the hole. During the last week of October, the polar vortex continued to be stable with the total column ozone images showing a relatively symmetrical ozone hole. The Antarctic stations of Mawson, Davis and Casey all spent time in, on the edge or out of the ozone hole in the last week of October, while Arrival heights was within the hole for the entire week.

The total column ozone images from 1 to 7 November show a relatively symmetrical ozone hole for 1-3 November, however an elongation of the ozone hole began on 4 November and is quite evident by 7 November along an axis of 60°E – 120°W. Mawson, Davis, Casey and Arrival Heights were all within or on the edge of the hole during 1-3 November, however during 4-7 November, Mawson, Davis & Casey all spent some time outside of the ozone hole, especially 5 November when all 3 stations were outside of the hole. In the second week of November a noticeable increase in the ozone levels within the hole can be seen in the total ozone column images and by 14 November the ozone hole was displaced off of the pole towards South America. Arrival Heights was within the ozone hole for the whole second week of November, but likely only partially on 14 November. Mawson and Davis spent time on the edge or completely out of the ozone hole during 7-14 November, while Casey was outside of the ozone hole for the entire period and under a ridge of high ozone (~375 DU) by 14 November. The total column ozone images show that the ozone hole was elongated on 15-18 November before becoming more symmetrical during 19-21 November, and has been predominantly offset from the pole towards the Atlantic Ocean. This has resulted in the Australian & New Zealand Antarctic stations spending considerable time outside of the ozone hole during the third week of November. The total column ozone images show the ozone hole becoming quite distorted during 22-25 November, with a tongue of low ozone reaching Heard Island in the Southern Indian Ocean on 24 November (and almost reaching Kerguelen Island), and Mawson & Davis being within the ozone hole while Casey and Arrival Heights were outside of the hole for this period. During 26-28 November the ozone hole became more symmetrical again but offset from the pole towards the Indian Ocean, resulting in Mawson, Davis & Casey stations being within the ozone hole during this period while Arrival heights was outside of the hole.

The total column ozone images from 29 November to 5 December show the ozone hole to still be predominantly offset from the pole towards the Indian Ocean, but by 5 December had become somewhat elongated along and axis of $60^{\circ}E - 120^{\circ}W$. Mawson and Davis spend the entire week within the ozone hole, while Casey and Arrival heights were often on the edge of or just within/outside of the hole during this period.



Figure 5 OMPS ozone hole images for 20 November – 5 December; the ozone hole boundary is indicated by the red 220 DU contour line. Green plus symbols indicate the location of the Australian Antarctic stations of Mawson (62.9°E, 67.6°S), Davis (78.0°E, 68.6°S) and Casey (110.5°E, 66.3°S), the New Zealand Antarctic station of Arrival Heights (166.7°E, 77.8°S) and Macquarie Island station (158.9°E, 54.5°S).

4 NASA MERRA heat flux and temperature

The MERRA 45-day mean 45-75°S heat fluxes at 50 & 100 hPa are shown in the left column of Figure 6. A less negative heat flux usually results in a colder polar vortex, while a more negative heat flux indicates heat transported towards the pole (via meteorological disturbances/waves) and results in a warming of the polar vortex. The corresponding 60-90°S zonal mean temperatures at 50 & 100 hPa are shown in the right column of Figure 6, these usually show an anti-correlation to the heat flux.

At 50 hPa, the type 1 PSC (HNO3.3H2O) formation threshold temperature (195 K) was reached in mid-June. At 100 hPa, the threshold temperature was reached at the end of June.

4.1 May, June, July, August

During May, 45-day mean 45-75°S heat flux at 50 hPa was in initially the 10-30th percentile moving into the 30-50th percentile of the 1979-2020 range towards the end of the month, while at 100 hPa it was predominantly in the 10-30th percentile, indicating more heat transport towards the pole than average. The corresponding 60-90°S zonal mean temperatures at both 50 & 100 hPa were very close to the average during May. From early-June to the end of July, there was a cooling event which saw the heat flux at both the 50 & 100 hPa levels move into record high levels for part of Jun and most of July. This resulted in the 60-90°S zonal mean temperatures at both 50 & 100 hPa levels for much of June and July. August (up to 24 August) saw the heat flux at both the 50 & 100 hPa levels return to close to the 70th percentile mark of the 1979-2020 range. Correspondingly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa moved to be at about the 10th percentile mark, indicating relatively cold temperatures in the mid to lower stratosphere for this time of year. The forecast is suggesting that these conditions will persist for the coming week.

The last week of August saw the 45-75°S heat flux at 50 & 100 hPa remain around the 70th percentile mark of the 1979-2020 range, while the 60-90°S zonal mean temperatures at both 50 & 100 hPa continued at about the 10th percentile mark, remaining relatively cold for this time of year. The forecast, however, is suggesting that there will be a small warming event in the coming week with the 45-75°S heat flux at both 50 & 100 hPa returning to close to the mean for this time of year, with the 60-90°S zonal mean temperatures at both 50 & 100 hPa returning modest increases before cooling slightly again.

4.2 September

The small warming event that was forecast did occur during the first week of September whereby the 45-75°S heat flux at both 50 & 100 hPa returned to close to the mean for this time of year. The 60-90°S zonal mean temperatures at both 50 & 100 hPa exhibited a corresponding increase to be in the lower 30th-50th percentile of 1979-2020 range before dropping back down into the 10th-30th percentile range by the end of the first week of September. The forecast is for similar conditions to continue, with the 50 hPa 60-90°S zonal mean temperature to drop into the lowest 10th percentile in the coming week.

The second week of September saw the 45-75°S heat flux at both 50 & 100 hPa move back into the 70-90th percentile of 1979-2020 range, indicating less heat transport towards the pole, which is consistent with a very stable polar vortex. The 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the 10th-30th percentile range during the second week of September, with the trace at 50 hPa ending at the 10th percentile mark. This continues the very cold stratospheric conditions seen so far in 2021 and along with other metrics is indicating potentially another super polar vortex in 2021 (pers. comm. Harry Hendon and colleagues).

By the end of third week of September the 45-75°S heat flux at both 50 & 100 hPa was at the 90th percentile mark of the 1979-2020 range, with the forecast suggesting that the heat flux at both levels will move into the upper 10th percentile in the coming week, and for the 100 hPa trace to be at record levels for this time of year. The 60-90°S zonal mean temperature at 50 hPa entered and remained in the lowest 10th percentile band of the 1979-2020 range during the third week of September, with the forecast predicting that in the coming week it will be at record levels for this time of year. The 100 hPa 60-90°S zonal mean temperature remained in the 10th-30th percentile range during the third week of September with the forecast suggesting it will enter the lowest 10th percentile band be the end of the fourth week of September.

The fourth week of September saw the 45-75°S heat flux at both 50 & 100 hPa move into the upper 10th percentile of the 1979-2020 range, which is unusually weak, with the 100 hPa trace at, or near, record levels for this time of year. The forecast is suggesting that the heat flux at both levels will remain in the upper 10th percentile in the coming week. The 60-90°S zonal mean temperature at 50 hPa was at, or near, record low levels during the fourth week of September, with the forecast predicting that in the coming week it will remain at, or near, record low levels for this time of year. The 100 hPa 60-90°S zonal mean temperature entered the lowest 10th percentile band during the fourth week of September, with the forecast suggesting it will remain there in the coming week. These conditions continue to indicate a very strong and stable polar vortex again this year.

4.3 October

The end of September and beginning of October saw the 45-75°S heat flux at both 50 & 100 hPa remain in the upper 10th percentile of the 1979-2020 range, which is unusually weak and indicates much less heat transport towards the pole than normal. The forecast is suggesting that the heat flux at both levels will remain in the upper 10th percentile in the coming week. The 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band at the end of September/beginning of October, with the forecast suggesting they will remain there in the coming week. These conditions continue to indicate a very strong and stable polar vortex again this year.

The second week of October saw similar conditions to the previous week prevail with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range, continuing the unusually weak heat flux which indicates much less heat transport towards the pole than normal. The forecast is suggesting that the heat flux at both levels will remain in the upper 10th percentile in the coming week. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the second week of October, with the forecast suggesting they will remain there in the coming week. These conditions continue to indicate a strong and stable polar vortex again this year.

The third week of October saw similar conditions to the previous week prevail with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range, but with a small dip in this metric a both levels indicating slightly more heat transport towards the pole. However, the forecast is suggesting that the heat flux at both levels will be at, or above, record levels in the coming week. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the third week of October, with the forecast suggesting the 50 hPa trace will enter record levels again at the end of the coming week, and the 100 hPa trace to be close to record low levels.

Following a similar pattern to previous weeks, the fourth week of October saw similar conditions prevail with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range and close or at record high levels for this time of year, indicating much less heat transport towards the pole than normal. The forecast is suggesting that the heat flux at both levels will remain in the upper 10th percentile in the coming week. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the fourth week of October, with the forecast suggesting

that the temperatures at both the 50 & 100 hPa levels will be at or close to record low levels again at the end of the coming week.

The last week of October saw similar conditions continue with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range, continuing the trend this season of much less heat transport towards the pole than normal. The forecast is suggesting that the heat flux at both levels will remain in the upper 10th percentile in the coming week. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the last week of October, with the forecast suggesting that the temperatures at both the 50 & 100 hPa removes the temperatures at both the solution of the temperatures at both temperatures at both the solution of the temperatures at both temperatures at both the solution of the temperatures at both t

4.4 November

The first week of November saw similar conditions continue with the 45-75°S heat flux at both 50 & 100 hPa remaining in the upper 10th percentile of the 1979-2020 range, continuing the trend this season of much less heat transport towards the pole than normal. The forecast is suggesting that the heat flux at both levels will remain in the upper 10th percentile in the coming week. Similarly, the 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the first week of November, with the 50 hPa trace being at or close to record low levels for this time of year. The forecast is suggesting that the temperatures at both the 50 & 100 hPa levels will increase considerably in the coming week but will remain in the lowest 10th percentile band.

The second week of November saw the 45-75°S heat flux at both 50 & 100 hPa remain in the upper 10th percentile of the 1979-2020 range, however these are now starting to deviate from the 2020 trajectory with the forecast suggesting that by the end of the coming week the heat flux will be at the 90th percentile mark of the 1979-2020 range. The 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the second week of November, however there were rapid increases in the temperatures at both levels. The forecast suggesting that by the end of the 1979-2020 range, and considerably warmer than in 2020.

During the third week of November the 45-75°S heat flux at both 50 & 100 hPa remained in the upper 10th percentile of the 1979-2020 range, however the forecast is suggesting that in the coming week the heat flux will move into the 70-90th percentile of the 1979-2020 range. The 60-90°S zonal mean temperatures at both 50 & 100 hPa remained in the lowest 10th percentile band during the second week of November, however both levels ended the week at the 10th percentile mark of the 1979-2020 range. The forecast suggesting that the temperatures at both levels will likely track 10th percentile mark of the 1979-2020 range in the coming week.

The fourth week of November saw the 45-75°S heat flux at both 50 & 100 hPa move into the 70-90th percentile of the 1979-2020 range, however, the forecast is suggesting that the heat flux will again become less negative in the coming week indicating less heat transport towards the pole. The 60-90°S zonal mean temperatures at both 50 & 100 hPa tracked close to the lower 10th percentile of the 1979-2020 range during the fourth week of November, with the forecast suggesting that they will both be close to or at record low levels again be the end of the coming week.

4.5 December

The first week of December saw the 45-75°S heat flux at 50 hPa remain in the 70-90th percentile of the 1979-2020 range, while at 100 hPa it moved just into the upper 10th percentile range. The forecast is suggesting that the heat flux will become more negative in the coming week indicating more heat transport towards the

pole. The 60-90°S zonal mean temperatures at both 50 & 100 hPa moved well and truly into the lower 10th percentile of the 1979-2020 range during the first week of December and were close to record low levels for this time of year. The forecast is suggesting that the 50 hPa trace will be close to record low levels again by the end of the coming week, while the 100 hPa trace is indicating an increase to be in the middle of the lower 10th percentile band.



Figure 6 MERRA2 45-day mean 45°S-75°S eddy heat flux and 60°S-90°S zonal mean temperature at 50 & 100 hPa. Images courtesy of NASA GSFC, downloaded 8 December 2021, data through to 15 December 2021 (data starting 2021-10-01 are from GEOS FP; 2021-12-07 onwards are forecasts): http://ozonewatch.gsfc.nasa.gov/meteorology/SH.html

5 Satellite Instrumentation

5.1 OMPS

OMPS (Ozone Mapping and Profiler Suite) is an ozone instrument on the Suomi National Polar-orbiting Partnership satellite (Suomi NPP), which was launched on 28 October 2011 and placed into a sunsynchronous orbit 824 km above the Earth. The partnership is between NASA, NOAA and DoD (Department of Defense), see https://www.jpss.noaa.gov/ for more details. OMPS continues the US program for monitoring the Earth's ozone layer using advanced hyperspectral instruments that measure sunlight in the ultraviolet and visible, backscattered from the Earth's atmosphere, and will contribute to observing the recovery of the ozone layer in coming years. For the 2021 ozone hole season, we will be using the OMPS total column ozone data and will be producing metrics from both OMI and OMPS Level 3 global gridded daily total ozone column products from NASA, and present both sets of results for comparison. The OMPS data used in this report are the Level 3 data created from Version 2.1 (V2.1) of the Nadir Mapper (NM) dataset from Suomi-NPP's Ozone Mapping and Profiler Suite (OMPS).

6 Archive of the weekly reports

The weekly Antarctic Ozone Hole reports for the 2021 ozone hole season are available from the Department of the Environment and Energy web page here:

http://www.environment.gov.au/protection/ozone/publications/antarctic-ozone-hole-summary-reports

7 Definitions

CFCs: chlorofluorocarbons, synthetic chemicals containing chlorine, once used as refrigerants, aerosol propellants and foam-blowing agents, that break down in the stratosphere (15-30 km above the earth's surface), releasing reactive chlorine radicals that catalytically destroy stratospheric ozone.

DU: Dobson Unit, a measure of the total ozone amount in a column of the atmosphere, from the earth's surface to the upper atmosphere, 90% of which resides in the stratosphere at 15 to 30 km.

Halons: synthetic chemicals containing bromine, once used as fire-fighting agents that break down in the stratosphere releasing reactive bromine radicals that catalytically destroy stratospheric ozone. Bromine radicals are about 50 times more effective than chlorine radicals in catalytic ozone destruction.

MERRA: is a NASA reanalysis for the satellite era using a major new version of the Goddard Earth Observing System Data Assimilation System Version 5 (GEOS-5). The project focuses on historical analyses of the hydrological cycle in a broad range of weather and climate time scales. It places modern observing systems (such as EOS suite of observations) in a climate context. Since these data are from a reanalysis, they are not up-to-date. So, NASA supplement with the GEOS-5 FP data that are also produced by the GEOS-5 model in near real time. These products are produced by the NASA Global Modeling and Assimilation Office (GMAO).

MERRA2: MERRA2 was introduced to replace the original MERRA dataset because of the advances made in the assimilation system that enable assimilation of modern hyperspectral radiance and microwave observations, along with GPS-Radio Occultation datasets. It also uses NASA ozone observations after 2005. Additional advances in both the GEOS-5 model and the GSI assimilation system are included in MERRA-2.

Ozone: a reactive form of oxygen with the chemical formula O3; ozone absorbs most of the UV radiation from the sun before it can reach the earth's surface.

Ozone Hole: ozone holes are examples of severe ozone loss brought about by the presence of ozone depleting chlorine and bromine radicals, whose levels are enhanced by the presence of PSCs (polar stratospheric clouds), usually within the Antarctic polar vortex. The chlorine and bromine radicals result from the breakdown of CFCs and halons in the stratosphere. Smaller ozone holes have been observed within the weaker Arctic polar vortex.

Polar night terminator: the delimiter between the polar night (continual darkness during winter over the Antarctic) and the encroaching sunlight. By the first week of October the polar night has ended at the South Pole.

Polar vortex: a region of the polar stratosphere isolated from the rest of the stratosphere by high west-east wind jets centred at about 60°S that develop during the polar night. The isolation from the rest of the atmosphere and the absence of solar radiation results in very low temperatures (< -78°C) inside the vortex.

PSCs: polar stratospheric clouds are formed when the temperatures in the stratosphere drop below -78°C, usually inside the polar vortex. This causes the low levels of water vapour present to freeze, forming ice crystals and usually incorporates nitrate or sulphate anions.

TOMS, OMI & OMPS: the Total Ozone Mapping Spectrometer (TOMS), Ozone Monitoring Instrument (OMI), and Ozone Mapping and Profiler Suite (OMPS) are satellite borne instruments that measure the amount of back-scattered solar UV radiation absorbed by ozone in the atmosphere; the amount of UV absorbed is proportional to the amount of ozone present in the atmosphere.

UV radiation: a component of the solar radiation spectrum with wavelengths shorter than those of visible light; most solar UV radiation is absorbed by ozone in the stratosphere; some UV radiation reaches the earth's surface, in particular UV-B which has been implicated in serious health effects for humans and animals; the wavelength range of UV-B is 280-315 nanometres.

8 Acknowledgements

The TOMS and OMI data are provided by the TOMS ozone processing team, NASA Goddard Space Flight Center, Atmosp heric Chemistry & Dynamics Branch, Code 613.3. The OMI instrument was developed and built by the Netherlands's Agency for Aerospace Programs (NIVR) in collaboration with the Finnish Meteorological Institute (FMI) and NASA. The OMI science team is led by the Royal Netherlands Meteorological Institute (KNMI) and NASA. The OMPS Level 3 data used in this report were created from a research dataset developed by NASA's NPP Ozone Science Team using nadir mapper measurements from Suomi-NPP's Ozone Mapping and Profiler Suite (OMPS). All data were downloaded from either the NASA Ozone Watch webpage https://ozonewatch.gsfc.nasa.gov/ or the NASA Earth Data Near Real Time data portal https://earthdata.nasa.gov/earth-observation-data/near-real-time/download-nrt-data/omps-nrt.

The NASA Ozone Watch webpage contains a lot of useful data and information on the Antarctic ozone hole.

As Australia's national science agency and innovation catalyst, CSIRO is solving the greatest challenges through innovative science and technology.

CSIRO. Unlocking a better future for everyone.

Contact us

1300 363 400 +61 3 9545 2176 csiroenquiries@csiro.au csiro.au

For further information

Climate Science Centre CSIRO Oceans and Atmosphere Paul Krummel +61 3 9239 4568 Paul.Krummel@csiro.au https://www.csiro.au/en/Research/OandA/Areas/Assessingour-climate/Climate-Science-Centre