

Australian National Insect Collection, O'Dwyer & Attiwill 1999, Clarke & O'Dwyer 2000 and Clarke & Whyte 2003 and other reliable sources.

The distribution of areas where the Golden Sun Moth is known to occur is from the ACT Government (2005). Golden Sun Moth (Synemon plana) habitat mapping, Environment ACT. Along with a 1km buffer of all post 1990 records.

The distribution of areas where the Golden Sun Moth is likely to occur is derived from a 3km buffer of pre 1990 records

The distribution of areas where the Golden Sun Moth may occur is derived from the NSW Ecosystems Database. The following ecosystem areas were mapped based on species occurrence; Boorowa Volcanics, Dalton Hills, Marilba Range, Burrinjuck Ridges, Doura Volcanics, Sato Alannels and Floodplain, Canberra Plains, Upper Murrumbidgee Channels and Floodplain and Minjary Hills and Ranges. Excluding the western part of the Murrumbidgee area an north east part of Tumut area as was very steep and mountainous.

The TOPO1M information in this product is copyright © Commonwealth of Australia, Geoscience Australia, 2002.

Datum: GDA 1994

02.55 10 15 20

The information presented in this map has been provided by a range of groups and agencies. While every effort has been made to ensure accuracy and completeness, no guarantee is given, nor responsibility taken by the Commonwealth for errors or omissions, and the Commonwealth does not accept responsibility in respect of any information or advice given in relation to, or as a consequence of, anything containing herein The map has been collated from a range of sources, with data at various resolutions. Data used are assumed to be correct as received from the data suppliers.



Australian Government

Department of the Environment, Water, Heritage and the Arts

Produced by Environmental Resources Information Network

Australian Government
Department of the Environment, Water, Heritage and the Arts

December 2008
COPYRIGHT Commonwealth of Australia, 2008