

Records are from the Species Profile and Threats Database, Atlas of Victorian Wildlife, Australian National Insect Collection, O'Dwyer & Attiwill 1999, Clarke & O'Dwyer 2000 and Clarke & Whyte 2003 and other reliable sources.

The distribution of areas were the Golden Sun Moth is known to occur is derived from a 1 km buffer of post1990 records and includes known sites. The distribution of areas were the Golden Sun Moth is likely to occur is derived from a 3km buffer of pre 1990 records. The distribution of areas were the Golden Sun Moth may occur is derived from the Victorian Pre-European Ecological Vegetation Class (EVC) mapping, DSE, 2008. The following EVC's were mapped;175, 132 and 55. EVCs were clipped to areas with elevation between 100 - 800m.

The TOPO10M information in this product is copyright © Commonwealth of Australia, Geoscience Australia, 2002.


Legend

Golden Sun Moth Records

△ pre 1990

post 1990

Distribution



0 12.5 25 50 75

Caveat:

The information presented in this map has been provided by a range of groups and agencies. While every effort has been made to ensure accuracy and completeness, no guarantee is given, nor responsibility taken by the Commonwealth for errors or omissions, and the Commonwealth does not accept responsibility in respect of any information or advice given in relation to, or as a consequence of, anything containing herein. The map has been collated from a range of sources, with data at various resolutions. Data used are assumed to be correct as received from the data suppliers.



Australian Government **Department of the Environment,** Water, Heritage and the Arts

Produced by Environmental Resources Information Network Australian Government

Department of the Environment, Water, Heritage and the Arts December 2008

COPYRIGHT Commonwealth of Australia, 2008