



NVIS Fact sheet

MVG 14 – Mallee woodlands and shrublands

Australia's native vegetation is a rich and fundamental element of our natural heritage. It binds and nourishes our ancient soils; shelters and sustains wildlife, protects streams, wetlands, estuaries, and coastlines; and absorbs carbon dioxide while emitting oxygen. The National Vegetation Information System (NVIS) has been developed and maintained by all Australian governments to provide a national picture that captures and explains the broad diversity of our native vegetation.

This is part of a series of fact sheets which the Australian Government developed based on NVIS Version 4.2 data to provide detailed descriptions of the major vegetation groups (MVGs) and other MVG types. The series is comprised of a fact sheet for each of the 25 MVGs to inform their use by planners and policy makers. An additional eight MVGs are available outlining other MVG types.

For more information on these fact sheets, including its limitations and caveats related to its use, please see: 'Introduction to the Major Vegetation Group (MVG) fact sheets'.

Overview

Typically, vegetation areas classified under MVG 14 – Mallee woodlands and shrublands:

- are semi-arid systems dominated by eucalypt species that produce multiple stems from an underground rootstock known as a lignotuber
- have some vegetation in non-arid regions that also supports eucalypts with mallee growth forms, but these are not part of MVG 14 or MVG 32
- have eucalypt as the most widespread tree component, however, species of *Callitris*, *Melaleuca*, *Acacia* and *Hakea* may co-occur as trees at varying densities and may co-dominate mallee communities
- feature an open tree or shrub layer (>10 per cent projective foliage cover, > 20 per cent crown cover) distinguishes MVG 14 from MVG 32 (Mallee open woodlands and Sparse mallee shrublands), which has a sparser tree or shrub layer (notionally <10 per cent foliage projective cover, <20 per cent crown cover). However, this distinction can be interpreted as largely artificial, as tree cover varies spatially at fine scales and temporally with fire regimes, while the occurrence of tree, shrub and ground layer species shows little relationship to variations in eucalypt cover
- comprise understorey composition strongly influenced by rainfall, soil types and fire regime and can be dominated by hummock grasses (spinifex), chenopods or other woody shrubs
- are among the most fire prone of all plant communities in semi-arid and arid zones.

Facts and Figures

Major Vegetation Group	MVG 14 - Mallee woodlands and shrublands
Major Vegetation Subgroups (number of NVIS descriptions)	27. Mallee with hummock grass 29. Mallee with a dense shrubby understorey 55. Mallee with an open shrubby understorey 61. Mallee with a tussock grass understorey
Typical NVIS structural formations	Closed mallee forest (tall, mid, low) Open mallee forest (tall, mid, low) Mallee woodland (tall, mid, low) Closed mallee shrubland (mid, low) Mallee shrubland (mid, low) Open mallee shrubland (mid, low)
Number of IBRA regions	39
Most extensive in IBRA region (Est. pre-1750 and present)	Present: Great Victoria Desert (SA, WA) Pre-1750: Murray Darling Depression (NSW, SA, Vic)
Estimated pre-1750 extent (km²)	318 217
Present extent (km²)	213 763
Area protected (km²)	87 391

Structure and physiognomy

- Woodlands and shrublands dominated by low, multi-stemmed, sclerophyllous eucalypts and occasionally co-dominated by small trees from other genera with a sparse to dense understorey.
- Height of eucalypt canopy rarely exceeds six m.
- Tree canopy cover varies with rainfall, topographic position, soil characteristics and particularly fire history, but projective foliage cover is notionally within the range 10 – 30 per cent and crown cover 20 – 50 per cent.
- Understorey structure also varies with rainfall, topographic position, soil characteristics and particularly fire history, and may be dominated by shrubs, hummock grasses, chenopods or tussock grasses. In drought the ground layer is sparse, while following heavy rainfall a prominent cover of ephemeral herbs with tussock grasses occurs.
- A soil crust forms on flat areas and local depressions, comprising cyanobacteria, lichens and mosses, but is largely absent from dune crests and slopes.
- The subgroups of this MVG differ in terms of canopy dominants, understorey composition, soil characteristics and topographic position.

Indicative flora

- Eucalypts of the semi-arid mallee woodlands belong almost entirely to sections *Bisectaria* and *Dumaria* within the subgenus *Symphyomyrtus*, although a few species belong to sections *Adnataria*, also within *Symphyomyrtus*. See further discussion in Notes below. Widespread mallee tree species include *Eucalyptus dumosa* (white mallee), *E. Socialis* (red mallee), *E. Gracilis* (yorrell), *E. oleosa* (red mallee), and *E. calycogona* (gooseberry mallee).
- Mallee communities in Western Australia are characterised by several eucalypts some of which are endemic to Western Australia, including *E. eremophila*, *E. moderata*, *E. oleosa*, *E. incrassata*, *E. foecunda*, *E. redunca* and *E. uncinata*. The understorey is predominantly shrubby with species of *Melaleuca* and *Acacia*, and occasional *Triodia*. They are widespread in the southern goldfield regions, extending eastward to the coast of the Great Australian Bight.
- Distinctions between the subgroups are based mainly on the perennial components of the understorey; hummock grass (*Triodia scariosa* and *Triodia irritans*), tussock grass (*Themeda*, *Poa* and *Astrelba*) and open or dense shrubby understoreys (including chenopods).

- Typical mallee understoreys include the following:
 - Shrubby understoreys dominated by Chenopods may include species of *Maireana*, *Sclerolaena*, *Enchylaena*, *Chenopodium* and *Zygophyllum*.
 - Other shrubby understoreys include a range of mostly non-sclerophyll shrubs from genera such as *Senna*, *Acacia*, *Dodonaea*, *Myoporum*, *Eremophila*, *Enchylaena*, *Sclerolaena*, *Melaleuca*, *Grevillea*, *Hakea*, and *Westringia*.
 - Some sclerophyllous shrub species may also form shrubby understoreys with *Callitris verrucosa* often prominent in either the tree or shrub layer. Typical shrub genera include *Acacia*, *Aotus*, *Baeckea*, *Leptospermum*, *Melaleuca*, *Beyeria*, *Brachyloma*, *Hibbertia*, *Cryptandra*, *Spyridium* and *Phebalium*. *Triodia* may occur sparsely among the shrubs.
 - *Triodia* (hummock) understoreys are characteristically dominated by *Triodia scariosa* and *Triodia irritans*, interspersed with small trees or shrubs of genera such as *Acacia*, *Callitris*, *Eremophila* and *Beyeria*.
 - Less common are dominant *Tussock* understoreys from the genera *Austrostipa*, *Austrodanthonia*, *Themeda*, *Poa* and *Astrelba*. Tussocks may also be interspersed among other dominant understorey growth forms.
- Ephemeral herbs from a wide range of plant families are abundant after rain including the *Poaceae* (mainly *Austrostipa*), *Asteraceae*, *Goodeniaceae*, *Solanaceae*, *Apiaceae*, *Brassicaceae* and *Zygophyllaceae*.

Environment

- Typically semi-arid Mediterranean with cool winters and hot dry summers, and annual rainfall of between 200 – 500 mm per annum occurring predominately in winter.
- Occurs on calcareous sandplains, predominantly with lateritic soils in Western Australia and on Aeolian landscapes in eastern Australia.
- Different geomorphic processes have shaped these mallee regions with much of the central and south-eastern mallee characterised by sand dune and swale systems.
- In many areas the variation in the thickness of surface sand above clay or calcareous hard pan affects the structural and floristic composition of mallee.
- Soils associated with mallee of arid Australia include solonized brown soils, siliceous sands, calcareous sands, solodized solonetz, soloths, earthy sands and calcareous red earths.



Mallee, near Lake Grace, WA (Photo: M. Fagg)

Geography

- Located in the semi-arid areas of southern Australia in south-west New South Wales, north-west Victoria, southern South Australia and Western Australia.
- Outliers in the Northern Territory, Queensland and northern Western Australia refer mostly to the growth form of pockets of individual trees in places such as mountain tops, rather than the typical extensive mallee plains of southern Australia.
- Largest area occurs in South Australia (95 828 km²).
- In the east, mallee is largely restricted to the sandplains and sand dunes of the western part of the Murray Basin with small areas occurring on landforms of aeolian origin further north.
- In the west, mallee occurs on three main land forms, the Yilgarn Plateau, the Stirling coastal zone and the Nullarbor Plain.
- In South Australia, mallee occurs on extensive dune fields of the Eyre Peninsula, the lower Murray and in the Gulf Ranges provinces.
- Understoreys in the temperate zones in southern Western Australia, South Australia, central-west Victoria and central New South Wales are likely to produce dense shrubby understoreys, becoming more open with increasing latitude and transitioning into hummock grasses in the semi-arid and arid zones in central Western Australia and South Australia, northern


Victoria, and south-western New South Wales. Pockets in the Northern Territory and Queensland are generally dominated by hummock grasses.

- Make up the most arid of the eucalypt-dominated communities of temperate Australia.

The image below outlines the location of this MVG group in Australia.

Change

- Approximately 33 per cent (>104 000 km²) of the estimated pre-1750 extent cleared accounting for 10 per cent of total clearing in Australia including large areas of southern Western Australia and South Australia, and northern Victoria.
- Losses have been disproportionate, and greatest by far in mallee with a shrubby (notably chenopod) understorey.
- Clearing for crops is ongoing, albeit at reduced rates. Legacies of fragmentation continue to degrade ecological functions.
- Removed from many cereal cropping and sheep grazing lands in the southern and eastern parts of Australia over the last 100 years, most recently irrigated vineyards and orchards.
- Clearing of the mallee in South Australia commenced in approximately 1900 following the widespread introduction of the stump-jump plough.



- Most of the South Australian Murraylands, Yorke and Eyre Peninsula areas were cleared before World War II. Tops of sandhills were cleared later in the 1960s as part of the rabbit control program.
- The Kangaroo Island mallee community was extensively cleared following World War II with the soldier settlement scheme.
- Overgrazing associated with pastoral activities and feral herbivores (notably rabbits and goats) have modified mallee in the inland and more arid areas of Australia beyond the rainfall limits of cropping and within the cropping region.
- Fire regimes involving short intervals between successive fires and fire events followed by droughts and/or overgrazing are greatly reducing plant diversity and the capacity of mallee to support its avian and mammal fauna.
- Threats include fragmentation of remaining areas, inappropriate fire regimes (e.g. fires too regular and/or too intense) and grazing impact on understorey.
- Associated issues in understanding and managing mallee areas for multiple values are yet to be fully explored.

Key values

- Many species of plants, birds, reptiles and invertebrates not found in any other ecosystems.
- Uniquely Australian form of semi-arid woodland.
- Remnant populations of many rare and endangered species of vertebrate and invertebrates (e.g. the Black-eared Miner and the Mallee Fowl).
- Significance to Indigenous communities.
- Conservation and tourism.

List of key management issues

- Control of vegetation clearing for mining, cropping and other transient uses.
- Weed control in remnant vegetation within agricultural landscapes.
- Management of total grazing pressure (including domestic, native and feral animals).
- Rising water tables in cropping areas affected by dryland salinity.
- Destruction of soil crusts by non-native hard hooved animals.
- Wind erosion of soils exposed by overgrazing or cropping.
- Fire frequency and interactions between fires and droughts.
- Livestock grazing and feral animal activity in recently burnt areas.
- Declining rainfall, especially in western and southern Australia.
- Many forms of mallee vegetation remain primarily in private ownership or under leasehold.
- Increased stewardship and multi-value management in response to pastoralist and mining access to natural resources.
- Long-term monitoring to inform future management strategies.



Mallee near Murray Bridge, SA (Photo: M. Fagg)

References

- Australian Surveying and Land Information Group (1990) *Atlas of Australian Resources. Volume 6 Vegetation*. AUSMAP, Department of Administrative Services, Canberra, 64pp. & 2 maps.
- Beadle N.C.W. (1981) *The Vegetation of Australia*. Cambridge Univ. Press, Cambridge, 690pp.
- Beard J.S., Beetson, G.R., Harvey J.M., Hopkins A.J.M. and Shepherd D.P. (2013) *The Vegetation of Western Australia at 1:3,000,000 Scale. Explanatory Memoir*. Second Edition. Science Division, Department of Parks and Wildlife, Western Australia.
- Blackburn G. and Wright M.J. (1989) Soils. In: *Mediterranean Landscapes in Australia: Mallee Ecosystems and their Management* (ed. Noble J.C. and Bradstock R.A.) pp. 34 - 53. CSIRO, Canberra.
- Bradstock R.A. and Cohn J.S. (2002) Fire regimes and biodiversity in semi-arid mallee ecosystems. In *Flammable Australia*. The fire regimes and biodiversity of a continent. (ed. Bradstock R.A., Williams J.E. and Gill A.M.) pp. 238 - 258. Cambridge University Press, Cambridge
- Brooker M.I.H. and Kleinig D.A. (1999) *Field guide to Eucalypts, South-eastern Australia, Volume 1, 2nd edition*. Bloomings Books, Melbourne, 353pp.
- Brooker M.I.H. and Kleinig D.A. (2001) *Field guide to Eucalypts, South-western and Southern Australia, Volume 2, 2nd edition*. Bloomings Books, Melbourne, 428pp.
- Haslem A., Callister K.E., Avitabile S.C., Griffioen P.A., Kelly, L.T., Nimmo D.G., Spence-Bailey L.M., Taylor R.S., Watson S.J., Brown L., Bennett A.F. and Clarke M.F. (2010) *A framework for mapping vegetation over broad spatial extents: A technique to aid land management across jurisdictional boundaries*. Landscape and Urban Planning, vol. 97, pp. 296-305.
- Hill K.D. (1989) Mallee Eucalypt communities: their classification and biogeography In: *Mediterranean Landscapes in Australia: Mallee Ecosystems and their Management* (ed. Noble J.C. and Bradstock R.A.) pp. 93 - 108. CSIRO, Canberra.
- Keith D.A. (2004) *Ocean Shores to Desert Dunes. The native vegetation of New South Wales and the ACT*. Department of Environment and Conservation (NSW), Hurstville.
- National Land and Water Resources Audit (2001) *Australian Native Vegetation Assessment 2001*. National Land and Water Resources Audit, Canberra, 332pp.
- Noble J.C. and Bradstock R.A., eds. (1989) *Mediterranean Landscapes in Australia, Mallee ecosystems and their management*. CSIRO, Melbourne, 485pp.
- Parsons R.F. (1994) *Eucalyptus* scrubs and shrublands. In: *Australian Vegetation* (ed. R.H. Groves) pp. 291-319. Cambridge Univ. Press, Cambridge.
- Wardell-Johnson G.W., Williams J.E., Hill K.D. and Cumming R. (1997) Evolutionary biogeography and contemporary distribution of eucalypts. In: *Eucalypt Ecology. Individuals to ecosystems* (eds. J.E. Williams and J.C.Z. Woinarski) pp. 92-128. Cambridge Univ. Press, Cambridge.
- Wasson R.J. (1989) Landforms. In: *Mediterranean Landscapes in Australia: Mallee Ecosystems and their Management* (ed. Noble J.C. and Bradstock R.A.) pp. 13 - 34. CSIRO, Canberra.
- Watson S.J., Taylor R.S., Nimmo D.G., Kelly L.T., Haslem A., Clarke M.F. and Bennett A.F. (2012) Effects of time-since-fire on bird species: how informative are generalized fire-response curves for conservation management? *Ecological Applications* vol. 22, pp. 685-696.

Data sources

Interim Biogeographic Regionalisation for Australia (IBRA),
Version 7.

National Vegetation Information System, Version 4.2.

Collaborative Australian Protected Areas Database –
CAPAD 2014 – Terrestrial.

Notes

- Many authors take a broad interpretation of ‘mallee woodland’ to include any vegetation dominated by multi-stemmed eucalypts. That concept encompasses additional vegetation types including woodlands on rocky hills and other stony substrates in semi-arid landscapes at temperate and subtropical latitudes that incorporate mallee growth forms from sections *Exsertaria* or *Adnataria*. It may also include mallee heaths in temperate humid or semi-humid climates on rocky or sandy substrates with multi-stemmed eucalypts from subgenus *Monocalyptus* or *Eudesmia*. All of these vegetation types are excluded from MVGs 14 and 32, and instead referred to in woodland and heathland groups. The ‘true’ mallee woodlands and shrublands included within MVGs 14 and 32 encompass those in semi-arid, winter-rain, temperate climates on sandy duplex substrates, dominated by eucalypts in sections *Bisectaria* and *Dumaria*.
- This fact sheet should be read in conjunction with MVG 32: Mallee open woodlands and sparse shrublands.

© Commonwealth of Australia, 2017.



This fact sheet is licensed by Commonwealth of Australia under a Creative Commons Attribution 4.0 International licence.

The views and opinions expressed in this publication are those of the authors and do not necessarily reflect those of the Australian Government or the Minister for the Environment and Energy.