




1 - branchlets segmented




Callitris intratropica

4 - leaves without a petiole, stem-clasping base



Calytrix arborescens

4* - leaves with a petiole




5 - *Calytrix achaeta*




5* - *Calytrix exstipulata*

7 - leaves alternate



7* - leaves whorled or opposite



GROUP E
Tree or shrub, no prickles, leaves absent,
or simple and ≤ 2 mm wide

1 - Branchlets segmented [3.3.2d], segments 2-3 mm long, each with a whorl of 4 tiny scale leaves < 1 mm long; mature plant a tree	<i>Callitris intratropica</i> [Cypress Pine]
1* - Branchlets not segmented; shrubs to tall shrubs	2
2 - Leaves absent [3.3.5b]; stems sub-divide into long, smooth, erect, slender, green twigs	<i>Pachynema junceum</i>
2* - Leaves present	3
3 - Most leaves < 3 mm long [3.3.5e]	4
3* - Most leaves > 4 mm long	6
4 - Leaves without petioles [3.3.7]; leaf bases stem-clasping	<i>Calytrix arborescens</i>
4* - Leaves with petioles ; leaf bases not stem-clasping	5
5 - Leaves with minute patent hairs [3.3.16h] on all surfaces; outer branches columnar [3.3.2e], often pendulous [3.3.3b]; flowers white	<i>Calytrix achaeta</i>
5* - Leaves with only a few minute (barely detectable) hairs on the margins; outer branches not columnar, erect spreading; flowers bright pink	<i>Calytrix exstipulata</i> [Turkey Bush]
6 - Leaves < 10 mm long	<i>Calytrix brownii</i> [Kerosene Bush]
6* - Most leaves > 10 mm long	7
7 - Leaves alternate [3.3.8a-d]	8
7* - Leaves opposite [3.3.8e] <i>or</i> whorled [3.3.8g]	9
8 - Leaves and stems smooth, not scaly [3.3.16e]	<i>Acacia gonocarpa</i>
8* - Leaves and stems densely encrusted with tiny circular scales	<i>Hibbertia scabra</i>
9 - Leaves opposite [3.3.8e] <i>or</i> in whorls of 3 [3.3.8g]; most older leaves < 17 mm long	<i>Verticordia cunninghamii</i> [Cunningham's Featherflower]
9* - Leaves in whorls of 4; most older leaves > 17 mm long	<i>Verticordia verticillata</i> [Featherflower]