

## APPENDIX 4

### THE RESPONSES OF NATIVE AUSTRALIAN PLANT SPECIES TO *Phytophthora cinnamomi*

A list of Australian plant responses to *Phytophthora cinnamomi* has been compiled<sup>1</sup> (Table A4.1) from published material and the unpublished records and observations of individual researchers. Comments, corrections and suggested additions should be sent to e.ogara@murdoch.edu.au

#### **'Are there species present for which the impact of *P. cinnamomi* would be significant?'**

The criteria for the level of threat to vegetation at a site, that warrants management of *P. cinnamomi*, is currently impossible to prescribe for nationally. The different States will need to exercise discretion on the criteria, and criteria need to be developed as a first step in planning the management of *P. cinnamomi*.

Table A4.1 contains species that range from highly susceptible to field resistant. In response to the question posed in the decision flowchart (Figure 5.1) 'Are there species present for which the impact of *P. cinnamomi* would be significant', it is suggested that the answer is 'yes' if:

1. there are species present that are listed in the table as moderately (MS) or highly susceptible (HS)
2. there are species and ecological communities present that are threatened and the extent of susceptibility to *P. cinnamomi* is unknown.

#### ***Threatened species and ecological communities***

The Australian Government *Environmental Protection & Biodiversity Conservation Act 1999* (EPBC Act) is national legislation that promotes the conservation of Australia's biodiversity. [Nationally threatened taxa](#), listed in schedules of the EPBC Act, are denoted in Table A4.1. General and spatial information on nationally listed taxa is available from the [Protected Matters Search Tool](#) on the Australian Government DEH website.

Each Australian State and Territory has its own environmental legislation for listing threatened taxa based on State/Territory boundaries. The status of species in individual States and Territories does not appear in Table A4.1; therefore the determination of the potential impact of *P. cinnamomi* at a site should also include consultation of the relevant State/Territory lists. Links to each of the State and Territory sites relating to listing of threatened taxa are provided below:

---

<sup>1</sup> The list was compiled by Dr Keith McDougall, Environment Protection and Regulation Division, Department of Environment and Conservation, PO Box 2115, Queanbeyan NSW 2620. Last updated 12<sup>th</sup> September, 2005.

[Australian Capital Territory](#)

[New South Wales](#)

- Profiles of flora and fauna of NSW are available in a [Wildlife Atlas](#)

[Northern Territory](#)

Queensland

- [Vegetation Management Act 1999 – Vegetation Management Regulation 2000](#)
- [Nature Conservation Act 1992/State Penalties Enforcement Act 1999 – Nature Conservation and other Legislation Amendment Regulation \(No. 1\) 2000](#)

[South Australia](#)

[Tasmania](#)

[Victoria](#)

[Western Australia](#)

### ***Important Caveats on the Use of Tables A4.1***

Table A4.1 is **not solely a host list**. Whilst it does contain the known Australian native hosts of *P. cinnamomi*, the fact that a species can be a host does not mean that it will display symptoms of infection in the wild. The responses of native plants to infection by *P. cinnamomi* are many and various:

- hosts of *P. cinnamomi* in the wild may show no obvious symptoms of infection
- the response of a species in the wild may depend on static site conditions (e.g. substrate and pH) and temporal conditions (e.g. rainfall and disturbances such as fire)
- species may be affected in some situations (e.g. in cultivation or glasshouse experiments) but largely unaffected in others (e.g. in the wild)
- there may be spatial variation in the response (e.g. *Hibbertia hypericoides* is highly susceptible to infection on the Swan Coastal Plain of WA but rarely affected in the adjoining jarrah forest)
- species may not be hosts of *P. cinnamomi* at all, but may be affected nonetheless by changes in habitat caused by the death of surrounding plants.

An effort has been made in Table A4.1 to indicate the field susceptibility of species to infection and spatial variation in susceptibility where they are known. **The list is indicative and not definitive.** We suggest that it is used as an indication of the potential impact of *P. cinnamomi* on native plants and vegetation, and **should not replace careful site evaluation** (e.g. sampling of roots and soil for the presence of the pathogen and long-term monitoring). As the information in Table A4.1 will require some interpretation it is strongly recommended that the following points are noted prior to consulting or using the information:

- the listing of a taxon in Table A4.1 as a host or as a susceptible species in one State or Territory does not necessarily mean that it is a host or is susceptible to infection across its range
- no attempt has been made in the list to evaluate the veracity of susceptibility ratings. Please read the cited reference to make this assessment yourself

- references provided in the list should be sought to clarify issues of variable susceptibility. In addition, the list is a work in progress - many more species will be added in the future, and many of the susceptibility ratings will be altered as we gain a better understanding of the effects of the pathogen over a greater time and over the entire area that it is capable of reaching

## ***Explanatory Notes on Table A4.1***

### **Species nomenclature**

In the list the name given to a taxon is that currently shown as accepted in the [Australian Plant Name Index](#). Taxa that have been split since the referenced work was published or for which no indication was given in a reference of the subspecific rank, are indicated by s.l. (*sensu lato*, in the broadest sense). Nationally threatened taxa, listed in schedules of the EPBC Act, are denoted in the table as **CE** (critically endangered), **E** (endangered) or **V** (vulnerable).

### **Distribution**

The distribution of taxa is indicated by the State or Territory in which they have been recorded: n = New South Wales and the Australian Capital Territory, nt = Northern Territory, q = Queensland, s = South Australia, t = Tasmania, v = Victoria, w = Western Australia; saf = South Africa (plantation species).

### **References**

The numbers in the body of the Table refer to the numbered references in the References section.

### **Isolation**

Species from which *P. cinnamomi* has been isolated have been separated into; those growing in the wild, those grown in cultivation (mostly botanic gardens), and those used in experiments to test for susceptibility. Experimental isolations were generally performed in a glasshouse environment. A few, indicated by a \*, were obtained by inoculating propagated plants in the field.

### **Susceptibility rating**

The susceptibility of a taxon, where known, is indicated by a rating adapted from previously used systems:

- **HS** – highly susceptible, i.e. species that are frequently and consistently killed in the wild following infection by *P. cinnamomi*, and/or appear to decline or be rare on infested sites (includes scale categories 10, 11, and 12 of Wills<sup>40</sup> and Barrett<sup>59</sup>, and groups 3 and 5 of Shearer and Dillon<sup>14</sup>)
- **MS** – moderately susceptible (or variable susceptibility), i.e. species that are often killed following infection by *P. cinnamomi* in the wild but many plants of which commonly survive (includes scale categories 7, 8 and 9 of Wills<sup>40</sup> and Barrett<sup>59</sup>, and group 4 of Shearer and Dillon<sup>14</sup>)
- **LS** – low susceptibility, i.e. species that are rarely but occasionally found dead on infested sites (includes scale categories 4, 5 and 6 of Wills<sup>40</sup> and Barrett<sup>59</sup>, and group 2 of Shearer and Dillon<sup>14</sup>)
- **S** – susceptible and thought to be affected, but degree of susceptibility not documented

- **SP** - susceptible but persistent, i.e. species that are frequently killed following infection by *P. cinnamomi* in the wild but which persist on affected sites through effective reproductive strategies
- **SV** – variable susceptibility; plants may be commonly killed on some infested sites but appear unaffected on others – this may be attributable to genetic differences between populations or differences in site characteristics that influence plant responses
- **FR** - field resistant (or tolerant), i.e. species that appear to be unaffected by *P. cinnamomi* in the wild when it is present and for which deaths in the field can rarely be associated with infection by *P. cinnamomi* (includes scale categories 1, 2 and 3 of Wills<sup>40</sup> and Barrett<sup>59</sup>, and group 1 of Shearer and Dillon<sup>14</sup>)
- **Q** – not known to be directly affected by *P. cinnamomi* but rarely found on affected sites (and may be affected either directly through infection or through changes in habitat).

**Table A4.1** A list of Australian native plants that are potential hosts of *Phytophthora cinnamomi*. Please consult the accompanying text before using this list.

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<b>ADIANTACEAE</b>					
<i>Cheilanthes austrotenuifolia</i>	n,t,s,t,v,w				FR <sup>59</sup>
<b>AGAVACEAE</b>					
<i>Cordyline murchisoniae</i>	q	16			
<b>AMARANTHACEAE</b>					
<i>Ptilotus declinatus</i>					FR <sup>21</sup>
<i>Ptilotus manglesii</i>	w	24			FR <sup>24</sup>
<b>ANNONACEAE</b>					
<i>Goniothalamus australis</i>	q				FR <sup>36</sup>
<b>ANTHERICACEAE</b>					
<i>Borya mirabilis</i> E	v	25	44		HS <sup>44</sup>
<i>Chamaescilla corymbosa</i> var. <i>corymbosa</i>	n,s,t,v,w	25			S <sup>43</sup> ,FR <sup>21</sup>
<i>Laxmannia grandiflora</i> subsp. <i>stirlingensis</i>	w				FR <sup>58</sup>
<i>Laxmannia jamesii</i> V	w				FR <sup>58</sup>
<i>Laxmannia orientalis</i>	s,t,v	25,29			
<i>Laxmannia sessiliflora</i>	n,s,t,v,w				
<i>Laxmannia squarrosa</i>	w				FR <sup>21</sup>
<i>Thysanotus dichotomus</i>	w	24			FR <sup>24</sup>
<i>Thysanotus multiflorus</i>	w			21	Q <sup>21</sup>
<i>Thysanotus thyrsoides</i>	w				Q <sup>21</sup>
<b>APIACEAE</b>					
<i>Actinotus bellidiooides</i>	t,v	17		33	FR <sup>61</sup>
<i>Actinotus helianthi</i>	n,q	1			
<i>Actinotus rhomboideus</i>	w				FR <sup>59</sup>
<i>Hydrocotyle hirta</i>	s,t,v,w	25			
<i>Pentapeltis peltigera</i>	w	24			FR <sup>24</sup> ,Q <sup>21</sup>
<i>Platysace compressa</i>	w	14,15			FR <sup>59</sup> ,S <sup>54</sup> , HS <sup>14,21</sup>
<i>Platysace heterophylla</i> s.l.	s,v	25			
<i>Platysace lanceolata</i> s.l.	n,q,v				FR <sup>22</sup>
<i>Platysace</i> sp. Stirling	w				FR <sup>58,59</sup>
<i>Platysace tenuissima</i>	w	24			FR <sup>24</sup>
<i>Xanthosia atkinsoniana</i>	n,w	24			FR <sup>21,24</sup>
<i>Xanthosia candida</i>	w	24			FR <sup>21,24</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Xanthosia dissecta</i>	n,s,t,v	25,29			
<i>Xanthosia huegelii</i>	w				FR <sup>21</sup>
<i>Xanthosia rotundifolia</i> s.l.	w				FR <sup>40,59</sup>
<i>Xanthosia tridentata</i>	n,t,v	1			
<b>APOCYNACEAE</b>					
<i>Alstonia muelleriana</i>	q				S <sup>36</sup>
<b>AQUIFOLIACEAE</b>					
<i>Sphenostemon lobosporus</i>	q				FR <sup>36</sup>
<b>ARALIACEAE</b>					
<i>Polyscias australiana</i>	nt,q				FR <sup>36</sup>
<i>Polyscias murrayi</i>	n,q,				S <sup>36</sup>
<b>ARAUCARIACEAE</b>					
<i>Wollemia nobilis</i>	n		12	11	
<b>ARECACEAE</b>					
<i>Archontophoenix cunninghamiana</i>	n,q	16			
<i>Oraniopsis appendiculata</i>	q				S <sup>36</sup>
<b>ASTELIACEAE</b>					
<i>Astelia australiana</i>	t,v	30			
<b>ASTERACEAE</b>					
<i>Argentipallium obtusifolium</i>	n,s,t,v,w	25,29			S <sup>43</sup>
<i>Brachyscome uliginosa</i>	s,v	25,29			S <sup>43</sup>
<i>Cassinia aculeata</i>	n,q,s,t,v	2,3,17			
<i>Helichrysum collinum</i>	n,q	3			
<i>Helichrysum macranthum</i>	w				FR <sup>59</sup>
<i>Hyalosperma cotula</i>	s,v,w				FR <sup>21</sup>
<i>Ixodia achillaeoides</i> subsp. <i>alata</i>	s,v	20			
<i>Lagenophora huegelii</i>	s,t,v,w				FR <sup>21</sup>
<i>Millotia tenuifolia</i> s.l.	n,s,t,v,w				FR <sup>21</sup>
<i>Olearia axillaris</i>	n,s,t,v,w				LS <sup>40</sup>
<i>Olearia ciliata</i> s.l.	n,q,s,t,v,w	29			
<i>Olearia pannosa</i> s.l.	s,v	32			MS <sup>32</sup>
<i>Olearia paucidentata</i>	w				FR <sup>21</sup>
<i>Olearia teretifolia</i>	s,v		3		
<i>Olearia</i> sp. aff. <i>axillaris</i> (Stirling Range)	w				FR <sup>59</sup>
<i>Ozothamnus obcordatus</i> s.l.	n,q,t,v		3		
<i>Podolepis gracilis</i>	w				FR <sup>21</sup>
<i>Podotheca angustifolia</i>	n,s,t,v,w				FR <sup>21</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Pterochaeta paniculata</i>	w				FR <sup>21</sup>
<i>Trichocline spathulata</i>	w				Q <sup>21</sup>
<i>Waitzia nitida</i>	w				FR <sup>21</sup>
<b>BALANOPACEAE</b>					
<i>Balanops australiana</i>	q				FR <sup>36</sup>
<b>BLANDFORDIACEAE</b>					
<i>Blandfordia punicea</i>	t	17,34			HS <sup>34</sup>
<b>BLECHNACEAE</b>					
<i>Blechnum wattsii</i>	n,q,s,t,v	17			
<b>BRUNONIACEAE</b>					
<i>Brunonia australis</i>	n,nt,q,s,t,v,w	25			
<b>CAESALPINIACEAE</b>					
<i>Labichea punctata</i>	w			21	Q <sup>21</sup>
<b>CAMPANULACEAE</b>					
<i>Isotoma hypocrateriformis</i>	w				FR <sup>21</sup>
<i>Lobelia gibbosa</i>	n,q,s,t,v,w				FR <sup>40</sup>
<i>Lobelia rhytidosperma</i>	w				FR <sup>21</sup>
<b>CASUARINACEAE</b>					
<i>Allocasuarina acutivalvis</i> s.l.	w	3			
<i>Allocasuarina campestris</i> s.l.	w	3			
<i>Allocasuarina crassa</i>	t			9	MS <sup>9</sup>
<i>Allocasuarina decussata</i>	w				FR <sup>59</sup>
<i>Allocasuarina duncanii</i>	t			9	HS <sup>9</sup>
<i>Allocasuarina eriochlamys</i> subsp. <i>grossa</i>	w				LS <sup>41</sup>
<i>Allocasuarina fibrosa</i> V	w				MS <sup>41</sup>
<i>Allocasuarina fraseriana</i>	w	2,14,15, 24,28			SP <sup>21,46, 51,53</sup> HS <sup>40,59</sup>
<i>Allocasuarina globosa</i>	w				LS <sup>41</sup>
<i>Allocasuarina grevilleoides</i>	w				MS <sup>41</sup>
<i>Allocasuarina helmsii</i>	s,w	3			
<i>Allocasuarina humilis</i>	w	40	3		S <sup>37,46,53</sup> FR <sup>40</sup>
<i>Allocasuarina lehmanniana</i> s.l.	w	40			FR <sup>40</sup>
<i>Allocasuarina littoralis</i>	n,q,t,v				SV <sup>42</sup>
<i>Allocasuarina microstachya</i>	w	40			FR <sup>40</sup>
<i>Allocasuarina monilifera</i>	t,v	17,34			S <sup>60</sup>
<i>Allocasuarina muelleriana</i> s.l.	s,v	25,29			S <sup>43</sup>
<i>Allocasuarina paludosa</i>	n,s,t,v	25			S <sup>43</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Allocasuarina pinaster</i>	w		3		
<i>Allocasuarina pusilla</i>	s,v	25,29			S <sup>43</sup>
<i>Allocasuarina ramosissima</i>	w				MS <sup>41</sup>
<i>Allocasuarina rigida</i> s.l.	n,q		3		
<i>Allocasuarina tessellata</i>	w				MS <sup>41</sup>
<i>Allocasuarina thuyoides</i>	w				S <sup>56</sup> ,MS <sup>40</sup>
<i>Allocasuarina tortiramula</i>	w				MS <sup>41</sup>
<i>Allocasuarina torulosa</i>	n,q	2			
<i>Allocasuarina trichodon</i>	w		3		FR <sup>59</sup>
<i>Allocasuarina verticillata</i>	n,s,t,v	2,8			FR <sup>60</sup>
<i>Casuarina cunninghamiana</i> subsp. <i>cunninghamiana</i>	n,q	2			
<i>Casuarina obesa</i>	n,s,v,w	50			FR <sup>50</sup>
<b>CENTROLEPIDACEAE</b>					
<i>Centrolepis aristata</i>	s,t,v,w				FR <sup>21</sup>
<b>CLUSIACEAE</b>					
<i>Garcinia</i> sp. (Davies Ck JG Tracey 14745)	q				FR <sup>36</sup>
<b>COLCHICACEAE</b>					
<i>Burchardia multiflora</i>	w	40			FR <sup>40</sup>
<i>Burchardia umbellata</i>	n,q,s,t,v,w	40			FR <sup>40</sup>
<b>CUNONIACEAE</b>					
<i>Anodopetalum biglandulosum</i>	t	17			S <sup>60</sup>
<i>Bauera rubioides</i>	n,q,s,t,v	8,17,34		33	LS <sup>61</sup>
<i>Bauera sessiliflora</i>	v		44		LS <sup>44</sup>
<i>Geissois biagiana</i>	q				FR <sup>36</sup>
<i>Gillbeea adenopetala</i>	q				S <sup>36</sup>
<i>Gillbeea whypalliana</i>	q				FR <sup>36</sup>
<b>CUPRESSACEAE</b>					
<i>Actinostrobus pyramidalis</i>	w	2a			S <sup>54</sup>
<i>Callitris preissii</i>	w		3		
<i>Callitris rhomboidea</i>	n,q,s,t,v	29			FR <sup>60</sup>
<b>CYPERACEAE</b>					
<i>Caustis dioica</i>	w	20			FR <sup>40</sup>
<i>Cyathochaeta avenacea</i>	w				FR <sup>21,40</sup>
<i>Cyathochaeta clandestina</i>	w	24			FR <sup>24</sup>
<i>Evandra aristata</i>	w				S <sup>46</sup> ,LS <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Gahnia grandis</i>	n,t,v	17			FR <sup>60</sup>
<i>Gahnia trifida</i>	s,t,v,w				FR <sup>40</sup>
<i>Gymnoschoenus sphaerocephalus</i>	n,s,t,v	17			FR <sup>61</sup>
<i>Lepidosperma angustatum</i>	w				FR <sup>21</sup>
<i>Lepidosperma brunonianum</i>	w				FR <sup>59</sup>
<i>Lepidosperma concavum</i>	s,t,v	17			FR <sup>42</sup>
<i>Lepidosperma effusum</i>	w				FR <sup>59</sup>
<i>Lepidosperma gladiatum</i>	n,s,t,v,w				FR <sup>59</sup>
<i>Lepidosperma laterale</i> s.l.	n,q,s,t,v	2			FR <sup>60</sup>
<i>Lepidosperma longitudinale</i>	n,q,s,t,v,w				FR <sup>58</sup>
<i>Lepidosperma scabrum</i>	w	24			FR <sup>24</sup>
<i>Lepidosperma squamatum</i>	w				FR <sup>21,59</sup>
<i>Lepidosperma tenue</i>	w	24			FR <sup>24</sup>
<i>Lepidosperma tetraquetrum</i>	w				FR <sup>53</sup>
<i>Lepidosperma urophorum</i>	n,q,v				FR <sup>42</sup>
<i>Lepidosperma viscidum</i>	s,v,w				FR <sup>40</sup>
<i>Mesomelaena graciliceps</i>	w				Q <sup>21</sup>
<i>Mesomelaena stygia</i> s.l.	w				FR <sup>40</sup>
<i>Mesomelaena tetragona</i>	w				FR <sup>21,24</sup>
<i>Ptilothrix deusta</i>	n,q				FR <sup>42</sup>
<i>Schoenus efoliatus</i>	w				FR <sup>58,59</sup>
<i>Schoenus imberbis</i>	n,v				FR <sup>42</sup>
<i>Schoenus</i> sp. Stirling	w				FR <sup>58,59</sup>
<i>Tetraria capillaris</i>	n,q,s,t,v,w				FR <sup>21,40</sup>
<i>Tetraria octandra</i>	w				FR <sup>21</sup>
<b>DASYPOGONACEAE</b>					
<i>Chamaexeros serra</i>	w	40			FR <sup>40</sup>
<i>Dasypogon bromeliifolius</i>	w	24			S <sup>37,46</sup> ,LS <sup>40,59</sup>
<i>Kingia australis</i>	w				FR <sup>22</sup>
<b>DENNSTAEDTIACEAE</b>					
<i>Pteridium esculentum</i>	n,q,s,t,v,w	2			FR <sup>40,59</sup> ,Q <sup>21</sup>
<b>DILLENIACEAE</b>					
<i>Hibbertia acerosa</i>	w	48,15,28			S <sup>48</sup>
<i>Hibbertia acicularis</i>	n,q,s,t,v	17,34			
<i>Hibbertia amplexicaulis</i>	w	14,28			S <sup>21,24,48</sup> ,HS <sup>14</sup>
<i>Hibbertia australis</i>	s,v	25			
<i>Hibbertia calycina</i>	n,t,v			9	HS <sup>9</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Hibbertia commutata</i>	w	28		21	LS <sup>40,59</sup> ,HS <sup>21</sup>
<i>Hibbertia cunninghamii</i>	w	15			
<i>Hibbertia desmophylla</i>	w	51			S <sup>51</sup>
<i>Hibbertia empetrifolia</i>	n,t,v	17,34			
<i>Hibbertia furfuracea</i>	w				S <sup>56</sup>
<i>Hibbertia glomerata</i> s.l.	w				HS <sup>21</sup>
<i>Hibbertia huegelii</i>	w			21	Q <sup>21</sup>
<i>Hibbertia humifusa</i>					FR <sup>44</sup>
<i>Hibbertia hypericoides</i>	w	14,24			S <sup>37,46</sup> ,SP <sup>21</sup> MS <sup>40</sup> ,HS <sup>14</sup>
<i>Hibbertia inconspicua</i>	w				S <sup>46</sup>
<i>Hibbertia lineata</i>	w	14,24			S <sup>24</sup>
<i>Hibbertia montana</i>	w	14,48,24			S <sup>24,48</sup> ,HS <sup>14</sup>
<i>Hibbertia obtusifolia</i>	n,q,t,v	3			
<i>Hibbertia procumbens</i>	n,t,v	17,34			S <sup>60</sup>
<i>Hibbertia prostrata</i>	s,t,v	17,25,29, 34			S <sup>43</sup>
<i>Hibbertia quadricolor</i>	w	14			S <sup>24</sup> ,HS <sup>14,21</sup>
<i>Hibbertia rhadinopoda</i>	w	26		21	SP <sup>21</sup>
<i>Hibbertia riparia</i>	n,q,s,t,v	17,20,25, 29,34			S <sup>60</sup>
<i>Hibbertia sericea</i> s.l.	n,q,s,v	17,34			
<i>Hibbertia silvestris</i>	w	14,24			FR <sup>24</sup>
<i>Hibbertia stricta</i>	s,v	4,7,25			S <sup>43</sup>
<i>Hibbertia subvaginata</i>	w	14,15,24, 37			HS <sup>14</sup>
<i>Hibbertia virgata</i>	n,s,t,v	5,25		9	S <sup>43</sup> , HS <sup>9</sup>
<b>DROSERACEAE</b>					
<i>Drosera erythrorhiza</i> s.l.	w				FR <sup>21,40</sup>
<i>Drosera huegelii</i>	w				FR <sup>59</sup>
<i>Drosera macrantha</i> s.l.	w				FR <sup>21</sup>
<i>Drosera pallida</i>	w				FR <sup>40</sup>
<i>Drosera platystigma</i>	w				FR <sup>21</sup>
<i>Drosera stolonifera</i> s.l.	w				FR <sup>59</sup> ,Q <sup>21</sup>
<b>ELAECARPACEAE</b>					
<i>Elaeocarpus eumundi</i>	n,q				FR <sup>36</sup>
<i>Elaeocarpus foveolatus</i>	q				S <sup>36</sup>
<i>Elaeocarpus holopetalus</i>	n,v				MS <sup>22</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Elaeocarpus largiflorens</i> subsp. <i>retinervis</i>	q				FR <sup>36</sup>
<i>Elaeocarpus sericopetalus</i>	q				S <sup>36</sup>
<i>Sloanea australis</i> subsp. <i>parviflora</i>	n,q				S <sup>36</sup>
<i>Sloanea macbrydei</i>	q				FR <sup>36</sup>
EPACRIDACEAE					
<i>Acrotriche cordata</i>	s,t,v,w			9	HS <sup>9</sup>
<i>Acrotriche fasciculiflora</i>	s				S <sup>57</sup>
<i>Acrotriche halmaturina</i>	s	19			
<i>Acrotriche patula</i>	s,w				HS <sup>41</sup>
<i>Acrotriche serrulata</i>	n,s,t,v	4,17,25, 29			S <sup>43</sup>
<i>Andersonia auriculata</i>	w				HS <sup>41</sup>
<i>Andersonia axilliflora</i> E	w	58			S <sup>49</sup> ,HS <sup>41,58,59</sup>
<i>Andersonia bifida</i>	w				HS <sup>41</sup>
<i>Andersonia caerulea</i>	w				S <sup>46,51</sup> ,HS <sup>40,59</sup>
<i>Andersonia carinata</i>	w				HS <sup>41</sup>
<i>Andersonia echinocephala</i>	w	58			HS <sup>41,58,59</sup> ,MS <sup>40</sup>
<i>Andersonia ferricola</i>	w	39			S <sup>39</sup>
<i>Andersonia grandiflora</i>	w				HS <sup>41</sup> ,MS <sup>58</sup>
<i>Andersonia heterophylla</i>	w	37			S <sup>37,46</sup>
<i>Andersonia lehmanniana</i>	w				S <sup>37,46</sup>
<i>Andersonia longifolia</i>	w				HS <sup>41</sup>
<i>Andersonia macranthera</i>	w				HS <sup>41</sup>
<i>Andersonia pinaster</i> V	w	58			HS <sup>40,41,58</sup>
<i>Andersonia setifolia</i>	w				HS <sup>41</sup>
<i>Andersonia simplex</i>	w	51			S <sup>51</sup> ,MS <sup>40</sup>
<i>Andersonia sprengelioides</i>	w				HS <sup>40,59</sup>
<i>Astroloba baxteri</i>	w				LS <sup>40</sup>
<i>Astroloba ciliatum</i>	w	48,28			S <sup>48</sup>
<i>Astroloba conostephioides</i>	s,v	25,29			S <sup>43</sup>
<i>Astroloba foliosum</i>	w				HS <sup>41</sup>
<i>Astroloba humifusum</i>	n,s,t,v,w	17,25,34			S <sup>43</sup> ,HS <sup>10</sup>
<i>Astroloba microcalyx</i>	w	14			
<i>Astroloba microphyllum</i>	w				LS <sup>41</sup>
<i>Astroloba pallidum</i>	w				FR <sup>21,24</sup>
<i>Astroloba pinifolium</i>	n,t,v	17,34			S <sup>60</sup>
<i>Astroloba xerophyllum</i>	w	37			S <sup>37,46,55</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Astroloma</i> sp. Cataby (EA Griffin 1022)	w				HS <sup>41</sup>
<i>Astroloma</i> sp. Eneabba (N Merchant s.n.)	w				HS <sup>41</sup>
<i>Astroloma</i> sp. Fitzgerald (GJ Keighery 8376)	w				LS <sup>41</sup>
<i>Astroloma</i> sp. Grass Patch (AJG Wilson 110)	w				LS <sup>41</sup>
<i>Astroloma</i> sp. Mt Lindesay	w				HS <sup>59</sup>
<i>Astroloma</i> sp. Nannup (RD Royce 3978)	w				HS <sup>41</sup>
<i>Astroloma</i> sp. 2504	w				HS <sup>40</sup>
<i>Brachyloma ciliatum</i>	s,t,v	25			S <sup>43</sup>
<i>Brachyloma daphnoides</i> s.l.	n,q,s,v	3,25,29			S <sup>43</sup>
<i>Brachyloma depressum</i>	t,v	29			HS <sup>29</sup>
<i>Coleanthera coelophylla</i>	w				HS <sup>41</sup>
<i>Coleanthera virgata</i>	w				LS <sup>41</sup>
<i>Conostephium marchantiorum</i>	w				LS <sup>41</sup>
<i>Conostephium minus</i>	w				HS <sup>41</sup>
<i>Conostephium pendulum</i>	w	14,24,37			S <sup>46</sup>
<i>Conostephium uncinatum</i>	w				LS <sup>41</sup>
<i>Cosmelia rubra</i>	w				MS <sup>40</sup>
<i>Cyathodes glauca</i>	t	2,17,34			
<i>Epacris acuminata</i> E	t	34			
<i>Epacris apsleyensis</i> E	t				HS <sup>9</sup>
<i>Epacris barbata</i> CE	t				HS <sup>61</sup>
<i>Epacris corymbiflora</i>	t	17		33	HS <sup>61</sup>
<i>Epacris curtisiae</i>	t			9	HS <sup>60</sup>
<i>Epacris exserta</i> E	t			9	MS <sup>9</sup>
<i>Epacris glabella</i> E	t			9	MS <sup>9</sup>
<i>Epacris grandis</i> E	t			9	HS <sup>9</sup>
<i>Epacris gunnii</i>	n,q,s,t,v	8,17			
<i>Epacris impressa</i>	n,s,t,v	2,4,13,17 18,25,29, 34	3		S <sup>43</sup> ,LS <sup>60</sup>
<i>Epacris lanuginosa</i>	t	17,34			S <sup>60</sup>
<i>Epacris limbata</i> CE	t			9	HS <sup>9</sup>
<i>Epacris marginata</i>	t			9	HS <sup>9</sup>
<i>Epacris myrtifolia</i>	t			9	HS <sup>9</sup>
<i>Epacris obtusifolia</i>	n,q,t,v	17			S <sup>60</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Epacris paludosa</i>	n,t,v			9	HS <sup>9</sup>
<i>Epacris purpurascens</i>	n	1			
<i>Epacris stuartii CE</i>	t			9	MS <sup>9</sup>
<i>Epacris virgata</i>	t			9	HS <sup>9</sup>
<i>Epacridaceae gen. nov. (aff. Melichrus)</i>	w				HS <sup>41</sup>
<i>Gaultheria hispida</i>	t	17,34			
<i>Leptecophylla juniperina</i>	t	17			S <sup>60</sup>
<i>Leptecophylla pendulosa</i>	t			9	HS <sup>9</sup>
<i>Leucopogon amplexens</i>	w				HS <sup>41</sup>
<i>Leucopogon apiculatus</i>	w				HS <sup>41</sup>
<i>Leucopogon atherolepis</i>	w	58			HS <sup>59</sup> ,MS <sup>58</sup>
<i>Leucopogon australis s.l.</i>	t,v,w	14,17,25			S <sup>37,43,46,53</sup> LS <sup>40,59</sup>
<i>Leucopogon blepharolepis</i>	w				HS <sup>41</sup>
<i>Leucopogon bracteolaris</i>	w				HS <sup>41</sup>
<i>Leucopogon brevicuspis</i>	w				HS <sup>41</sup>
<i>Leucopogon breviflorus</i>	w				HS <sup>41</sup>
<i>Leucopogon capitellatus</i>	w	14,28			S <sup>24,37,46,48</sup> MS <sup>21,40</sup>
<i>Leucopogon collinus</i>	s,t,v	17,34			HS <sup>60</sup>
<i>Leucopogon concinnus</i>	w				S <sup>46</sup>
<i>Leucopogon concurvus</i>	s				S <sup>20</sup>
<i>Leucopogon conostephiooides</i>	w	14,24,37			S <sup>46</sup> ,LS <sup>40</sup> ,HS <sup>14</sup>
<i>Leucopogon cordifolius</i>	s,v,w				HS <sup>41</sup>
<i>Leucopogon corifolius</i>	w				MS <sup>40</sup>
<i>Leucopogon cryptanthus</i>	w				HS <sup>41</sup>
<i>Leucopogon cucullatus</i>	w				LS <sup>40</sup>
<i>Leucopogon cymbiformis</i>	w				MS <sup>40</sup>
<i>Leucopogon denticulatus</i>	w				HS <sup>41</sup>
<i>Leucopogon distans var. contractus</i>	w				S <sup>40</sup>
<i>Leucopogon distans s.l.</i>	w				HS <sup>40</sup>
<i>Leucopogon elegans</i>	w	51			S <sup>51</sup>
<i>Leucopogon ericoides</i>	n,q,s,t,v	17,25,29, 34	3		S <sup>43</sup> ,HS <sup>60</sup>
<i>Leucopogon esquamatus</i>	n,t,v			9	MS <sup>9</sup>
<i>Leucopogon flavescens</i>	w				S <sup>46,51</sup>
<i>Leucopogon florulentus</i>	w				HS <sup>41</sup>
<i>Leucopogon gibbosus</i>	w				HS <sup>40,59</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Leucopogon glabellus</i>	w	15			
<i>Leucopogon glacialis</i>	s,v	25,29			S <sup>43</sup>
<i>Leucopogon glaucifolius</i>	w				HS <sup>41</sup>
<i>Leucopogon gnaphalooides E</i>	w	58			HS <sup>58,59</sup>
<i>Leucopogon gracillimus</i>	w				S <sup>46</sup>
<i>Leucopogon interruptus</i>	w				HS <sup>41</sup>
<i>Leucopogon lanceolatus s.l.</i>	n,q,s,t,v	2		9	FR <sup>22</sup>
<i>Leucopogon lasiophyllus</i>	w				MS <sup>40,59</sup> , HS <sup>41</sup>
<i>Leucopogon lasiostachyus</i>	w	15			LS <sup>40,59</sup>
<i>Leucopogon macreraei</i>	n,v		3		
<i>Leucopogon marginatus E</i>	w				HS <sup>41</sup>
<i>Leucopogon microphyllus var. pilibundus</i>	n,v	3			
<i>Leucopogon multiflorus</i>	w				HS <sup>41</sup>
<i>Leucopogon nutans</i>	w	14,22			S <sup>24,46</sup> ,HS <sup>14</sup> LS <sup>40</sup> ,SP <sup>21</sup>
<i>Leucopogon obtectus E</i>	w				S <sup>52</sup> ,HS <sup>41</sup>
<i>Leucopogon oliganthus</i>	w				HS <sup>41</sup>
<i>Leucopogon oxycedrus</i>	w	48,28			SP <sup>21,48</sup> ,HS <sup>40</sup>
<i>Leucopogon parviflorus</i>	n,q,s,t,v,w				LS <sup>59</sup> , S <sup>46</sup>
<i>Leucopogon pendulus</i>	w				FR <sup>40</sup>
<i>Leucopogon pleurandroides</i>	w				HS <sup>41</sup>
<i>Leucopogon plumulifolius</i>	w				HS <sup>41</sup>
<i>Leucopogon pagonocalyx</i>	w				HS <sup>41</sup>
<i>Leucopogon polymorphus</i>	w				S <sup>37,53</sup>
<i>Leucopogon polystachyus</i>	w				HS <sup>41</sup>
<i>Leucopogon propinquus</i>	w	14,28			FR <sup>59</sup> ,S <sup>24,37,46,48</sup>
<i>Leucopogon pulchellus</i>	w	15			S <sup>54</sup>
<i>Leucopogon revolutus</i>	w				S <sup>46</sup>
<i>Leucopogon tamariscinus</i>	w				HS <sup>41</sup>
<i>Leucopogon unilateralis</i>	w				LS <sup>59</sup>
<i>Leucopogon verticillatus</i>	w	14,15,28			S <sup>24,46,48</sup> ,HS <sup>14,21</sup>
<i>Leucopogon virgatus</i>	n,q,s,t,v	3,17,18, 25,29			S <sup>43</sup>
<i>Leucopogon sp. Cascades (MA Burgman 3700)</i>	w				HS <sup>41</sup>
<i>Leucopogon sp. Clyde Hill (MA Burgman 1207)</i>	w				HS <sup>41</sup>
<i>Leucopogon sp. Condingup (MA</i>	w				HS <sup>41</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
Burgman 1377)					
<i>Leucopogon</i> sp. Coujinup (MA Burgman 1085)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. Dundas (MA Burgman 1482)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. Kau Rock (MA Burgman 1126)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. Munglinup (KR Newbey 8123)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. Peak Charles (MA Burgman 1476)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. Roberts Swamp (KR Newbey 8173)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. Yanneymooning (F Mollemans 3797)	w				HS <sup>41</sup>
<i>Leucopogon</i> sp. 4	w				LS <sup>40</sup>
<i>Lissanthe strigosa</i> s.l.	n,q,s,t,v	4,25			
<i>Lysinema ciliatum</i>	w			37	S <sup>37</sup> ,HS <sup>40,59</sup>
<i>Lysinema conspicuum</i>	w				MS <sup>40</sup>
<i>Lysinema elegans</i>	w				MS <sup>41</sup>
<i>Lysinema lasianthum</i>	w				LS <sup>41</sup>
<i>Melichrus urceolatus</i>	n,q,v	3			
<i>Monotoca elliptica</i>	n,t,v	17,34			S <sup>60</sup>
<i>Monotoca glauca</i>	t,v	2,17,34			S <sup>60</sup>
<i>Monotoca leucantha</i>	w				HS <sup>41</sup>
<i>Monotoca linifolia</i> susbp. <i>linifolia</i>	t	34			
<i>Monotoca oligarrhenoides</i>	w				HS <sup>59</sup>
<i>Monotoca scoparia</i>	n,q,t,v	3,25,29			
<i>Monotoca submutica</i> s.l.	t	17,34			
<i>Monotoca tamariscina</i>	w	15			S <sup>46</sup>
<i>Monotoca</i> sp. aff. <i>elliptica</i> (D. Albrecht pers. comm.)	n,v	13			HS <sup>13</sup>
<i>Monotoca</i> sp. Mt Maxwell (KR Newbey 4727)	w				LS <sup>41</sup>
<i>Prionotes cerinthoides</i>	t	17			
<i>Richea dracophylla</i>	t	34			
<i>Richea milliganii</i>	t	17,34			
<i>Richea pandanifolius</i>	t	17,34			HS <sup>60</sup>
<i>Sphenotoma dracophylloides</i>	w				HS <sup>40,59</sup>
<i>Sphenotoma drummondii</i> E	w	58			HS <sup>58,59</sup> ,LS <sup>41</sup>
<i>Sphenotoma gracilis</i>	w				HS <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Sphenotoma parviflora</i>	w				LS <sup>41</sup>
<i>Sphenotoma squarrosa</i>	w	15	53		S <sup>53</sup> ,HS <sup>40,59</sup>
<i>Sphenotoma</i> sp. Stirling	w	58			HS <sup>58,59</sup>
<i>Sprengelia incarnata</i>	n,s,t,v	8,13,17, 25,34		33	SP <sup>61</sup>
<i>Styphelia adscendens</i>	n,s,t,v	25,29,34			S <sup>60</sup>
<i>Styphelia pulchella</i>	w				LS <sup>41</sup>
<i>Styphelia tenuiflora</i>	w	14,24,46			S <sup>24,46</sup> ,Q <sup>21</sup> ,HS <sup>22</sup>
<i>Trochocarpa disticha</i>	t	17			
<i>Trochocarpa gunnii</i>	t	17			
<i>Trochocarpa parviflora</i>	w				LS <sup>41</sup>
<i>Woollsia pungens</i>	n,q	1			
<b>ESCALLONIACEAE</b>					
<i>Anopterus glandulosus</i>	t	17,34			
<b>EUCRYPHIACEAE</b>					
<i>Eucryphia lucida</i>	t	17			
<i>Eucryphia milliganii</i>	t	17	3		
<i>Eucryphia moorei</i>	n		3		
<b>EUPHORBIACEAE</b>					
<i>Amperea ericoides</i>	w	15			
<i>Amperea xiphoclada</i>	n,q,s,t,v	17,25,34			S <sup>43,60</sup> , FR <sup>22</sup>
<i>Antidesma erostre</i>	q				FR <sup>36</sup>
<i>Hylandia dockrillii</i>	q				FR <sup>36</sup>
<i>Macaranga subdentata</i>	q				FR <sup>36</sup>
<i>Mallotus polyadenos</i>	q				S <sup>36</sup>
<i>Monotaxis occidentalis</i>	w				Q <sup>21</sup>
<i>Phyllanthus calycinus</i>	s,w	24			FR <sup>21,24</sup>
<i>Phyllanthus hirtellus</i>	n,v	3			
<i>Poranthera corymbosa</i>	n,q,v				
<i>Poranthera microphylla</i>	n,nt,q,s,t,v,w	25			
<i>Ricinocarpus glaucus</i>	w		3		
<i>Ricinocarpus pinifolius</i>	n,q,t,v	17			
<i>Stachystemon vermicularis</i>	w				FR <sup>21</sup>
<b>FABACEAE</b>					
<i>Aotus ericoides</i>	n,q,t,v,w	15,17,25, 34			S <sup>37,43,53,54</sup> ,HS <sup>60</sup>
<i>Aotus genistoides</i>	w				MS <sup>59</sup> ,S <sup>58</sup>
<i>Aotus passerinoides</i>	w	15			S <sup>53,54</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Bossiaea aquifolium</i>	w	14			SP <sup>21</sup>
<i>Bossiaea cinerea</i>	n,s,v	17,25,34			S <sup>43</sup> ,HS <sup>60</sup>
<i>Bossiaea eriocarpa</i>	w	14,15,24			S <sup>51</sup>
<i>Bossiaea linophylla</i>	w	40			FR <sup>40</sup>
<i>Bossiaea obcordata</i>	n,q,t,v			9	LS <sup>9</sup>
<i>Bossiaea ornata</i>	w	14,28			S <sup>46,48</sup> ,SP <sup>21</sup>
<i>Bossiaea prostrata</i>	n,q,s,t,v	25,29			S <sup>43,60</sup>
<i>Bossiaea rufa</i>	w	40			FR <sup>40</sup>
<i>Bossiaea webbii</i>	w	40			FR <sup>40,59</sup>
<i>Castanospermum australe</i>	q			16	
<i>Chorizema aciculare</i> s.l.	w	40			FR <sup>40</sup>
<i>Chorizema carinatum</i>	w				MS <sup>40</sup>
<i>Chorizema rhombeum</i>	w				Q <sup>21</sup>
<i>Daviesia brevifolia</i>	n,s,v	25,29			S <sup>43</sup>
<i>Daviesia bursarioides</i> E	w				MS <sup>41</sup>
<i>Daviesia chapmanii</i>	w				MS <sup>41</sup>
<i>Daviesia debilior</i> subsp. <i>sinuans</i>	w				MS <sup>41</sup>
<i>Daviesia decurrens</i>	w	15			S <sup>24</sup> , SP <sup>21</sup>
<i>Daviesia dielsii</i>	w				MS <sup>41</sup>
<i>Daviesia epiphyllum</i>	w				MS <sup>41</sup>
<i>Daviesia euphorbioides</i> E	w				MS <sup>41</sup>
<i>Daviesia glossosema</i>	w	58			HS <sup>58</sup>
<i>Daviesia incrassata</i> s.l.	w				S <sup>37</sup>
<i>Daviesia inflata</i>	w				S <sup>56</sup> ,HS <sup>40</sup>
<i>Daviesia latifolia</i>	n,q,t,v	17,25,34			S <sup>43</sup>
<i>Daviesia lineata</i>	w				MS <sup>41</sup>
<i>Daviesia megacalyx</i> E	w				S <sup>49</sup> ,LS <sup>41</sup>
<i>Daviesia mesophylla</i>	w				HS <sup>58</sup>
<i>Daviesia microcarpa</i> E	w				LS <sup>41</sup>
<i>Daviesia microphylla</i>	w				LS <sup>41</sup>
<i>Daviesia mimosoides</i> s.l.	n,q,v	2,3			
<i>Daviesia oppositifolia</i>	w				LS <sup>40</sup>
<i>Daviesia obovata</i>	w	58			HS <sup>58</sup>
<i>Daviesia ovata</i>	w				LS <sup>41</sup>
<i>Daviesia oxylobium</i>	w				MS <sup>41</sup>
<i>Daviesia pauciflora</i>	w				LS <sup>41</sup>
<i>Daviesia physodes</i>	w				S <sup>56</sup> ,MS <sup>41</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Daviesia polyphylla</i>	w	14			
<i>Daviesia preissii</i>	w				HS <sup>21</sup>
<i>Daviesia pseudophylla E</i>	w	58			MS <sup>58</sup> ,S <sup>49</sup> ,LS <sup>41</sup>
<i>Daviesia pteroclada</i>	w				MS <sup>41</sup>
<i>Daviesia purpurascens</i>	w				LS <sup>41</sup>
<i>Daviesia rhombifolia</i>	w	14,24			S <sup>24</sup>
<i>Daviesia speciosa E</i>	w				MS <sup>41</sup>
<i>Daviesia spiralis</i>	w				MS <sup>41</sup>
<i>Daviesia ulicifolia</i>	n,q,s,t,v,w	2,17,25, 34			S <sup>43,60</sup>
<i>Daviesia wyattiana</i>	n,q,v				HS <sup>13</sup>
<i>Daviesia</i> sp. [CAM] (KR Newbey 8162)	w				LS <sup>41</sup>
<i>Daviesia</i> sp. [PLE] (AS George 10288)	w				LS <sup>41</sup>
<i>Daviesia</i> sp. 4	w				HS <sup>40</sup>
<i>Dillwynia glaberrima</i>	n,t,v	17,25,29, 34	3		S <sup>43</sup> ,HS <sup>60</sup>
<i>Dillwynia phylloides</i>	n,q,v	2,3,25			S <sup>43</sup>
<i>Dillwynia sericea</i>	n,q,s,t,v	17,25,29, 34			S <sup>43</sup> ,HS <sup>60</sup>
<i>Dillwynia uncinata</i>	s,v,w	15			
<i>Dillwynia</i> sp. A	w				FR <sup>21</sup>
<i>Eutaxia densifolia</i>	w				LS <sup>40</sup>
<i>Gastrolobium bilobum</i>	w				FR <sup>59</sup>
<i>Gastrolobium crenulatum</i>	w				S <sup>58</sup>
<i>Gastrolobium leakeanum</i>	w				HS <sup>58,59</sup> ,LS <sup>40</sup>
<i>Gastrolobium luteifolium</i>	w	58			HS <sup>58</sup>
<i>Gastrolobium mondurup</i>	w				HS <sup>58</sup> ,MS <sup>59</sup>
<i>Gastrolobium papilio</i>	w				S <sup>49</sup>
<i>Gastrolobium pulchellum</i>	w	58			HS <sup>58</sup> , MS <sup>59</sup> ,S <sup>46</sup>
<i>Gastrolobium spinosum</i>	w				LS <sup>40</sup>
<i>Gastrolobium tetragonophyllum</i>	w				MS <sup>40</sup>
<i>Gastrolobium rubrum</i>	w				MS <sup>58,59</sup>
<i>Gompholobium capitatum</i>	w	24			FR <sup>24</sup>
<i>Gompholobium confertum</i>	w	15			S <sup>58</sup> ,HS <sup>40</sup>
<i>Gompholobium ecostatum</i>	s,t,v	25,29			
<i>Gompholobium huegelii</i>	n,q,t,v	17,25,34			
<i>Gompholobium knightianum</i>	w	14			SP <sup>21</sup>
<i>Gompholobium marginatum</i>	w				FR <sup>21</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Gompholobium polymorphum</i>	w			21	FR <sup>21</sup>
<i>Gompholobium preissii</i>	w				FR <sup>21</sup>
<i>Hovea chorizemifolia</i>	w	24		21	FR <sup>24,59</sup> ,Q <sup>21</sup>
<i>Hovea corrickeriae</i>	t,v			9	MS <sup>9</sup>
<i>Hovea elliptica</i>	w	15,54			S <sup>54</sup>
<i>Hovea linearis</i>	n,q,s,t,v	25,29			S <sup>43</sup>
<i>Hovea pungens</i>	w		53		S <sup>53</sup>
<i>Jacksonia alata</i>	w				FR <sup>21</sup>
<i>Jacksonia calycina</i>	w				S <sup>58</sup>
<i>Jacksonia carduacea</i>	w				MS <sup>41</sup>
<i>Jacksonia floribunda</i>	w	37			S <sup>37,46,54</sup>
<i>Jacksonia furcellata</i>	w	24			HS <sup>24</sup>
<i>Jacksonia grevilleoides</i>	w				MS <sup>40</sup>
<i>Jacksonia horrida</i>	w				S <sup>46</sup>
<i>Jacksonia sericea</i>	w				MS <sup>41</sup>
<i>Jacksonia spinosa</i>	w	51			S <sup>51</sup> ,LS <sup>40</sup>
<i>Jacksonia sternbergiana</i>	w	24			HS <sup>24</sup>
<i>Jacksonia sp. aff. furcelata</i>	w				MS <sup>40</sup>
<i>Kennedia coccinea</i>	w	14,24		21	FR <sup>24</sup> ,MS <sup>21</sup>
<i>Kennedia prostrata</i>	n,s,t,v	29,24			FR <sup>24</sup>
<i>Latrobea genistoides</i>	w				S <sup>51</sup>
<i>Latrobea hirtella</i>	w				S <sup>51</sup>
<i>Mirbelia dilatata</i>	w	24			FR <sup>24</sup>
<i>Oxylobium arborescens</i>	n,q,t,v	2,8,17			
<i>Oxylobium ellipticum</i>	n,t,v	2,17,35			FR <sup>22</sup> HS <sup>35</sup>
<i>Oxylobium ilicifolium</i>	n,q,v	2			
<i>Phyllota diffusa</i>	t	17,34			HS <sup>34</sup>
<i>Platylobium formosum</i>	n,q,t,v	25,34			S <sup>43</sup> ,HS <sup>60</sup>
<i>Platylobium obtusangulum</i>	s,t,v	4,17,29, 34			S <sup>43</sup>
<i>Pultenaea altissima</i>	n,v		3		
<i>Pultenaea benthamii</i>	n,v	13			
<i>Pultenaea cunninghamii</i>	n,q,v	2			
<i>Pultenaea daphnoides</i>	n,q,s,t,v	2,13,17,3 4			HS <sup>22,34</sup>
<i>Pultenaea ericifolia</i>	w				FR <sup>40</sup>
<i>Pultenaea graveolens</i>	v	32			HS <sup>32</sup>
<i>Pultenaea gunnii</i>	s,t,v	17,25,34			S <sup>43</sup> ,HS <sup>34</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Pultenaea hibbertioides</i>	t,v			9	HS <sup>9</sup>
<i>Pultenaea humilis</i>	n,t,v	4,25,29			S <sup>43</sup>
<i>Pultenaea involucrata</i>	s	18			
<i>Pultenaea juniperina</i> s.l.	n,t,v	17,34			HS <sup>34</sup>
<i>Pultenaea mollis</i>	v	9,25			S <sup>43</sup>
<i>Pultenaea palacea</i> var. <i>sericea</i>	n,t,v			9	HS <sup>9</sup>
<i>Pultenaea pedunculata</i>	n,s,t,v	4,25,34			S <sup>43</sup>
<i>Pultenaea procumbens</i>	n,v	3			
<i>Pultenaea prostrata</i>	s,t,v			9	HS <sup>9</sup>
<i>Pultenaea pycnocephala</i>	n,q	13			
<i>Pultenaea reticulata</i>	w	2,15			HS <sup>40,59</sup>
<i>Pultenaea scabra</i>	n,s,v	25			S <sup>43</sup>
<i>Pultenaea stricta</i>	s,t,v	17,25,34			S <sup>43</sup> ,HS <sup>34</sup>
<i>Pultenaea subalpina</i>	v		3,44		HS <sup>44</sup>
<i>Pultenaea subspicata</i>	n,v		3		
<i>Pultenaea trifida</i>	s	19			
<i>Sphaerolobium acanthos</i>	v		44		MS <sup>44</sup>
<i>Sphaerolobium medium</i>	w				FR <sup>21</sup>
<b>FAGACEAE</b>					
<i>Nothofagus cunninghamii</i>	t,v	2,17,25			S <sup>43</sup>
<b>GLEICHENIACEAE</b>					
<i>Gleichenia dicarpa</i>	n,q,t,v	25			S <sup>43</sup>
<b>GOODENIACEAE</b>					
<i>Coopernochla barbata</i>	n,v				
<i>Dampiera alata</i>	w				S <sup>37</sup>
<i>Dampiera fasciculata</i>	w				FR <sup>59</sup>
<i>Dampiera linearis</i>	w	15			FR <sup>21</sup>
<i>Goodenia caerulea</i>	w				FR <sup>21,40,59</sup>
<i>Goodenia geniculata</i>	n,q,s,t,v	25			
<i>Goodenia hederacea</i> s.l.	n,q,v	25,29			S <sup>43</sup>
<i>Goodenia humilis</i>	n,s,t,v	25,29			S <sup>43</sup>
<i>Goodenia lanata</i>	t,v	25,29			S <sup>43</sup>
<i>Goodenia ovata</i>	n,q,s,t,v	25			
<i>Goodenia scapigera</i>	w				FR <sup>40,59</sup>
<i>Lechenaultia biloba</i>	w	14,24	3		FR <sup>21,24</sup>
<i>Lechenaultia floribunda</i>	w		3		
<i>Lechenaultia formosa</i>	w		3		

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Lechenaultia hirsuta</i>	w		3		
<i>Scaevola calliptera</i>	w			21	Q <sup>21</sup>
<i>Scaevola striata</i> s.l.	w	24			FR <sup>24</sup>
<i>Scaevola thesioides</i> s.l.	w		3		
<i>Velleia foliosa</i>	w				FR <sup>58</sup>
<b>GROSSULARIACEAE</b>					
<i>Polyosma alangiacea</i>	q				FR <sup>36</sup>
<b>HAEMODORACEAE</b>					
<i>Anigozanthus flavidus</i>	w				FR <sup>54</sup>
<i>Anigozanthus manglesii</i>	w				FR <sup>22</sup>
<i>Anigozanthus rufus</i>	w				FR <sup>54</sup>
<i>Conostylis aculeata</i> s.l.	w	24			FR <sup>24</sup>
<i>Conostylis misera</i> E	w				FR <sup>58</sup>
<i>Conostylis pusilla</i>	w	24			FR <sup>24</sup>
<i>Conostylis serrulata</i>	w	24			FR <sup>21,24</sup>
<i>Conostylis setigera</i> s.l.	w	15			FR <sup>21,40</sup>
<i>Conostylis setosa</i>	w	24			FR <sup>21,40</sup>
<i>Macropidia fuliginosa</i>	w		54		FR <sup>54</sup>
<b>HALORAGACEAE</b>					
<i>Gonocarpus benthamii</i> subsp. Stirling	w				FR <sup>58</sup>
<i>Gonocarpus mezianus</i>	s,v	25			S <sup>43</sup>
<i>Gonocarpus rудis</i>	w				FR <sup>58</sup>
<i>Gonocarpus tetragynus</i>	n,q,s,t,v	17,34			
<i>Gonocarpus teucrioides</i>	n,q,t,v	2,17,34			FR <sup>42</sup>
<i>Haloragodendron monospermum</i>	n		3		
<b>ICACINACEAE</b>					
<i>Apodytes brachystylis</i>	q				FR <sup>36</sup>
<i>Citronella smythii</i>	q				FR <sup>36</sup>
<i>Irvingbaileya australis</i>	q				FR <sup>36</sup>
<b>IRIDACEAE</b>					
<i>Diplarrena moraea</i>	n,t,v	8,13,17			
<i>Isophysis tasmanica</i>	t	34			HS <sup>61</sup>
<i>Patersonia babianoides</i>	w				Q <sup>21</sup>
<i>Patersonia fragilis</i>	n,q,s,t,v	17,34			
<i>Patersonia glabrata</i>	n,q,v				S <sup>42</sup>
<i>Patersonia occidentalis</i>	s,t,v,w	14,24			FR <sup>59</sup> ,S <sup>24,37,46,53</sup> FR <sup>60</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Patersonia pygmaea</i>	w	28			HS <sup>28</sup> ,FR <sup>21,24</sup>
<i>Patersonia rufa</i>	w	14			S <sup>46</sup> ,SP <sup>21</sup>
<i>Patersonia sericea</i>	n,q,v	2			
<i>Patersonia umbrosa</i>	w	14			FR <sup>59</sup> ,S <sup>46</sup>
<b>LAMIACEAE</b>					
<i>Gmelina fasciculiflora</i>	q				FR <sup>36</sup>
<i>Hemigenia curvifolia</i>	w	24			FR <sup>24</sup>
<i>Hemigenia ramosissima</i>	w				FR <sup>21</sup>
<i>Prostanthera cuneata</i>	n,t,v		3		
<i>Prostanthera decussata</i>	v	32			S <sup>32</sup>
<i>Prostanthera lasianthos</i>	n,q,t,v	17	3		
<i>Prostanthera ovalifolia</i>	n,q		3		
<i>Prostanthera ringens</i>	n,q		3		
<i>Prostanthera saxicola</i> var. <i>montana</i>	n		3		
<b>LAURACEAE</b>					
<i>Beilschmiedia bancroftii</i>	q				S <sup>36</sup>
<i>Beilschmiedia collina</i>	q	16			
<i>Beilschmiedia tooram</i>	q				S <sup>36</sup>
<i>Cassytha glabella</i>	n,q,s,t,v,w				FR <sup>22</sup>
<i>Cinnamomum oliveri</i>	q	16			HS <sup>45</sup>
<i>Cryptocarya angulata</i>	q				FR <sup>36</sup>
<i>Cryptocarya corrugata</i>	q				FR <sup>36</sup>
<i>Cryptocarya densiflora</i>	q				FR <sup>36</sup>
<i>Cryptocarya corrugata</i>	q	16			FR <sup>36</sup>
<i>Cryptocarya glaucescens</i>	q	16			
<i>Cryptocarya leucophylla</i>	q				FR <sup>36</sup>
<i>Cryptocarya lividula</i>	q				FR <sup>36</sup>
<i>Cryptocarya mackinnoniana</i>	q				S <sup>36</sup>
<i>Cryptocarya putida</i>	q				FR <sup>36</sup>
<i>Endiandra bessaphila</i>	q				S <sup>36</sup>
<i>Endiandra dichrophylla</i>	q				FR <sup>36</sup>
<i>Endiandra monothyra</i> subsp. <i>monothyra</i>	q				FR <sup>36</sup>
<i>Endiandra montana</i>	q				FR <sup>36</sup>
<i>Endiandra sankeyana</i>	q				FR <sup>36</sup>
<i>Endiandra wolfei</i>	q				FR <sup>36</sup>
<b>LENTIBULARIACEAE</b>					

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Utricularia multifida</i>	w				FR <sup>40</sup>
LINDSAEACEAE					
<i>Lindsaea linearis</i>	n,q,s,t,v,w				FR <sup>40,60</sup>
LOGANIACEAE					
<i>Logania serpyllifolia</i> s.l.	w			21	FR <sup>40,59</sup> ,Q <sup>21</sup>
<i>Mitrasacme pilosa</i>	n,s,t,v	17			
LORANTHACEAE					
<i>Nuytsia floribunda</i>	w				FR <sup>22</sup>
LYCOPODIACEAE					
<i>Lycopodium deuterodensum</i>	n,q,s,t,v	35			SV <sup>22</sup>
MELIACEAE					
<i>Synoum glandulosum</i> subsp. <i>paniculatum</i>	q			16	FR <sup>36</sup>
MIMOSACEAE					
<i>Acacia aculeatissima</i>	n,v	25			
<i>Acacia axillaris</i> V	t			9	MS <sup>9</sup>
<i>Acacia barbinervis</i> s.l.	w	24			FR <sup>21,24</sup>
<i>Acacia baxteri</i>	w				FR <sup>56</sup>
<i>Acacia browniana</i> s.l.	w	24			FR <sup>24</sup>
<i>Acacia browniana</i> var. <i>intermedia</i>	w				FR <sup>40</sup>
<i>Acacia buxifolia</i> subsp. <i>buxifolia</i>	n,q,v	3			
<i>Acacia campylophylla</i>	w	24			S <sup>24</sup>
<i>Acacia cyclops</i>	w				FR <sup>40</sup>
<i>Acacia dealbata</i>	n,t,v	2			FR <sup>60</sup>
<i>Acacia drummondii</i> s.l.	w	24			FR <sup>24,59</sup>
<i>Acacia extensa</i>	w				FR <sup>24</sup>
<i>Acacia genistifolia</i>	n,t,v	2,3			
<i>Acacia horridula</i>	w	3			
<i>Acacia lateriticola</i>	w				FR <sup>24,Q<sup>21</sup></sup>
<i>Acacia melanoxylon</i>	n,q,s,t,v	17			FR <sup>60</sup>
<i>Acacia mitchelli</i>	n,s,v	25			S <sup>43</sup>
<i>Acacia mucronata</i> s.l.	n,q,t,v	17,25			S <sup>43</sup>
<i>Acacia myrtifolia</i>	n,q,s,t,v,w	17,18,25, 29			S <sup>43,55</sup> ,MS <sup>40,59</sup>
<i>Acacia nervosa</i>	w				FR <sup>21,24</sup>
<i>Acacia obovata</i>	w				FR <sup>21</sup>
<i>Acacia oxycedrus</i>	n,s,v	25			S <sup>43</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Acacia paradoxa</i>	n,q,s,t,v,w				S <sup>57</sup>
<i>Acacia pataczekii</i>	t			9	LS <sup>9</sup>
<i>Acacia preissiana</i>	w				FR <sup>24</sup>
<i>Acacia pulchella</i> s.l.	w	24, 50	3		FR <sup>50</sup> ,LS <sup>40</sup>
<i>Acacia retinodes</i> s.l.	s,v			34	
<i>Acacia siculiformis</i>	n,t,v			9	MS <sup>9</sup>
<i>Acacia stenoptera</i>	w	14,24			
<i>Acacia suaveolens</i>	n,q,s,t,v	17,25			S <sup>43</sup> ,FR <sup>42</sup> ,LS <sup>60</sup>
<i>Acacia terminalis</i> s.l.	n,t,v				
<i>Acacia ulicifolia</i>	n,q,t,v	17			
<i>Acacia urophylla</i>	w			21	FR <sup>24,53</sup> ,Q <sup>21</sup>
<i>Acacia veronica</i>	w				FR <sup>58,59</sup>
<i>Acacia verticillata</i> s.l.	n,s,t,v	2,8,17,25			
<i>Paraserianthes lophantha</i>	w				FR <sup>22</sup>
<b>MONIMIACEAE</b>					
<i>Atherosperma moschatum</i>	n,t,v	17			
<i>Daphnandra repandula</i>	q				FR <sup>36</sup>
<b>MYRISTICACEAE</b>					
<i>Myristica insipida</i>	q			16	
<b>MYRSINACEAE</b>					
<i>Rapanea achradiifolia</i>	q				S <sup>36</sup>
<b>MYRTACEAE</b>					
<i>Acmena resa</i>	q			16	FR <sup>36</sup>
<i>Agonis floribunda</i>	w				LS <sup>59</sup>
<i>Agonis hypericifolia</i>	w	15			MS <sup>40,59</sup>
<i>Agonis linearifolia</i>	w	51			S <sup>51</sup> ,LS <sup>40</sup>
<i>Agonis parviceps</i>	w				LS <sup>40,59</sup>
<i>Agonis spathulata</i>	w				LS <sup>40,59</sup>
<i>Angophora costata</i>	n,q	12			
<i>Angophora hispida</i>	n				
<i>Astartea fascicularis</i>	w	40	3		FR <sup>40</sup>
<i>Astartea heteranthera</i>	w	50	3		FR <sup>50</sup>
<i>Austromyrtus</i> sp. (Gillies BG 1484)	q				FR <sup>36</sup>
<i>Baeckea camphorosmae</i>	w				FR <sup>21,24,50</sup>
<i>Baeckea leptocaulis</i>	t	17,34		33	HS <sup>34</sup> ,MS <sup>61</sup>
<i>Baeckea pachyphylla</i>	w	20			FR <sup>40</sup>
<i>Euryomyrtus ramosissima</i> subsp.	n,t,v	25			

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>prostrata</i>					
<i>Beaufortia anisandra</i>	w	51			S <sup>51</sup> ,MS <sup>40,59</sup>
<i>Beaufortia decussata</i>	w				LS <sup>40,59</sup>
<i>Beaufortia elegans</i>	w	37			
<i>Beaufortia emprtrifolia</i>	w				LS <sup>40</sup>
<i>Beaufortia eriocephala</i>	w				LS <sup>41</sup>
<i>Beaufortia micrantha</i>	w				MS <sup>40</sup>
<i>Beaufortia sparsa</i>	w	15	3		
<i>Calothamnus affinis</i>	w	40			FR <sup>40</sup> ,LS <sup>58</sup>
<i>Calothamnus crassus</i>	w	58			LS <sup>58,59</sup>
<i>Calothamnus quadrifidus</i>	w	40			FR <sup>40</sup>
<i>Calothamnus sanguineus</i>	w	40			FR <sup>40</sup>
<i>Calothamnus villosus</i>	w				S <sup>37</sup>
<i>Calytrix alpestris</i>	s,v		3		
<i>Calytrix asperula</i>	w	40			FR <sup>40</sup>
<i>Calytrix flavescens</i>	w	15			FR <sup>40</sup>
<i>Calytrix leschenaultii</i>	w	40			FR <sup>40</sup>
<i>Calytrix tenuiramea</i>	w	40			FR <sup>40</sup>
<i>Calytrix tetragona</i>	n,q,s,t,v,w	17,25,29, 34	3		S <sup>43</sup> ,SV <sup>60</sup>
<i>Chamelaucium ciliatum</i>	w		3		
<i>Chamelaucium erythrochlora</i>	w				S <sup>52</sup>
<i>Chamelaucium griffinii</i>	w				S <sup>52</sup>
<i>Chamelaucium roycei</i>	w				S <sup>52</sup>
<i>Corymbia calophylla</i>	w			21	FR <sup>24,50,53</sup>
<i>Corymbia ficifolia</i>	w				FR <sup>50</sup>
<i>Corymbia gummifera</i>	n,q,v	2			SV <sup>42</sup>
<i>Corymbia maculata</i>	n,q,v			31	FR <sup>22</sup>
<i>Corymbia tessellaris</i>	n,q			31	
<i>Darwinia citriodora</i>	w				FR <sup>50</sup>
<i>Darwinia collina E</i>	w	58			MS <sup>58</sup> ,S <sup>49,52</sup>
<i>Darwinia hypericifolia</i>	w				LS-MS <sup>58</sup>
<i>Darwinia leiostyla</i>	w				FR <sup>40</sup> ,LS-MS <sup>58</sup>
<i>Darwinia macrostegia</i>	w				S <sup>49</sup>
<i>Darwinia meboldii</i>	w				S <sup>49,52</sup>
<i>Darwinia oxylepis E</i>	w				HS <sup>58</sup> ,S <sup>46,52</sup>
<i>Darwinia squarrosa V</i>	w				LS <sup>59</sup> ,MS <sup>58</sup> ,S <sup>46</sup>
<i>Darwinia vestita</i>	w				FR <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Darwinia wittwerorum</i> E	w	58			MS <sup>58</sup> ,S <sup>46</sup>
<i>Darwinia</i> sp. Stirling Range V	w				MS <sup>58</sup>
<i>Eremaea pauciflora</i> s.l.	w	37	3		
<i>Eucalyptus accedens</i>	w				FR <sup>50</sup>
<i>Eucalyptus acmenoides</i>	q	2			
<i>Eucalyptus amygdalina</i>	t	2,17			FR <sup>60</sup>
<i>Eucalyptus andrewsii</i> s.l.	n,q			31	
<i>Eucalyptus angulosa</i>	w				FR <sup>40</sup>
<i>Eucalyptus aromaphloia</i>	n,v	25			S <sup>43</sup>
<i>Eucalyptus astringens</i>	w				FR <sup>50</sup>
<i>Eucalyptus baxteri</i>	n,s,v	2,4,18,25 29	3	31	S <sup>43</sup>
<i>Eucalyptus botryoides</i>	n,v			31	
<i>Eucalyptus buprestium</i>	w				FR <sup>40</sup>
<i>Eucalyptus camaldulensis</i>	n,q,s,v,w			31	FR <sup>50</sup>
<i>Eucalyptus cladocalyx</i>	s			31	
<i>Eucalyptus cloeziana</i>	q			31	
<i>Eucalyptus coccifera</i>	t	17			FR <sup>60</sup>
<i>Eucalyptus conferruminata</i>	w				FR <sup>50</sup>
<i>Eucalyptus consideniana</i>	n,v	25			S <sup>43</sup>
<i>Eucalyptus cordata</i>	t	17			FR <sup>60</sup>
<i>Eucalyptus dalrympleana</i> s.l.	n,t,v			31	FR <sup>60</sup>
<i>Eucalyptus decipiens</i> s.l.	w				LS <sup>40</sup>
<i>Eucalyptus decurva</i>	w				FR <sup>40</sup>
<i>Eucalyptus delegatensis</i>	n,t,v	8,17,25			S <sup>43</sup> ,FR <sup>60</sup>
<i>Eucalyptus diversicolor</i>	w	2			
<i>Eucalyptus dives</i>	n,v	2,25			S <sup>43</sup>
<i>Eucalyptus doratoxylon</i>	w				FR <sup>59</sup>
<i>Eucalyptus erectifolia</i>	w				FR <sup>59</sup>
<i>Eucalyptus eugenoides</i>	n,q	2			
<i>Eucalyptus falcata</i>	w				FR <sup>40</sup>
<i>Eucalyptus fastigata</i>	n,v,saf	25	27		S <sup>43</sup> ,HS <sup>27</sup>
<i>Eucalyptus fibrosa</i> s.l.	n,q			31	
<i>Eucalyptus forrestiana</i>	w				FR <sup>50</sup>
<i>Eucalyptus fraxinoides</i>	n,v,saf		27		HS <sup>27</sup>
<i>Eucalyptus gardneri</i>	w				FR <sup>50</sup>
<i>Eucalyptus globoidea</i>	n,v	2,25			S <sup>43</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Eucalyptus globulus</i> s.l.	n,q,s,t,v	2		31	FR <sup>60</sup>
<i>Eucalyptus gomphocephala</i>	w				FR <sup>50</sup>
<i>Eucalyptus goniocalyx</i>	n,s,v			31	
<i>Eucalyptus grandis</i>	n,q	2			
<i>Eucalyptus imlayensis</i> E	n				?HS <sup>22</sup>
<i>Eucalyptus krusana</i>	w				FR <sup>50</sup>
<i>Eucalyptus laeliae</i>	w				FR <sup>50</sup>
<i>Eucalyptus lehmannii</i>	w				FR <sup>56</sup>
<i>Eucalyptus ligulata</i> subsp. <i>stirlingica</i>	w				LS <sup>58</sup>
<i>Eucalyptus macrorhyncha</i>	n,s,v	2,4,25,29			S <sup>43</sup>
<i>Eucalyptus mannifera</i>	n,v	2			
<i>Eucalyptus marginata</i>	w	2,14,15, 24	46,53	31	S <sup>46,53</sup> ,MS <sup>24</sup> LS <sup>40,59</sup>
<i>Eucalyptus megacarpa</i>	w				FR <sup>50</sup>
<i>Eucalyptus niphophila</i>	n,v		3		
<i>Eucalyptus nitens</i>	t,v	8,25			S <sup>43</sup> ,LS <sup>60</sup>
<i>Eucalyptus nitida</i>	t	17			FR <sup>60</sup>
<i>Eucalyptus oblique</i>	n,q,s,t,v	2,17,18, 19,25,29		31	S <sup>43</sup> ,LS <sup>60</sup>
<i>Eucalyptus occidentalis</i>	w				FR <sup>50</sup>
<i>Eucalyptus ovata</i> s.l.	n,s,t,v	17			FR <sup>60</sup>
<i>Eucalyptus pachyloma</i>	w				FR <sup>40</sup>
<i>Eucalyptus paniculata</i>	n,q			31	
<i>Eucalyptus patens</i>	w				FR <sup>50</sup>
<i>Eucalyptus pauciflora</i> s.l.	n,t,v	25		31	FR <sup>22</sup>
<i>Eucalyptus pilularis</i>	n,q	2		31	
<i>Eucalyptus piperita</i> s.l.	n	2			
<i>Eucalyptus polyanthemos</i>	n,v	3			
<i>Eucalyptus preissiana</i>	w				FR <sup>40</sup>
<i>Eucalyptus pulchella</i>	t	17			FR <sup>60</sup>
<i>Eucalyptus racemosa</i> s.l.	n			31	
<i>Eucalyptus radiata</i>	n,q,t,v	2,4,25		31	S <sup>43</sup>
<i>Eucalyptus regnans</i>	t,v	2,25			S <sup>43</sup> ,FR <sup>60</sup>
<i>Eucalyptus rossii</i>	n	2			
<i>Eucalyptus rufa</i>	w,				FR <sup>50</sup>
<i>Eucalyptus saligna</i>	n,q	2			
<i>Eucalyptus sieberi</i>	n,t,v	2,17,25		31	S <sup>43</sup> ,LS <sup>60</sup>
<i>Eucalyptus smithii</i>	n,v,saf		27		HS <sup>27</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Eucalyptus spathulata</i>	w				FR <sup>50</sup>
<i>Eucalyptus staeri</i>	w				LS <sup>40,59</sup>
<i>Eucalyptus talyuberlup</i>	w				FR <sup>59</sup>
<i>Eucalyptus tenuiramis</i>	t	17	3		
<i>Eucalyptus todtiana</i>	w		37,53		S <sup>37,53</sup>
<i>Eucalyptus uncinata</i>	w				FR <sup>40</sup>
<i>Eucalyptus viminalis</i>	n,q,s,t,v	2,25		31	FR <sup>60</sup>
<i>Eucalyptus wandoo</i>	w				FR <sup>50</sup>
<i>Eucalyptus willisii</i> s.l.	s,v	25			S <sup>43</sup>
<i>Homalospermum firmum</i>	w				FR <sup>59</sup>
<i>Hypocalymma angustifolium</i> s.l.	w	15,28,24	3		FR <sup>21,24</sup>
<i>Hypocalymma cordifolium</i> s.l.	w	15			
<i>Hypocalymma myrtifolium</i>	w				FR <sup>40,59</sup>
<i>Hypocalymma phillipsii</i>	w				FR <sup>59</sup>
<i>Hypocalymma robustum</i>	w	14,15,24			S <sup>24,37,46</sup>
<i>Hypocalymma speciosum</i>	w				FR <sup>40</sup>
<i>Hypocalymma strictum</i> s.l.	w				HS <sup>40</sup>
<i>Kunzea ericifolia</i>	v	14,24	3		
<i>Kunzea montana</i>	w	58			LS <sup>58,59</sup>
<i>Kunzea parvifolia</i>	n,q,v	25			
<i>Kunzea pomifera</i>	s,v		3		
<i>Kunzea preissiana</i>	w				FR <sup>40</sup>
<i>Kunzea recurva</i> s.l.	w		3		MS <sup>40</sup>
<i>Kunzea sulphurea</i>	w				S <sup>46</sup>
<i>Leptospermum continentale</i>	s	25			S <sup>43</sup>
<i>Leptospermum coriaceum</i>	n,s,v		3		
<i>Leptospermum ellipticum</i>	w	15			
<i>Leptospermum erubescens</i>	w				FR <sup>40</sup>
<i>Leptospermum glaucescens</i>	t	17,34			HS <sup>34</sup> , VS or MS <sup>60</sup>
<i>Leptospermum juniperinum</i>	n,s,v	13,18,19	3		
<i>Leptospermum lanigerum</i>	n,q,s,t,v	3			
<i>Leptospermum myrsinoides</i>	n,s,v	4,25			S <sup>43</sup>
<i>Leptospermum scoparium</i>	n,t,v	8,17			FR <sup>22</sup>
<i>Leptospermum trinervium</i>	n,q,v				LS <sup>42</sup>
<i>Lophostemon confertus</i>	n,q	2			
<i>Melaleuca cuticularis</i>	w				FR <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Melaleuca diosmifolia</i>	w				FR <sup>50</sup>
<i>Melaleuca elliptica</i>	w		3		
<i>Melaleuca gibbosa</i>	s,t,v,w	17			
<i>Melaleuca holosericea</i>	w				FR <sup>56</sup>
<i>Melaleuca incana</i>	w		3		
<i>Melaleuca lanceolata</i>	w				FR <sup>50</sup>
<i>Melaleuca laxiflora</i>	w				FR <sup>53</sup>
<i>Melaleuca macronychia</i>	w		3		FR <sup>50</sup>
<i>Melaleuca microphylla</i>	w				FR <sup>59</sup>
<i>Melaleuca nesophila</i>	w				FR <sup>50</sup>
<i>Melaleuca pentagona</i> s.l.	w		3		FR <sup>50</sup>
<i>Melaleuca pritzellii</i>	w				LS <sup>58</sup>
<i>Melaleuca pulchella</i>	w				FR <sup>50</sup>
<i>Melaleuca scabra</i>	w	37			S <sup>37,46</sup>
<i>Melaleuca seriata</i>	w				FR <sup>40</sup>
<i>Melaleuca spathulata</i>	w				FR <sup>50</sup>
<i>Melaleuca squamea</i>	n,s,t,v	17,34	3	33	HS <sup>34</sup>
<i>Melaleuca squarrosa</i>	n,s,t,v	17,25			S <sup>43</sup> , FR <sup>21</sup>
<i>Melaleuca suberosa</i>	w				FR <sup>40</sup>
<i>Melaleuca subfalcata</i>	w				S <sup>40</sup>
<i>Melaleuca thymoides</i>	w	14,24			MS <sup>59</sup>
<i>Melaleuca uncinata</i>	n,nt,s,v,w		3		
<i>Melaleuca violacea</i>	w				FR <sup>50</sup>
<i>Pericalymma ellipticum</i>	w	14,24	37,53		S <sup>37,53</sup>
<i>Phymatocarpus maxwellii</i>	w		3		
<i>Regelia inops</i>	w		3		LS <sup>40</sup>
<i>Rhodamnia blairiana</i>	q				S <sup>36</sup> FR <sup>36</sup>
<i>Rhodamnia sessiliflora</i>	q				FR <sup>36</sup>
<i>Scholtzia involucrata</i>	w	37			S <sup>46</sup>
<i>Syzygium cormiflorum</i>	q				FR <sup>36</sup>
<i>Syzygium erythrodoxum</i>	q			16	
<i>Syzygium kuranda</i>	q				S <sup>36</sup>
<i>Syzygium johnsonii</i>	q				FR <sup>36</sup>
<i>Syzygium wesa</i>	q			16	FR <sup>36</sup>
<i>Thryptomene calycina</i>	s,v	2	44		MS <sup>44</sup>
<i>Thryptomene micrantha</i>	s,t,v	34			HS <sup>60</sup>
<i>Thryptomene saxicola</i>	w				S <sup>37</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Verticordia carinata</i> V	w	58			LS-MS <sup>58</sup>
<i>Verticordia chrysantha</i>	w		3		
<i>Verticordia densiflora</i>	w	14,37, 49,53			S <sup>37,49,53</sup>
<i>Verticordia habrantha</i>	w				FR <sup>40</sup>
<i>Verticordia huegelii</i>	w	14		37, 53	S <sup>37,53</sup>
<i>Verticordia nitens</i>	w	37			S <sup>37,46</sup>
<i>Verticordia plumosa</i>	w	14	3		
<i>Waterhousia unipunctata</i>	q				FR <sup>36</sup>
<i>Wehlia coarctata</i>	w		3		
OCHNACEAE					
<i>Brackenridgea nitida</i> subsp. <i>australiana</i>	q	16			FR <sup>36</sup>
OLACACEAE					
<i>Olax benthamiana</i>	w				Q <sup>21</sup>
<i>Olax phyllanthi</i>	w				FR <sup>40</sup>
OLEACEAE					
<i>Chionanthus axillaris</i>	q				FR <sup>36</sup>
ORCHIDACEAE					
<i>Caladenia flava</i>	w				FR <sup>21,59</sup>
<i>Cryptostylis ovata</i>	w				FR <sup>40</sup>
<i>Drakea confluens</i> E	w				FR <sup>58</sup>
<i>Elythranthera brunonis</i>	w				FR <sup>21,59</sup>
<i>Eriochilus dilatatus</i>	w				FR <sup>40</sup>
<i>Leporella fimbriata</i>	s,t,vw				FR <sup>40</sup>
<i>Mecopodium parvifolium</i>	w				FR <sup>21</sup>
<i>Pterostylis concinna</i>	n,s,t,v	25			S <sup>43</sup>
<i>Pterostylis vittata</i>	s,v,t,w				FR <sup>59</sup>
<i>Pterostylis aff. nana</i>	w				FR <sup>59</sup>
<i>Pyrorchis nigricans</i>	w				FR <sup>21</sup>
<i>Thelymitra crinita</i>	w				FR <sup>21</sup>
<i>Thelymitra pauciflora</i> s.l.	n,q,s,t,v,w				FR <sup>59</sup>
PHORMIACEAE					
<i>Agrostocrinum scabrum</i>	w			21	FR <sup>59</sup> S <sup>21</sup>
<i>Dianella longifolia</i> s.l.	n,nt,q,s,t,v,w	25		9,34	
<i>Dianella revoluta</i> s.l.	n,q,s,t,v,w	14,24			S <sup>24</sup>
<i>Dianella tasmanica</i>	n,t,v	8,17			
<i>Johnsonia lupulina</i> s.l.	w				FR <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Tricoryne elatior</i>	n,nt,q,s,t,v,w				FR <sup>21</sup>
<b>PHYLLOCLADACEAE</b>					
<i>Phyllocladus aspleniifolius</i>	t	17,34	3		
<b>PITTOSPORACEAE</b>					
<i>Billardiera drummondiana</i>	w				FR <sup>24</sup>
<i>Billardiera fraseri</i>	w				FR <sup>21</sup>
<i>Pittosporum phillyreoides</i>	n,nt,q,s,v,w				FR <sup>50</sup>
<i>Rhytidosporum procumbens</i>	n,q,t,v		2		
<i>Sollya drummondii</i>	w				FR <sup>58,59</sup>
<b>POACEAE</b>					
<i>Agrostis aemula</i>	n,q,s,t,v,w				FR <sup>59</sup>
<i>Amphipogon amphipogonoides</i>	w				Q <sup>21</sup>
<i>Amphipogon laguroides</i> s.l.	w				FR <sup>59</sup>
<i>Anisopogon avenaceus</i>	n,v				FR <sup>42</sup>
<i>Austrostipa compressa</i>	w				FR <sup>40</sup>
<i>Austrostipa flavescens</i>	n,s,t,v,w				LS <sup>40</sup>
<i>Deyeuxia drummondii</i> E	w				FR <sup>58</sup>
<i>Entolasia stricta</i>	n,q				FR <sup>42</sup>
<i>Neurachne alopecuroidea</i>	s,v,w				FR <sup>21</sup>
<i>Poa poiformis</i>	n,q,s,v,w				FR <sup>54</sup>
<i>Poa porphyroclados</i>	w				FR <sup>59</sup>
<i>Poa sieberiana</i>	n,q,t,v		44		FR <sup>44</sup>
<i>Tetrarrhena juncea</i>	n,q,t,v				FR <sup>22</sup>
<i>Tetrarrhena laevis</i>	w			21	FR <sup>59</sup> ,Q <sup>21</sup>
<i>Themeda triandra</i>	n,nt,q,s,t,v,w	29			S <sup>43</sup>
<b>PODOCARPACEAE</b>					
<i>Microstrobos fitzgeraldii</i>	n		3		
<i>Podocarpus druynianus</i>	w	15,46, 53	3		S <sup>46,53</sup>
<i>Podocarpus lawrencei</i>	n,t,v	17	3		
<i>Prumnopitys amara</i>	q			16	
<b>POLYGALACEAE</b>					
<i>Comesperma calymega</i>	s,t,v,w	40			FR <sup>21</sup>
<i>Comesperma confertum</i>	w	40			FR <sup>40</sup>
<i>Comesperma ericinum</i> s. l.	n,q,t,v				
<i>Comesperma virgatum</i>	w			21	Q <sup>21</sup> ,FR <sup>24</sup>
<b>PROTEACEAE</b>					
<i>Adenanthes apiculatus</i>	w		3		

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Adenanthes barbiger</i>	w	14,28			S <sup>46</sup> ,HS <sup>14</sup>
<i>Adenanthes cacomorphus</i>	w				HS <sup>41</sup>
<i>Adenanthes cuneatus</i>	w				S <sup>46,51</sup> HS <sup>40</sup> LS <sup>59</sup>
<i>Adenanthes cunninghamii</i> E	w				S <sup>49,52</sup> ,MS <sup>40,41</sup>
<i>Adenanthes cygnorum</i> s.l.	w				S <sup>37,46</sup> ,HS <sup>40</sup>
<i>Adenanthes cygnorum</i> subsp. <i>chamaephyton</i>	w				HS <sup>41</sup>
<i>Adenanthes detmoldii</i> s.l.	w				S <sup>46</sup> ,HS <sup>41</sup>
<i>Adenanthes dobagii</i> E	w				S <sup>49</sup> ,HS <sup>41</sup>
<i>Adenanthes ellipticus</i> V	w				S <sup>52</sup> ,HS <sup>41,59</sup>
<i>Adenanthes eyrei</i> E	w				HS <sup>41</sup>
<i>Adenanthes filifolius</i>	w	58			S <sup>56</sup> ,MS-HS <sup>40,58,59</sup>
<i>Adenanthes glabrescens</i> subsp. <i>exasperatus</i>	w				HS <sup>41</sup>
<i>Adenanthes gracilipes</i>	w				HS <sup>41</sup>
<i>Adenanthes ileticos</i>	w				S <sup>52</sup> ,HS <sup>41</sup>
<i>Adenanthes labillardierei</i>	w				HS <sup>41,59</sup>
<i>Adenanthes linearis</i>	w				HS <sup>40,41</sup>
<i>Adenanthes macropodiana</i>	s	19			
<i>Adenanthes meisneri</i>	w				S <sup>46</sup>
<i>Adenanthes obovatus</i>	w	2,14,15, 24			S <sup>37,46,53</sup> ,HS <sup>40,59</sup>
<i>Adenanthes oreophilus</i>	w				S <sup>56</sup> ,HS <sup>40,59</sup>
<i>Adenanthes pungens</i> subsp. <i>effusus</i> E	w				S <sup>49</sup> ,HS <sup>41</sup>
<i>Adenanthes pungens</i> subsp. <i>pungens</i> V	w				S <sup>49</sup> ,HS <sup>41</sup>
<i>Adenanthes sericeus</i>	w				S <sup>46</sup>
<i>Adenanthes terminalis</i>	s,v				S <sup>57</sup>
<i>Adenanthes velutinus</i> E	w				HS <sup>41</sup>
<i>Agastachys odorata</i>	t	17,34			HS <sup>34</sup>
<i>Banksia aculeata</i>	w				MS <sup>40</sup> ,S <sup>58</sup>
<i>Banksia ashbyi</i>	w	47	47		S <sup>47</sup>
<i>Banksia aspleniifolia</i>	q		3		
<i>Banksia attenuata</i>	w	14,15,24, 37	3	23*	S <sup>46,53</sup> ,HS <sup>23,40</sup>
<i>Banksia audax</i>	w			23*	
<i>Banksia baueri</i>	w	46, 47		23*	HS <sup>59</sup> ,S <sup>23,46,47</sup> , MS <sup>40</sup>
<i>Banksia baxteri</i>	w	46, 47		23*	S <sup>46,47</sup> ,HS <sup>40,59</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Banksia benthamiana</i>	w			23*	HS <sup>23,41</sup>
<i>Banksia brownii E</i>	w	58	49, 47, 52	23*	S <sup>49,47,52</sup> HS <sup>23,40,41,58,59</sup>
<i>Banksia burdettii</i>	w		47	23*	S <sup>47</sup>
<i>Banksia caleyi</i>	w		47, 3	23*	S <sup>47</sup> ,MS <sup>40</sup>
<i>Banksia candolleana</i>	w		47	23*	S <sup>47</sup>
<i>Banksia chamaephyton</i>	w				HS <sup>41</sup>
<i>Banksia coccinea</i>	w		46, 47	23*	S <sup>46,47</sup> ,HS <sup>23,40,59</sup>
<i>Banksia cuneata E</i>	w		52	23*	S <sup>52</sup> ,HS <sup>23,40,41</sup>
<i>Banksia dryandroides</i>	w			23*	HS <sup>40</sup>
<i>Banksia elderiana</i>	w		3	23*	
<i>Banksia elegans</i>	w		3		HS <sup>41</sup>
<i>Banksia epica</i>	w				HS <sup>41</sup>
<i>Banksia ericifolia</i>	n	3,6		23*	SV <sup>42</sup>
<i>Banksia gardneri</i> var. <i>brevidentata</i>	w				MS <sup>40</sup>
<i>Banksia gardneri</i> var. <i>gardneri</i>	w				MS <sup>40</sup>
<i>Banksia gardneri</i> var. <i>hiemalis</i>	w			23*	
<i>Banksia goodii V</i>	w				S <sup>49,52</sup> ,MS <sup>41</sup>
<i>Banksia grandis</i>	w	14,15,21, 24,28	3	21,23*	S <sup>24,46,47,53</sup> HS <sup>14,23,40,59</sup>
<i>Banksia hookeriana</i>	w		47,	23□	S <sup>47</sup> ,HS <sup>23</sup>
<i>Banksia ilicifolia</i>	w	14,15,24, 37,46,53			S <sup>46,53</sup> ,HS <sup>23,40</sup>
<i>Banksia integrifolia</i> s.l.	n,q,t,v	2		23*	
<i>Banksia laevigata</i>	w			23*	S <sup>23</sup> ,HS <sup>41</sup>
<i>Banksia laricina</i>	w	37	47,	23*	S <sup>47</sup> ,HS <sup>23,40</sup>
<i>Banksia lemanniana</i>	w		3	23*	
<i>Banksia lindleyana</i>	w		47,	23*	S <sup>47</sup>
<i>Banksia littoralis</i>	w	14,15	46, 53	23*	S <sup>46,53</sup> ,HS <sup>40</sup>
<i>Banksia lullfitzii</i>	w			23*	HS <sup>41</sup>
<i>Banksia marginata</i>	n,s,t,v	2,4,8,13, 17,18,19, 25,29,34		23*	S <sup>43</sup>
<i>Banksia media</i>	w		46	23*	S <sup>46</sup>
<i>Banksia meisneri</i> var. <i>ascendens</i>	w				HS <sup>41</sup>
<i>Banksia menziesii</i>	w	14,15,24, 37		23*	S <sup>46,47,53</sup> ,HS <sup>23,40</sup>
<i>Banksia micrantha</i>	w				MS <sup>40,41</sup>
<i>Banksia nutans</i>	w			23*	S <sup>46, 47</sup> ,HS <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Banksia oblongifolia</i>	n,q			23*	
<i>Banksia occidentalis</i> s.l.	w		3	23*	S <sup>40,46,47</sup> ,HS <sup>40,41</sup>
<i>Banksia oligantha</i> E	w				S <sup>49,52</sup> ,HS <sup>41</sup>
<i>Banksia oreophila</i>	w	58			HS <sup>40,58,59</sup>
<i>Banksia ornata</i>	s,v		3	23*	HS <sup>23</sup>
<i>Banksia paludosa</i> subsp. <i>paludosa</i>	n			23*	SV <sup>42</sup>
<i>Banksia petiolaris</i>	w			23*	
<i>Banksia pilostylis</i>	w			23*	
<i>Banksia preamorsa</i>	w			23*	S <sup>46</sup>
<i>Banksia prionotes</i>	w			23*	S <sup>46,47</sup> ,HS <sup>23,40</sup>
<i>Banksia pulchella</i>	w			23*	HS <sup>23</sup>
<i>Banksia quercifolia</i>	w	15		23*	S <sup>46,51</sup> ,HS <sup>40,59</sup>
<i>Banksia repens</i>	w			23*	MS <sup>40</sup>
<i>Banksia saxicola</i>	v		44		HS <sup>44</sup>
<i>Banksia scabrella</i>	w				HS <sup>41</sup>
<i>Banksia sceptrum</i>	w			23*	HS <sup>23</sup>
<i>Banksia seminuda</i>	w				S <sup>46</sup> ,HS <sup>40</sup>
<i>Banksia serrata</i>	n,s,t,v	2,17,25, 34			S <sup>43</sup> ,SV <sup>42,60</sup>
<i>Banksia solandri</i>	w	58			MS <sup>40</sup> , HS <sup>41,58,59</sup>
<i>Banksia speciosa</i>	w		3	23*	S <sup>46,47</sup> ,HS <sup>23,40</sup>
<i>Banksia sphaerocarpa</i> s.l.	w		3	23*	S <sup>46,47</sup> MS <sup>59</sup> , HS <sup>23,40,41,58</sup>
<i>Banksia spinulosa</i> var. <i>cunninghamii</i>	n,q,v	13,25			S <sup>43</sup>
<i>Banksia telmatiae</i> a	w	37			S <sup>37,46</sup>
<i>Banksia tricuspis</i>	w				S <sup>52</sup> ,HS <sup>41</sup>
<i>Banksia verticillata</i> V	w			23*	S <sup>52</sup> ,HS <sup>40,41,58,59</sup>
<i>Banksia victoriae</i>	w			23*	HS <sup>23</sup>
<i>Banksia violacea</i>	w			23*	HS <sup>40,59</sup>
<i>Buckinghamia celsissima</i>	q			16	
<i>Cardwellia sublimis</i>	q				S <sup>36</sup>
<i>Carnarvonia araliifolia</i> var. <i>montana</i>	q	16			
<i>Cennarrhenes nitida</i>	t	17,34			HS <sup>34</sup>
<i>Conospermum caeruleum</i> s.l.	w				MS <sup>59</sup>
<i>Conospermum coerulescens</i> subsp. <i>dorrienii</i>	w				MS <sup>59</sup>
<i>Conospermum eatoniae</i>	w				LS <sup>41</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Conospermum hookeri</i>	t	34			MS <sup>60</sup>
<i>Conospermum mitchellii</i>	v	25,29			S <sup>43</sup>
<i>Conospermum scaposum</i>	w				LS <sup>41</sup>
<i>Conospermum spectabile</i>	w				S <sup>58</sup>
<i>Conospermum stoechadis</i>	w				S <sup>37,46</sup>
<i>Conospermum todii</i>	w				S <sup>52</sup>
<i>Conospermum triplinervium</i>	w	37		37	S <sup>37</sup>
<i>Conospermum undulatum</i>	w				LS <sup>41</sup>
<i>Darlingia darlingiana</i>	q	16			S <sup>36</sup>
<i>Dryandra acanthopoda</i>	w				HS <sup>41</sup>
<i>Dryandra anatona</i> E	w	58			S <sup>49</sup> ,HS <sup>41,58</sup>
<i>Dryandra arctotidis</i>	w				S <sup>46</sup> ,MS <sup>40</sup>
<i>Dryandra armata</i> s.l.	w	14			HS <sup>59</sup> ,S <sup>46</sup>
<i>Dryandra baxteri</i>	w				HS <sup>59</sup>
<i>Dryandra bipinnatifida</i>	w				S <sup>46</sup>
<i>Dryandra calophylla</i>	w				MS <sup>58</sup>
<i>Dryandra carduacea</i>	w	14			
<i>Dryandra cirsoides</i>	w				MS <sup>40</sup>
<i>Dryandra comosa</i>	w				HS <sup>41</sup>
<i>Dryandra concinna</i>	w	58			HS <sup>58,59</sup>
<i>Dryandra conferta</i> var. <i>parva</i>	w				HS <sup>58</sup>
<i>Dryandra cynaroides</i>	w				HS <sup>41</sup>
<i>Dryandra erythrocephala</i> var. <i>inopinata</i>	w				HS <sup>41</sup>
<i>Dryandra falcata</i>	w				HS <sup>40</sup>
<i>Dryandra ferruginea</i> subsp. <i>pumila</i>	w				HS <sup>58</sup>
<i>Dryandra foliolata</i>	w	58			HS <sup>58,59</sup>
<i>Dryandra foliosissima</i>	w				HS <sup>41</sup>
<i>Dryandra formosa</i>	w	46, 51, 58			S <sup>46,51</sup> ,MS-HS <sup>40,58,59</sup>
<i>Dryandra fraseri</i> var. <i>oxycedra</i>	w				HS <sup>41</sup>
<i>Dryandra hirsuta</i>	w	58			MS <sup>40,58</sup>
<i>Dryandra horrida</i>	w				HS <sup>41</sup>
<i>Dryandra idiogenes</i>	w				HS <sup>41</sup>
<i>Dryandra ionthocarpa</i> E	w				HS <sup>41</sup>
<i>Dryandra kippistiana</i> var. <i>paenepeccata</i>	w				HS <sup>41</sup>
<i>Dryandra lepidorhiza</i>	w				HS <sup>41</sup>
<i>Dryandra lindleyana</i> s.l.	w				S <sup>46</sup> ,SP <sup>21</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Dryandra mimica</i> E	w				S <sup>52</sup> ,HS <sup>41</sup>
<i>Dryandra montana</i> E	w	58			S <sup>49</sup> ,HS <sup>41,58,59</sup>
<i>Dryandra mucronulata</i>	w				HS <sup>40</sup>
<i>Dryandra nivea</i> s.l.	w	14,28			S <sup>37,46,53</sup> MS <sup>40,59</sup>
<i>Dryandra octotriginta</i>	w				HS <sup>41</sup>
<i>Dryandra plumosa</i> subsp. <i>denticulata</i>	w				MS-HS <sup>40,58,59</sup>
<i>Dryandra polycephala</i>	w				MS <sup>41</sup>
<i>Dryandra porrecta</i>	w				HS <sup>41</sup>
<i>Dryandra praemorsa</i>	w		3		
<i>Dryandra preissii</i>	w				HS <sup>41</sup>
<i>Dryandra pseudoplumosa</i>	w				HS <sup>58</sup> ,MS <sup>40</sup>
<i>Dryandra pteridifolia</i>	w				S <sup>46</sup> ,HS <sup>40</sup>
<i>Dryandra pulchella</i>	w				HS <sup>41</sup>
<i>Dryandra quercifolia</i>	w				HS <sup>40,59</sup>
<i>Dryandra sclerophylla</i>	w				HS <sup>41</sup>
<i>Dryandra seneciifolia</i>	w				S <sup>46</sup> ,MS-HS <sup>41,58</sup>
<i>Dryandra serra</i>	w				S <sup>46</sup> ,HS <sup>41,58</sup>
<i>Dryandra serratuloides</i> subsp. <i>perissa</i>	w				S <sup>49</sup>
<i>Dryandra serratuloides</i> s.l.	w				S <sup>49,52</sup> ,HS <sup>41</sup>
<i>Dryandra sessilis</i>	w	14,28			SP <sup>21</sup> ,S <sup>46</sup> ,MS <sup>40</sup>
<i>Dryandra shanklandiorum</i>	w				HS <sup>41</sup>
<i>Dryandra speciosa</i>	w				HS <sup>41</sup>
<i>Dryandra squarrosa</i> subsp. <i>argillacea</i>	w				S <sup>49</sup>
<i>Dryandra squarrosa</i> s.l.	w				S <sup>46</sup>
<i>Dryandra subpinnatifida</i>	w				HS <sup>41</sup>
<i>Dryandra tenuifolia</i>	w				HS <sup>40</sup>
<i>Dryandra tortifolia</i>	w				HS <sup>41</sup>
<i>Dryandra trifontinalis</i>	w				HS <sup>41</sup>
<i>Dryandra wonganensis</i>	w				HS <sup>41</sup>
<i>Franklandia fucifolia</i>	w				MS <sup>40</sup>
<i>Franklandia triaristata</i>	w				LS <sup>41</sup>
<i>Grevillea acrobotrys</i> subsp. <i>uniformis</i>	w				LS <sup>41</sup>
<i>Grevillea alpina</i>	n,v	2,25,29			S <sup>43</sup>
<i>Grevillea aneura</i>	w				LS <sup>41</sup>
<i>Grevillea annulifera</i>	w				LS <sup>41</sup>
<i>Grevillea aquifolium</i>	s,v	25,29			S <sup>43</sup>
<i>Grevillea asparagoides</i>	w				LS <sup>41</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Grevillea astericosa</i>	w				LS <sup>41</sup>
<i>Grevillea batrachiooides E</i>	w				LS <sup>41</sup>
<i>Grevillea baxteri</i>	w				LS <sup>41</sup>
<i>Grevillea brachystylis</i> subsp. <i>australis</i> V	w				LS <sup>41</sup>
<i>Grevillea brachystylis</i> subsp. <i>brachystylis</i>	w				LS <sup>41</sup>
<i>Grevillea bracteosa</i>	w				LS <sup>41</sup>
<i>Grevillea calliantha E</i>	w				S <sup>52</sup> , LS <sup>41</sup>
<i>Grevillea candicans</i>	w				LS <sup>41</sup>
<i>Grevillea christinae E</i>	w				LS <sup>41</sup>
<i>Grevillea chrysophaea</i>	v	32			HS <sup>32</sup>
<i>Grevillea cirsifolia</i>	w				S <sup>52</sup>
<i>Grevillea confertifolia</i>	v		44		MS <sup>44</sup>
<i>Grevillea costata</i>	w				LS <sup>41</sup>
<i>Grevillea crowleyae</i>	w				LS <sup>41</sup>
<i>Grevillea curviloba</i> s.l.	w				LS <sup>41</sup>
<i>Grevillea decora</i>	q		3		
<i>Grevillea depauperata</i>	w				LS <sup>41</sup>
<i>Grevillea donaldiana</i>	w				LS <sup>41</sup>
<i>Grevillea dryandroides</i> s.l.	w				LS <sup>41</sup>
<i>Grevillea erectiloba</i>	w				LS <sup>41</sup>
<i>Grevillea eriobotrya</i>	w				LS <sup>41</sup>
<i>Grevillea fasciculata</i>	w				FR <sup>40</sup> , MS <sup>59</sup>
<i>Grevillea fistulosa</i>	w				LS <sup>41</sup>
<i>Grevillea flexuosa</i> V	w				LS <sup>41</sup>
<i>Grevillea fulgens</i>	w				LS <sup>41</sup>
<i>Grevillea fuscolutea</i>	w				LS <sup>41</sup>
<i>Grevillea georgeana</i>	w				LS <sup>41</sup>
<i>Grevillea glabrata</i> subsp. <i>dissectifolia</i>	w				LS <sup>41</sup>
<i>Grevillea glabrata</i> subsp. <i>ornithopoda</i>	w				LS <sup>41</sup>
<i>Grevillea globosa</i>	w				LS <sup>41</sup>
<i>Grevillea granulosa</i>	w				LS <sup>41</sup>
<i>Grevillea hookeriana</i>	w				FR <sup>50</sup>
<i>Grevillea inconspicua</i>	w				LS <sup>41</sup>
<i>Grevillea infundibularis E</i>	w				LS <sup>41</sup>
<i>Grevillea involucrata E</i>	w				LS <sup>41</sup>
<i>Grevillea irrasa</i> subsp. <i>irrasa</i>	n	13			HS <sup>22</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Grevillea juniperina</i> s.l.	n		3		
<i>Grevillea kenneallyi</i>	w				LS <sup>41</sup>
<i>Grevillea lanigera</i>	n,v		3		
<i>Grevillea latifolia</i>	w				LS <sup>41</sup>
<i>Grevillea lavandulacea</i> s.l.	s,v				S <sup>20</sup>
<i>Grevillea leptopoda</i>	w				LS <sup>41</sup>
<i>Grevillea lissopleura</i>	w				LS <sup>41</sup>
<i>Grevillea longistyla</i>	q		3		
<i>Grevillea lullfitzii</i>	w				LS <sup>41</sup>
<i>Grevillea makinsonii</i>	w				LS <sup>41</sup>
<i>Grevillea manglesioides</i> subsp. <i>papillosa</i>	w				LS <sup>41</sup>
<i>Grevillea mariottii</i>	w				LS <sup>41</sup>
<i>Grevillea maxwellii</i> E	w				LS <sup>41</sup>
<i>Grevillea microstegia</i>	v		44		LS <sup>44</sup>
<i>Grevillea miniata</i>	w				LS <sup>41</sup>
<i>Grevillea minutiflora</i>	w				LS <sup>41</sup>
<i>Grevillea miquelianiana</i> s.l.	n,v		3		
<i>Grevillea mucronulata</i>	n		3		
<i>Grevillea murex</i> E	w				LS <sup>41</sup>
<i>Grevillea nana</i> subsp. <i>abbreviata</i>	w				LS <sup>41</sup>
<i>Grevillea oleoides</i>	n				S <sup>22</sup>
<i>Grevillea olivacea</i>	w				LS <sup>41</sup>
<i>Grevillea phanerophlebia</i>	w				LS <sup>41</sup>
<i>Grevillea phillipsiana</i>	w				LS <sup>41</sup>
<i>Grevillea pieroniae</i>	w				MS <sup>58</sup>
<i>Grevillea pilosa</i> subsp. <i>dissecta</i>	w				LS <sup>41</sup>
<i>Grevillea pilulifera</i>	w				FR <sup>21</sup>
<i>Grevillea pimeleoides</i>	w				LS <sup>41</sup>
<i>Grevillea polybractea</i>	n,v		3		
<i>Grevillea prostrata</i>	w				LS <sup>41</sup>
<i>Grevillea psilantha</i>	w				LS <sup>41</sup>
<i>Grevillea quinquenervis</i>	s	19			
<i>Grevillea rogersii</i>	s	19			
<i>Grevillea rogersoniana</i>	w				LS <sup>41</sup>
<i>Grevillea rosieri</i>	w				LS <sup>41</sup>
<i>Grevillea rosmarinifolia</i> s.l.	n,s,v		3		

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Grevillea royceri</i>	w				LS <sup>41</sup>
<i>Grevillea rудis</i>	w				LS <sup>41</sup>
<i>Grevillea saccata</i> Benth.	w				S <sup>52</sup>
<i>Grevillea scabra</i>	w				LS <sup>41</sup>
<i>Grevillea scabrida</i>	w				LS <sup>41</sup>
<i>Grevillea scapigera</i> E	w				LS <sup>41</sup>
<i>Grevillea secunda</i>	w				LS <sup>41</sup>
<i>Grevillea spinosissima</i>	w				LS <sup>41</sup>
<i>Grevillea steiglitziana</i>	v	25			S <sup>43</sup>
<i>Grevillea stenostachya</i>	w				LS <sup>41</sup>
<i>Grevillea subtiliflora</i>	w				LS <sup>41</sup>
<i>Grevillea synapheae</i>	w	24			FR <sup>24</sup>
<i>Grevillea tenuiloba</i>	w				LS <sup>41</sup>
<i>Grevillea tetrapleura</i>	w				LS <sup>41</sup>
<i>Grevillea thelemanniana</i>	w				LS <sup>41</sup>
<i>Grevillea triloba</i>	w				LS <sup>41</sup>
<i>Grevillea tripartita</i>	w				MS <sup>40</sup>
<i>Grevillea victoriae</i> s.l.	n,v		3		
<i>Grevillea williamsonii</i> E	v		44		HS <sup>44</sup>
<i>Grevillea wilsonii</i>	w				FR <sup>21</sup>
<i>Grevillea wittweri</i>	w				LS <sup>41</sup>
<i>Hakea aculeata</i> V	w				LS <sup>41</sup>
<i>Hakea ambigua</i>	w				MS <sup>40,59</sup>
<i>Hakea amplexicaulis</i>	w	24			FR <sup>24</sup>
<i>Hakea bakeriana</i>	n		3		
<i>Hakea baxteri</i>	w				S <sup>46</sup> ,MS <sup>40,59</sup>
<i>Hakea bicornata</i>	w				LS <sup>41</sup>
<i>Hakea ceratophylla</i>	w				LS <sup>40,59</sup>
<i>Hakea conchifolia</i>	w		3		
<i>Hakea corymbosa</i>	w				FR <sup>40</sup>
<i>Hakea crassifolia</i>	w				MS <sup>40,59</sup>
<i>Hakea crassinervia</i>	w				LS <sup>41</sup>
<i>Hakea cucullata</i>	w				S <sup>46</sup> ,MS <sup>40,59</sup>
<i>Hakea dactyloides</i>	n,q,v				S <sup>22</sup>
<i>Hakea elliptica</i>	w				MS <sup>59</sup>
<i>Hakea flabellifolia</i>	w				S <sup>56</sup>
<i>Hakea kippistiana</i>	w		3		

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Hakea lasiantha</i>	w				MS <sup>40</sup>
<i>Hakea lasiocarpa</i>	w				MS <sup>58</sup>
<i>Hakea laurina</i>	w				LS <sup>40</sup>
<i>Hakea lemanniana</i>	w				MS <sup>40</sup>
<i>Hakea lissocarpa</i>	w	24,28			FR <sup>21,24</sup>
<i>Hakea longiflora</i>	w				LS <sup>41</sup>
<i>Hakea marginata</i>	w				HS <sup>40</sup>
<i>Hakea megalosperma</i> V	w				LS <sup>41</sup>
<i>Hakea myrtoides</i>	w				LS <sup>41</sup>
<i>Hakea neurophylla</i>	w				LS <sup>41</sup>
<i>Hakea nodosa</i>	s,t,v	25			S <sup>43</sup>
<i>Hakea oleifolia</i>	w		3		S <sup>46</sup>
<i>Hakea pendanicarpa</i> subsp. <i>crassifolia</i>	w				S <sup>40,46</sup>
<i>Hakea pendens</i>	w				LS <sup>41</sup>
<i>Hakea petiolaris</i>	w				FR <sup>50</sup>
<i>Hakea platysperma</i>	w		3		
<i>Hakea prostrata</i>	w				S <sup>46</sup>
<i>Hakea rigida</i>	w				LS <sup>41</sup>
<i>Hakea rubrifolia</i>	w				LS <sup>40</sup>
<i>Hakea ruscifolia</i>	w	14			FR <sup>21</sup> ,LS <sup>40</sup>
<i>Hakea salicifolia</i>	n,q				
<i>Hakea sericea</i>	n,t,v	2,17,25			
<i>Hakea spathulata</i>	w				LS <sup>41</sup>
<i>Hakea stenocarpa</i>	w				FR <sup>21</sup>
<i>Hakea trifurcata</i>	w				S <sup>46</sup> ,LS <sup>40</sup>
<i>Hakea tuberculata</i>	w				LS <sup>58</sup>
<i>Hakea ulicina</i>	n,t,v	25,34		9	S <sup>43</sup>
<i>Hakea undulata</i>	w		3		S <sup>46</sup> ,MS <sup>40</sup>
<i>Hakea varia</i>	w				LS <sup>40,59</sup>
<i>Hakea victoria</i>	w		3		LS <sup>40,59</sup>
<i>Isopogon alcicornis</i>	w				HS <sup>41</sup>
<i>Isopogon anemonifolius</i>	n				SV <sup>42</sup>
<i>Isopogon asper</i>	w		3		
<i>Isopogon attenuatus</i>	w	15		54	LS <sup>59</sup> ,S <sup>54</sup>
<i>Isopogon axillaris</i>	w				S <sup>46</sup> ,HS <sup>40</sup>
<i>Isopogon baxteri</i>	w				HS <sup>58</sup> ,LS <sup>59</sup> MS <sup>40</sup>
<i>Isopogon buxifolius</i> var. <i>obovatus</i>	w				HS <sup>40,59</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Isopogon ceratophyllum</i>	s,t,v	18,25,29, 34			S <sup>43</sup>
<i>Isopogon drummondii</i>	w				HS <sup>41</sup>
<i>Isopogon formosus</i>	w	15,46			HS <sup>59</sup> ,S <sup>46</sup> ,LS <sup>40</sup>
<i>Isopogon heterophyllum</i>	w				MS <sup>40</sup>
<i>Isopogon latifolius</i>	w	58			HS <sup>58</sup> ,MS <sup>40</sup>
<i>Isopogon petiolaris</i>	n,q		3		
<i>Isopogon polycephalus</i>	w		3		
<i>Isopogon scabriusculus</i>	w				MS <sup>40</sup>
<i>Isopogon sphaerocephalus</i>	w	14,28			S <sup>24,46,48</sup> MS <sup>40,59</sup>
<i>Isopogon teretifolius</i> var. <i>petrophiloides</i>	w				MS <sup>40</sup>
<i>Isopogon tridens</i>	w				HS <sup>41</sup>
<i>Isopogon trilobus</i>	w				HS <sup>40</sup>
<i>Isopogon tripartitus</i>	w				MS <sup>40</sup>
<i>Isopogon uncinatus</i> E	w	58			S <sup>49,52</sup> ,HS <sup>41,58</sup>
<i>Lambertia echinata</i> subsp. <i>echinata</i> E	w				S <sup>46,49,52</sup> ,HS <sup>41,58</sup>
<i>Lambertia echinata</i> subsp. <i>occidentalis</i>	w				S <sup>49</sup>
<i>Lambertia ericifolia</i>	w				HS <sup>58</sup> ,MS <sup>40</sup>
<i>Lambertia fairallii</i> E	w	58			S <sup>49,52</sup> ,HS <sup>41,58,59</sup>
<i>Lambertia formosa</i>	n				SV <sup>42</sup>
<i>Lambertia inermis</i> s.l.	w				S <sup>56</sup> ,HS <sup>40,59</sup>
<i>Lambertia multiflora</i>	w				S <sup>56</sup>
<i>Lambertia orbifolia</i> E	w	58			S <sup>49,52</sup> ,HS <sup>40,41,58</sup>
<i>Lambertia rariflora</i> s.l.	w				LS <sup>41</sup>
<i>Lambertia uniflora</i>	w				HS <sup>40,59</sup>
<i>Lomatia fraseri</i>	n,v	13			
<i>Lomatia fraxinifolia</i>	q				S <sup>36</sup>
<i>Lomatia ilicifolia</i>	n,q,v	25			S <sup>43</sup>
<i>Lomatia tasmanica</i> CE	t			34	S <sup>61</sup>
<i>Opistholepis heterophylla</i>	q				S <sup>36</sup>
<i>Orites diversifolia</i>	t	17			
<i>Persoonia baeckeoides</i>	w				LS <sup>41</sup>
<i>Persoonia brachystylis</i>	w				LS <sup>41</sup>
<i>Persoonia brevirhachis</i>	w				LS <sup>41</sup>
<i>Persoonia chapmaniana</i>	w				LS <sup>41</sup>
<i>Persoonia cornifolia</i>	n,q				HS <sup>22</sup>
<i>Persoonia elliptica</i>	w				S <sup>46</sup> ,HS <sup>22</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Persoonia gunnii</i>	t	17			
<i>Persoonia hakeiformis</i>	w				LS <sup>41</sup>
<i>Persoonia juniperina</i>	s,t,v	17,25,29			S <sup>43</sup> , S <sup>60</sup>
<i>Persoonia kararae</i>	w				LS <sup>41</sup>
<i>Persoonia leucopogon</i>	w				LS <sup>41</sup>
<i>Persoonia levigata</i>	n,v	3,6			FR <sup>22</sup>
<i>Persoonia linearis</i>	n,q,v				FR <sup>22</sup>
<i>Persoonia longifolia</i>	w	14,15,28			S <sup>46,53,54</sup> HS <sup>21,40,59</sup>
<i>Persoonia micranthera</i> E	w	58			HS <sup>58,59</sup> ,LS <sup>41</sup>
<i>Persoonia microcarpa</i>	w				LS <sup>40</sup>
<i>Persoonia muelleri</i> var. <i>densifolia</i>	t			9	HS <sup>9</sup>
<i>Persoonia papillosa</i>	w				LS <sup>41</sup>
<i>Persoonia rufa</i>	w				LS <sup>41</sup>
<i>Persoonia scabra</i>	w				LS <sup>41</sup>
<i>Persoonia silvatica</i>	n,v				S <sup>22</sup>
<i>Persoonia sulcata</i>	w				LS <sup>41</sup>
<i>Persoonia trinervis</i>	w				MS <sup>40</sup>
<i>Petrophile anceps</i>	w				HS <sup>58</sup>
<i>Petrophile biloba</i>	w		53		S <sup>53</sup>
<i>Petrophile biternata</i>	w				MS <sup>41</sup>
<i>Petrophile canescens</i>	n,q				
<i>Petrophile crispata</i>	w				MS <sup>41</sup>
<i>Petrophile divaricata</i>	w				HS <sup>40,59</sup>
<i>Petrophile diversifolia</i>	w				S <sup>46</sup> ,HS <sup>40,59</sup>
<i>Petrophile drummondii</i>	w	37			S <sup>37</sup>
<i>Petrophile ericifolia</i>	w				HS <sup>40</sup>
<i>Petrophile fastigiata</i>	w		3		
<i>Petrophile incurvata</i>	w				MS <sup>41</sup>
<i>Petrophile linearis</i>	w	14,24			S <sup>37,46</sup>
<i>Petrophile longifolia</i>	w				HS <sup>40</sup>
<i>Petrophile media</i>	w				HS <sup>40</sup>
<i>Petrophile multisecta</i>	s	19			
<i>Petrophile plumosa</i>	w				MS <sup>41</sup>
<i>Petrophile pulchella</i>	n	1			
<i>Petrophile seminuda</i>	w		3		MS <sup>40</sup>
<i>Petrophile serruriae</i>	w				S <sup>46</sup> ,HS <sup>40</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Petrophile sessilis</i>	n,q	3,6			
<i>Petrophile squamata</i>	w		3		S <sup>46</sup> ,HS <sup>40</sup>
<i>Petrophile striata</i>	w	14			SP <sup>21</sup>
<i>Petrophile stricta</i>	w				S <sup>37</sup>
<i>Stenocarpus sinuatus</i>	n,q				FR <sup>36</sup>
<i>Stirlingia latifolia</i>	w	14,24			S <sup>46</sup> ,MS <sup>40</sup>
<i>Stirlingia tenuifolia</i> s.l.	w				S <sup>40</sup>
<i>Stirlingia tenuifolia</i> var. <i>anethifolia</i>	w				MS <sup>40</sup>
<i>Stirlingia tenuifolia</i> var. <i>tenuifolia</i>	w				MS <sup>40</sup>
<i>Synaphea petiolaris</i> s.l.	w	24			S <sup>46</sup> ,FR <sup>24</sup>
<i>Synaphea polymorpha</i>	w				S <sup>40</sup> ,MS <sup>40,59</sup>
<i>Telopea mongaensis</i>	n		3, 12		
<i>Telopea speciosissima</i>	n		3, 12		
<i>Telopea truncata</i>	t	17			
<i>Xylomelum angustifolium</i>	w				S <sup>46</sup>
<i>Xylomelum occidentale</i>	w	15			S <sup>37,46</sup>
<b>RANUNCULACEAE</b>					
<i>Clematis pubescens</i>	w				FR <sup>24</sup> ,Q <sup>21</sup>
<b>RESTIONACEAE</b>					
<i>Anarthria gracilis</i>	w	40			FR <sup>40</sup>
<i>Anarthria prolifera</i>	w	40			FR <sup>40</sup>
<i>Anarthria scabra</i>	w	40			FR <sup>40</sup>
<i>Calorophus exsulcus</i>	w				FR <sup>21</sup>
<i>Chordifex abortivus</i> E	w	58			FR <sup>58</sup>
<i>Chordifex isomorphus</i>	w				FR <sup>58</sup>
<i>Chordifex monocephalus</i>	w	17		33	
<i>Desmocladus fasciculatus</i>	w				FR <sup>21</sup> ,LS <sup>40</sup> MS <sup>59</sup>
<i>Desmocladus flexuosa</i>	w				FR <sup>21,59</sup>
<i>Harperia confertospicata</i>	w				FR <sup>58</sup>
<i>Leptocarpus tenax</i>	w				FR <sup>40</sup>
<i>Lepyrodia scariosa</i>	n,q				FR <sup>42</sup>
<i>Restio confertospicatus</i>	w				FR <sup>40</sup>
<i>Restio laxocarya</i>	w	14			MS <sup>21</sup>
<b>RHAMNACEAE</b>					
<i>Alphitonia petriei</i>	n,q				S <sup>36</sup>
<i>Alphitonia whitei</i>	q				S <sup>36</sup>
<i>Cryptandra pumila</i>	w				LS <sup>41</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Cryptandra tomentosa</i>	s,t,v	25			S <sup>43</sup>
<i>Pomaderris intermedia</i>	n,t,v	3			
<i>Trymalium floribundum</i>	w		3		
<i>Trymalium ledifolium</i>	w	14,28			S <sup>48,53</sup> ,SP <sup>21</sup>
<b>ROSACEAE</b>					
<i>Acaena echinata</i>	n,s,t,v,w	25			
<b>RUBIACEAE</b>					
<i>Antirhea</i> sp. (Mt Lewis BG 5733)	q				S <sup>36</sup>
<i>Atractocarpus fitzalanii</i> subsp. <i>tenuipes</i>	q				FR <sup>36</sup>
<i>Opercularia echinocephala</i>	w	28			Q <sup>21</sup>
<i>Opercularia vaginata</i>	w				Q <sup>21</sup>
<i>Opercularia varia</i>	n,s,t,v	25			
<b>RUTACEAE</b>					
<i>Acronychia oblongifolia</i>	q	16			
<i>Antirhea</i> sp. (Mt Lewis BG 5733)	q				
<i>Asterolasia phebaloides</i> V	s,v		44		HS <sup>44</sup>
<i>Boronia anemonifolia</i>	n,t,v		3		
<i>Boronia baeckeacea</i>	w		3		
<i>Boronia citriodora</i>	t,v	17,34	3		HS <sup>34</sup>
<i>Boronia crenulata</i> s.l.	w	40			FR <sup>40,59</sup>
<i>Boronia crenulata</i> subsp. <i>viminea</i>	w		3		
<i>Boronia fastigiata</i>	w			21	S <sup>21</sup>
<i>Boronia microphylla</i>	n,q				
<i>Boronia nana</i> var. <i>hyssopifolia</i>	v	25			
<i>Boronia nana</i> var. <i>nana</i>	v	25			
<i>Boronia parviflora</i>	n,q,s,t,v	17,34			HS <sup>34</sup> , LS or SV <sup>61</sup>
<i>Boronia pilosa</i> s.l.	s,t,v	17,34	3		HS <sup>34</sup> , MS <sup>61</sup>
<i>Boronia revoluta</i>	w				S <sup>49</sup>
<i>Boronia spathulata</i>	w				FR <sup>24</sup>
<i>Brombya platynema</i>	q				FR <sup>36</sup>
<i>Correa decumbens</i>	q		3		
<i>Correa pulchella</i>	s	20			
<i>Correa reflexa</i>	s	17,20,25, 29			S <sup>43</sup>
<i>Crowea angustifolia</i> s.l.	w				S <sup>46</sup>
<i>Crowea angustifolia</i> var. <i>platyphylla</i>	w		3		
<i>Crowea exalata</i>	n,v		3		

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Crowea saligna</i>	n,q		3		
<i>Flindersia bourjotiana</i>	q				S <sup>36</sup>
<i>Flindersia brayleyana</i>	q			16	
<i>Flindersia pimenteliana</i>	q				FR <sup>36</sup>
<i>Leionema phylicifolium</i>	n,v		3		
<i>Leionema ralstonii</i> V	n		22		MS <sup>22</sup>
<i>Melicope elleryana</i>	q				FR <sup>36</sup>
<i>Muiriantha hassellii</i>	w				FR <sup>58</sup>
<i>Nematolepis squamea</i> s.l.	n,q,t,v	2,8,17,34			HS <sup>34</sup>
<i>Phebalium daviesii</i>	t			9	HS <sup>9</sup>
<i>Philoteca myoporoides</i>	n,q,v		3		
<i>Philoteca spicata</i>	w	24	3		FR <sup>21,24</sup>
<i>Philoteca virgata</i>	n,t,v	17			
<i>Zieria laevigata</i> s.l.	n,q		3		
<b>SANTALACEAE</b>					
<i>Exocarpus cupressiformis</i>	n,q,s,t,v	3			
<i>Leptomeria cunninghamii</i>	w	24			FR <sup>24</sup> ,Q <sup>21</sup>
<i>Leptomeria eriocoides</i>	w				FR <sup>40</sup>
<b>SAPINDACEAE</b>					
<i>Dodonaea boroniifolia</i>	n,q,v		3		
<i>Dodonaea viscosa</i> s.l.	n,nt,q,s,t,v,w	3			FR <sup>50</sup>
<i>Jagera pseudorhus</i>	q			16	
<i>Mischocarpus macrocarpus</i>	q				FR <sup>36</sup>
<b>SAPOTACEAE</b>					
<i>Pouteria brownlessiana</i>	q				S <sup>36</sup>
<i>Pouteria euphlebia</i>	q				FR <sup>36</sup>
<i>Pouteria papyracea</i>	q				FR <sup>36</sup>
<i>Pouteria pearsoniorum</i>	q				FR <sup>36</sup>
<b>SELAGINELLACEAE</b>					
<i>Selaginella uliginosa</i>	n,q,t,v	25			S <sup>43</sup>
<b>SOLANACEAE</b>					
<i>Anthocercis racemosa</i>	w		3		
<i>Solanum oldfieldii</i>	w		3		
<b>STACKHOUSIACEAE</b>					
<i>Tripterococcus brunonis</i>	w				FR <sup>21</sup>
<b>STERCULIACEAE</b>					
<i>Argyrodendron actinophyllum</i>	q			16	

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
Species					
<i>Franciscodendron laurifolium</i>	q				S <sup>36</sup>
<i>Lasiopetalum floribundum</i>	w	15,28			S <sup>24,48,53</sup> ,MS <sup>21</sup>
<i>Lasiopetalum glabratum</i>	w	14			
<i>Lasiopetalum membranifolium</i>	w				FR <sup>58</sup>
<i>Thomasia grandiflora</i>	w	15		37	S <sup>37</sup>
<i>Thomasia pauciflora</i>	w	15			
<i>Thomasia</i> sp. Toolbrunup	w				FR <sup>58,59</sup>
<b>STYLDIACEAE</b>					
<i>Levenhookia pusilla</i>	s,w				FR <sup>21</sup>
<i>Levenhookia stipitata</i>	s,w				FR <sup>21</sup>
<i>Stylium amoenum</i>	w	26		21	HS <sup>21</sup>
<i>Stylium brunonianum</i>	w				FR <sup>40</sup>
<i>Stylium calcaratum</i>	s,v,w				FR <sup>21</sup>
<i>Stylium graminifolium</i> s. l.	n,q,s,t,v	17,34	3	33	MS <sup>60</sup>
<i>Stylium imbricatum</i>	w				FR <sup>40,59</sup>
<i>Stylium hispidum</i>	w				FR <sup>21</sup>
<i>Stylium junceum</i>	w			21	Q <sup>21</sup>
<i>Stylium keigheryi</i>	w				FR <sup>59</sup>
<i>Stylium piliferum</i> subsp. <i>minor</i>	w				FR <sup>40</sup>
<i>Stylium repens</i>	w				FR <sup>21</sup>
<i>Stylium scandens</i>	w				FR <sup>40</sup>
<i>Stylium schoenoides</i>	w				Q <sup>21</sup>
<i>Stylium spathulatum</i>	w	51			S <sup>51</sup>
<i>Stylium spinulosum</i> subsp. <i>montanum</i>	w				FR <sup>59</sup>
<i>Stylium verticillatum</i>	w				FR <sup>40,58,59</sup>
<i>Stylium</i> sp. Stirling Range	w				FR <sup>59</sup>
<b>SYMPLOCACEAE</b>					
<i>Symplocos ampulliformis</i>	q				FR <sup>36</sup>
<i>Symplocos cochinchinensis</i> var. <i>gittonsi</i>	q				FR <sup>36</sup>
<i>Symplocos stawellii</i>	q	16			
<b>TAXODIACEAE</b>					
<i>Athrotaxis selaginoides</i>	t	17			FR <sup>60</sup>
<b>THYMELAEACEAE</b>					
<i>Pimelea ferruginea</i>	w		3		
<i>Pimelea hispida</i>	w				FR <sup>40,59</sup>
<i>Pimelea humilis</i>	n,s,t,v				

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>					
<i>Pimelea imbricata</i> var. <i>piligera</i>	w		3		
<i>Pimelea ligustrina</i> s.l.	n,q,s,t,v	25			
<i>Pimelea linifolia</i> s.l.	n,q,s,t,v	25,29			S <sup>43</sup> ,FR <sup>22</sup>
<i>Pimelea pagophila</i> V	v		44		HS <sup>44</sup>
<i>Pimelea rosea</i>	w				MS <sup>40</sup>
<i>Pimelea suaveolens</i>	w	14,24			S <sup>24</sup> ,SP <sup>21</sup>
<b>TREMANDRACEAE</b>					
<i>Tetratheca ciliata</i>	n,s,t,v	25,29,34			S <sup>43</sup>
<i>Tetratheca gunnii</i> CE	t	34			MS <sup>60</sup>
<i>Tetratheca hirsuta</i>	w	15		21	HS <sup>21</sup>
<i>Tetratheca labillardierei</i>	n,t,v	17			
<i>Tetratheca pilosa</i> s.l.	n,s,t,v	17,18,25			S <sup>18,43</sup>
<i>Tetratheca procumbens</i>	t	17,34			
<i>Tetratheca setigera</i>	w				HS <sup>40,59</sup>
<i>Tetratheca subaphylla</i>	n,v	13			HS <sup>22</sup>
<i>Tremandra stelligera</i>	w				S <sup>46</sup>
<b>VIOLACEAE</b>					
<i>Hybanthus floribundus</i>	w			21	Q <sup>21</sup>
<b>WINTERACEAE</b>					
<i>Bubbia semecarpoides</i>	q				FR <sup>36</sup>
<i>Tasmannia lanceolata</i>	n,t,v	8,17,34			
<i>Tasmannia purpurascens</i> V	n	35			HS <sup>35</sup>
<b>XANTHOPHYLLACEAE</b>					
<i>Xanthophyllum octandrum</i>	q				S <sup>36</sup>
<b>XANTHORRHOEACAE</b>					
<i>Lomandra caespitosa</i>	w			21	
<i>Lomandra confertifolia</i> s.l.	n,q,v				FR <sup>42</sup>
<i>Lomandra filiformis</i>	n,q,v	29			
<i>Lomandra hermaphrodita</i>	w				FR <sup>21</sup>
<i>Lomandra integra</i>	w			21	
<i>Lomandra longifolia</i>	n,q,s,v	29			FR <sup>60</sup>
<i>Lomandra nigricans</i>	w				FR <sup>59</sup>
<i>Lomandra obliqua</i>	n,q				FR <sup>42</sup>
<i>Lomandra odora</i>	w	14			S <sup>46</sup>
<i>Lomandra pauciflora</i>	w				FR <sup>40,59</sup>
<i>Lomandra preissii</i>	w				FR <sup>40</sup>
<i>Lomandra sonderi</i>	w	14,28,46,		21	S <sup>46,48</sup> MS <sup>21</sup>

FAMILY	Distribution	Isolation			Susceptibility Rating
		In wild	In cultivation	By experiment	
<i>Species</i>		48			HS <sup>14</sup>
<i>Lomandra sparteia</i>	w				FR <sup>21</sup>
<i>Xanthorrhoea arenaria</i> V	t			9	HS <sup>9,60</sup>
<i>Xanthorrhoea australis</i>	n,s,t,v	2,4,13,17 22,29,34	44		S <sup>43</sup> ,HS <sup>22,61</sup>
<i>Xanthorrhoea bracteata</i>	t	34			
<i>Xanthorrhoea brevistyla</i>	w				HS <sup>40,41</sup>
<i>Xanthorrhoea drummondii</i>	w				S <sup>46</sup>
<i>Xanthorrhoea glauca</i> subsp. <i>glauca</i>	n,q	13			HS <sup>13</sup>
<i>Xanthorrhoea gracilis</i>	w	14,24,28, 37,46,48, 53			SP <sup>21</sup> S <sup>24,37,46,48,53</sup> HS <sup>14,40</sup>
<i>Xanthorrhoea latifolia</i>	w	38			
<i>Xanthorrhoea nana</i>	w				S <sup>46</sup>
<i>Xanthorrhoea platyphylla</i>	w	58			HS <sup>40,58,59</sup>
<i>Xanthorrhoea preissii</i>	w	2,14,22, 24,28,37, 46,48,53			SP <sup>21</sup> ,S <sup>37,46,48,53</sup> HS <sup>14,40,59</sup>
<i>Xanthorrhoea quadrangulata</i>	s	?			
<i>Xanthorrhoea resinifera</i>	n,q,v	13			
<i>Xanthorrhoea semiplana</i> var. <i>semiplana</i>	s	18			
<i>Xanthorrhoea semiplana</i> var. <i>tateana</i>	s	18			
<b>XYRIDACEAE</b>					
<i>Xyris exilis</i> V	w				FR <sup>58,59</sup>
<b>ZAMIACEAE</b>					
<i>Macrozamia communis</i>	n	2			S <sup>22</sup>
<i>Macrozamia riedlei</i>	w	2,14,15, 24			SP <sup>21</sup> S <sup>46,53</sup> HS <sup>40</sup>

## A4 REFERENCES

- 1 Fraser LR (1956) *Phytophthora cinnamomi* attacking native plants. *Australian Plant Disease Recorder* **8**, 12.
- 2 Pratt BH, Heather WA (1973) The origin and distribution of *Phytophthora cinnamomi* Rands in Australian native plant communities and the significance of its association with particular plant species. *Australian Journal of Biological Science* **26**, 559-573.
- 3 Taylor PA (1974) Ecological studies on the occurrence of *Phytophthora cinnamomi* on Black Mountain, ACT PhD Thesis, Australian National University, Canberra.
- 4 Weste G (1974) *Phytophthora cinnamomi* – the cause of severe disease in certain native communities in Victoria. *Australian Journal of Botany* **22**, 1-8.
- 5 Weste G (1981) Changes in the vegetation of sclerophyll shrubby woodland associated with invasion by *Phytophthora cinnamomi*. *Australian Journal of Botany* **29**, 261-276.
- 6 Gerrettson-Cornell L (1986) *Phytophthora cinnamomi* in New South Wales. Forestry Commission of NSW. Government Printer, Sydney.
- 7 Kennedy J, Weste G (1986) Vegetation change associated with invasion by *Phytophthora cinnamomi* on monitored sites in the Grampians, western Victoria. *Australian Journal of Botany* **34**, 251-279.
- 8 Podger FD, Brown MJ (1989) Vegetation damage caused by *Phytophthora cinnamomi* on disturbed sites in temperate rainforest in western Tasmania. *Australian Journal of Botany* **37**, 443-480.
- 9 Barker PCJ, Wardlaw TJ (1995) Susceptibility of selected rare plants to *Phytophthora cinnamomi*. *Australian Journal of Botany* **43**, 379-386.
- 10 Weste G (1997) The changing status of disease caused by *Phytophthora cinnamomi* in Victorian open forests, woodlands and heathlands. *Australasian Plant Pathology* **26**, 1-9.
- 11 Bullock S, Gunn LV, Summerell BA (2000) Pathogens of the Wollemi Pine, *Wollemia nobilis*. *Australasian Plant Pathology* **29**, 211-214.
- 12 Brett Summerell (Royal Botanic Gardens Sydney) pers. comm. (Brett.Summerell@rbgsyd.nsw.gov.au)
- 13 McDougall KL, Summerell BA (2003) The impact of *Phytophthora cinnamomi* on the flora and vegetation of New South Wales – a re-appraisal. In 'Phytophthora in Forests and Natural Ecosystems'. 2<sup>nd</sup> International IUFRO Working Party 7.02.09 Meeting, Albany, Western Australia, October 2001. Eds. JA McComb, GESTJ Hardy and IC Tommerup; pages 49-56. (Murdoch University Print: Murdoch, Western Australia).
- 14 Shearer BL, Dillon M (1995) Susceptibility of plant species in *Eucalyptus marginata* forest to infection by *Phytophthora cinnamomi*. *Australian Journal of Botany* **43**, 113-134. (16)
- 15 Podger FD (1968) Aetiology of jarrah dieback – a disease of dry sclerophyll *Eucalyptus marginata* Sm. forests in Western Australia. M.Sc. Thesis, University of Melbourne.
- 16 Brown BN (1998) Occurrence and impact of *Phytophthora cinnamomi* and other *Phytophthora* species in rainforests of the Wet Tropics World Heritage Area, and of the Mackay region, Qld. In 'Patch Deaths in Tropical Queensland Rainforests:

- Association and Impact of *Phytophthora cinnamomi* and Other Soil Borne Pathogens'. (Ed. PA Gadek) pp41-76. (Cooperative Research Centre for Tropical Rainforest Ecology and Management, Cairns).
- 17 Podger FD, Palzer C, Wardlaw TJ (1990) A guide to the distribution of *Phytophthora cinnamomi* and its effects on native vegetation. *Tasforests* **2**, 13-20.
  - 18 Lee TC, Wicks TS (1977) *Phytophthora cinnamomi* in native vegetation in South Australia. *Australasian Plant Pathology Society Newsletter* **6**, 22-23.
  - 19 Vickery FJ (1997) The distribution of *Phytophthora cinnamomi* on Kangaroo Island. PhD thesis, University of New England, Armidale.
  - 20 South Australian Research and Development Institute (SARDI), *pers. comm.*, pirsa.sardi@saugov.sa.gov.au
  - 21 McDougall KL (1997) Vegetation patterns in the northern jarrah forest of Western Australia in relation to dieback history and the current distribution of *Phytophthora cinnamomi*. PhD Thesis, Murdoch University, Western Australia.
  - 22 Keith McDougall (NSW Department of Environment and Conservation) *pers. comm.*, keith.mcdougall@environment.nsw.gov.au
  - 23 McCredie TA, Dixon KW, Sivasithamparam K (1985) Variability in the resistance of *Banksia* L. f. species to *Phytophthora cinnamomi* Rands. *Australian Journal of Botany* **33**, 629-637. (only plants dying after inoculation are listed)
  - 24 Shearer BL, Dillon M (1996) Susceptibility of plant species in *Banksia* woodlands on the Swan Coastal Plain, Western Australia, to infection by *Phytophthora cinnamomi*. *Australian Journal of Botany* **44**, 433-445. (15)
  - 25 David Cahill (Deakin University) *pers. comm.*, cahill@deakin.edu.au
  - 26 McDougall KL, Hardy GESTJ, Hobbs RJ (2001) Additions to the host range of *Phytophthora cinnamomi* in the jarrah (*Eucalyptus marginata*) forest of Western Australia. *Australian Journal of Botany* **49**, 193-198. (12)
  - 27 University of Pretoria (2002) The Forestry and Agricultural Biotechnology Institute (Tree Pathology Co-operative Program, *Phytophthora* root and collar rot of cold tolerant eucalypts pamphlet, 2002).
  - 28 Gardner JH, Rokich PA (1987) *Phytophthora cinnamomi* in operational and rehabilitated bauxite mine areas in south-western Australia. Environmental Research Bulletin Number 13. Alcoa of Australia Limited: Booragoon.
  - 29 Weste G, Brown K, Kennedy J, Walshe T (2002) *Phytophthora cinnamomi* infestation – a 24-year study of vegetation change in forests and woodlands of the Grampians, western Victoria. *Australian Journal of Botany* **50**, 247-274.
  - 30 Burgman MA and Ferguson IS (1995) Rainforest in Victoria: a review of the scientific basis of current and proposed protection measures. Forests Service Technical Reports 95-4. Department of Conservation and Natural Resources, Victoria.
  - 31 Podger FD, Batini F (1971) Susceptibility to *Phytophthora cinnamomi* root-rot of thirty six species of *Eucalyptus*. *Australian Forest Research* **5**, 9-20.
  - 32 Peters D, Weste G (1997) The impact of *Phytophthora cinnamomi* on six rare native tree and shrub species in the Brisbane Ranges, Victoria. *Australian Journal of Botany* **45**, 975-995.
  - 33 Podger FD (1989) Comparative pathogenicity of fourteen Australian isolates of *Phytophthora cinnamomi* determined on transplants of Tasmanian temperate heathland. *Australian Journal of Botany* **37**, 491-500.
  - 34 Schahinger R, Rudman T, Wardlaw T (2003) Conservation of Tasmanian plant species and communities threatened by *Phytophthora cinnamomi*. Strategic

- Regional Plan for Tasmania. Nature Conservation Branch Technical Report 03/03. Tasmanian Department of Primary Industries, Water and Environment.
- 35** McDougall KL, Summerell BA (2003) *Phytophthora cinnamomi* causing disease in subalpine vegetation in New South Wales. *Australasian Plant Pathology* **32**, 1-3.
  - 36** Worboys S, Gadek PA, Abell S, Jensen R (2003) Impact of dieback on rainforest canopies. In Rainforest Dieback Mapping and Assessment. (Ed. PA Gadek, S Worboys) pp23-68. (Cooperative Research Centre for Tropical Rainforest Ecology and Management, Cairns).
  - 37** Hill (1990) Dieback Diseases and other *Phytophthora* spp. in the Northern Kwongan. In *Nature, Conservation, Landscape and Recreational Values of the Lesueur Area. A report to the Environmental Protection Authority from the Department of Conservation and Land Management.* eds. A Burbidge, SD Hopper & S van Leeuwen, EPA, WA.
  - 38** Elaine Davison (Curtin University) *pers. comm.*, e.davison@curtin.edu.au
  - 39** WA Department of Conservation and Land Management. (undated) List of Priority One Flora. [http://www.calm.wa.gov.au/plants\\_animals/pdf\\_files/cfr\\_pri1.pdf](http://www.calm.wa.gov.au/plants_animals/pdf_files/cfr_pri1.pdf)
  - 40** Wills RT (1993) The ecological impact of *Phytophthora cinnamomi* in the Stirling Range National Park, Western Australia. *Australian Journal of Ecology* **18**, 145-159. (unpublished data from this paper).
  - 41** Wills RT, Keighery GJ (1994) Ecological impact of plant disease on plant communities. *Journal of the Royal Society of Western Australia* **77**, 127-131.
  - 42** Howard CG, McDougall KL, Summerell BA and Burgess LW (2004) Spatial relationship between *Phytophthora cinnamomi* presence and coastal vegetation health in south eastern Australia. Poster presentation to IUFRO conference.
  - 43** Victorian National Parks Service (1995) 2.7.2 *Phytophthora cinnamomi* control in parks, National Parks and Conservation Reserves Guidelines and Procedures Manual, Volume 1, Victorian Department of Conservation and Natural Resources, Melbourne
  - 44** Reiter N, Weste, G, Guest D (2004) The risk of extinction resulting from disease caused by *Phytophthora cinnamomi* to endangered, vulnerable or rare species endemic to the Grampians, western Victoria. *Australian Journal of Botany* **52**, 425-433.
  - 45** Stuart Worboys, *pers. comm.*, sworboys@ghd.com.au
  - 46** Alcoa of Australia (2002) Indicators of *P. cinnamomi* used by Interpreters. Environment Research Bulletin, Alcoa World Alumina, Booragoon, Western Australia.
  - 47** Cho JJ (1983) Variability in susceptibility of some *Banksia* Species to *Phytophthora cinnamomi* and their distribution in Australia. *Plant Disease*, **67**, 869-871.
  - 48** Edmiston RJ (1989) Plants Resistant to Dieback. Department of Conservation and Land Management. 1-89.
  - 49** Environment Australia (2001) Threat Abatement Plan for Dieback caused by the root-rot fungus *Phytophthora cinnamomi*.
  - 50** Gardner JH, Rokich PA (1987) *Phytophthora cinnamomi* in operational and rehabilitated bauxite mine area in South-western Australia. Alcoa of Australia Environment Research Bulletin. 13.
  - 51** Hart R (1983) Report on Dieback due to *Phytophthora cinnamomi* in Two People's Bay Nature Reserve. Department of Fisheries and Wildlife.

- 52** Keighery G (1992) The impact of *Phytophthora* species on rare plants. In *Dieback-What is the Future?*, Eds. MJ Freeman, R Hart & M Ryall, Northern Sandplains Dieback Working Group, Perth. pp29-36.
- 53** Podger FD (1972) *Phytophthora cinnamomi*, a cause of lethal disease in indigenous plant communities in Western Australia. *Phytopathology*, **62**, 972-981.
- 54** Titze JF and Palzer CR (1969) Host list of *Phytophthora cinnamomi* Rands with special reference to Western Australia. Forestry and Timber Bureau, Department of National Development. Technical Note No. 1.
- 55** Tynan KM, Scott ES and Sedgley M (1998) Evaluation of *Banksia* species for response to *Phytophthora* infection. *Plant Pathology*, **47**, 446-455.
- 56** Tynan KM, Wilkinson CJ, Holmes JM, Dell B, Colquhoun IJ, McComb JA, Hardy GESTJ (2001) The long-term ability of phosphite to control *Phytophthora cinnamomi* in two native plant communities of Western Australia. *Australian Journal of Botany*, **49**, 761- 770.
- 57** Renate Velzeboer (South Australian Department of Environment and Heritage), *pers. comm.* (Velzeboer.Renate@sa.gov.au)
- 58** Sarah Barrett and Malcolm Grant (Department of Conservation and Land Management, WA), *pers. comm.* (sarahba@calm.wa.gov.au, malcomg@calm.wa.gov.au)
- 59** Barrett S (1996) Biological survey of mountains of south western Australia. Project No. AW03. Australian Nature Conservation Agency.
- 60** Tim Rudman (Department of Primary Industries, Water and Environment, Tasmania) *pers. comm.* - personal observations (Tim.Rudman@dpiwe.tas.gov.au)
- 61** Tim Rudman (Department of Primary Industries, Water and Environment, Tasmania) unpublished data from disease monitoring plots (Tim.Rudman@dpiwe.tas.gov.au)