Importation of cooked duck meat from Thailand

Draft review

July 2019



© Commonwealth of Australia 2019

Ownership of intellectual property rights

Unless otherwise noted, copyright (and any other intellectual property rights, if any) in this publication is owned by the Commonwealth of Australia (referred to as the Commonwealth).

Creative Commons licence

All material in this publication is licensed under a Creative <u>Commons Attribution 4.0 International Licence</u>, except content supplied by third parties, logos and the Commonwealth Coat of Arms.

Inquiries about the licence and any use of this document should be emailed to copyright@agriculture.gov.au.



Cataloguing data

This publication (and any material sourced from it) should be attributed as: Australian Department of Agriculture 2019, *Importation of cooked duck meat from Thailand*, Department of Agriculture, Canberra. CC BY 4.0.

This publication is available at agriculture.gov.au/biosecurity/risk-analysis

Australian Government Department of Agriculture

GPO Box 858 Canberra ACT 2601

Switchboard: +61 2 6272 3933 or 1800 900 090

Web agriculture.gov.au

Email animal@agriculture.gov.au

The Australian Government acting through the Department of Agriculture has exercised due care and skill in preparing and compiling the information and data in this publication. Notwithstanding, the Department of Agriculture, its employees and advisers disclaim all liability, including liability for negligence and for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying on any of the information or data in this publication to the maximum extent permitted by law.

Stakeholder submissions on draft reports

This draft report has been issued to give all interested parties an opportunity to comment on relevant technical biosecurity issues, with supporting rationale. A final report will then be produced taking into consideration any comments received.

Submissions should be sent to the Australian Government Department of Agriculture following the conditions specified within the related Biosecurity Advice, which is available at: <u>agriculture.gov.au/biosecurity/risk-analysis/memos</u>.

Contents

Glo	ssary	v
Sum	nmary	
1	Introd	uction2
	1.1	Australia's biosecurity framework
	1.2	Background2
	1.3	Scope
	1.4	Existing conditions
	1.5	Duck meat industry 4
	1.6	Next Steps 4
2	Metho	d6
	2.1	Review of hazard identification
	2.2	Risk assessment
	2.3	Risk management
	2.4	Risk communication 13
3	Hazard	l identification15
4	Risk re	views23
	4.1	Duck hepatitis A virus
	4.2	Duck Tembusu virus
	4.3	Duck virus enteritis (duck plague) 41
	4.4	Waterfowl parvoviruses 49
5	Requir	ements for the importation of cooked duck meat from Thailand
	5.1	Eligibility
	5.2	Documentation
	5.3	Certification
	5.4	Annex 1
6	Review	v of processes65
	6.1	Review of conditions
	0.1	

Figures

Figure 1. Components of the unrestricted risk estimate	7
Figure 2. Hazard identification and refinement	16
Figure 3. Inactivation and environmental stability of Zika virus	34

Tables

Table 1. Matrix for combining qualitative likelihoods	9
Table 2. Rules for determining the overall effect of establishment and/or spread	. 11
Table 3. Likely consequences: a combination of the likelihood and overall effect of establishment and/or spread	. 12
Table 4. Risk estimation matrix	12
Table 5. Hazard identification	. 17

Glossary

Term or abbreviation	Definition
Appropriate level of protection (ALOP)	The level of protection deemed appropriate by the Member establishing a sanitary or phytosanitary measure to protect human, animal or plant life or health within its territory
AS	Attenuated strain
Bean geese	Anser fabalis
Biosecurity	The prevention of the entry, establishment or spread of unwanted pests and infectious disease agents to protect human, animal or plant health or life, and the environment
Cherry valley duck	A breed of domestic duck
Chicken	Gallus gallus domesticus
CM IRA/chicken meat IRA 2008	Generic Import Risk Analysis Report for Chicken Meat 2008
Code	OIE Terrestrial Animal Health Code 2017
CRISPR	Clustered regularly interspaced short palindromic repeats
DEDS	Duck Egg Drop Syndrome
The department	The Australian Government Department of Agriculture
DHAV-1	Duck hepatitis A virus type 1
DNA	Deoxyribonucleic acid
Domestic duck	Anas platyrhynchos domesticus
DPV	Duck parvovirus
DTMUV	Duck Tembusu virus
DVE	Duck virus enteritis
DVEV	Duck virus enteritis virus
ELISA	Enzyme-linked immunosorbent assay
Endemic	Belonging to, native to, or prevalent in a particular geography, area or environment
FSANZ	Food Standards Australia New Zealand
FMDV	Foot and mouth disease virus
GPV	Goose parvovirus
Host	An organism that harbours a parasite, mutual partner, or commensal partner, typically providing nourishment and shelter
ICTV	International Committee on Taxonomy of Viruses
Import permit	Official document authorising importation of a commodity in accordance with specified sanitary import requirements
Mallard	Anas platyrhynchos
MDPV	Muscovy duck parvovirus

Term or abbreviation	Definition
Mule duck	Cairina moschata and Anas platyrhynchos domesticus cross breed
Muscovy duck	Cairina moschata
OIE	World Organisation for Animal Health
Pathogen	A biological agent that can cause disease to its host
Pathway	Any means that allows the entry or spread of a pest
PCR	Polymerase chain reaction
Pekin duck	A breed of domestic duck
Pest	Any species, strain or biotype of plant, animal, or pathogenic agent injurious to plants or animals
PFU	Plaque forming unit
Quarantine	Official confinement of regulated articles for observation and research or for further inspection, testing or treatment
Quarantine pest	A pest of potential economic importance to the area, not yet present there, or present but not widely distributed and being officially controlled
Restricted risk	Risk estimate with sanitary measure(s) applied
RNA	Ribonucleic acid
RT-PCR	Reverse transcription polymerase chain reaction
Spp	Species
Stakeholders	Government agencies, individuals, community or industry groups or organizations, whether in Australia or overseas, including the proponent/applicant for a specific proposal, who have an interest in the issues and import conditions
Surveillance	An official process which collects and records data on pest occurrence or absence by surveying, monitoring or other procedures
Tsiay duck	A breed of domestic duck
TMUV	Tembusu virus
Unrestricted risk	Risk estimate in the absence of sanitary measure(s)
VS	Virulent strain
White-fronted geese	Anser albifrons

Summary

The department has prepared this draft review of biosecurity requirements in response to a request by Thailand for market access to Australia for cooked duck meat.

This draft review takes into account new and relevant peer-reviewed scientific information and advice from international scientific experts.

Australia currently only permits the importation of canned or retorted duck meat products that meet specific temperature and time requirements during the manufacturing process.

This draft review proposes that the importation of cooked duck meat to Australia from Thailand be permitted, subject to biosecurity measures.

This draft review identifies hazards that could be present in cooked duck meat from Thailand and therefore proposes the following to achieve Australia's appropriate level of protection (ALOP):

- The duck meat must have been sourced from domestic ducks (*Anas platyrhynchos domesticus* and *Cairina moschata*) that were hatched, raised and slaughtered in Thailand.
- The meat must be limited to muscle meat and overlying skin and fat.
- The ducks from which the meat was derived must have passed ante- and post-mortem veterinary inspection under official veterinary supervision.
- The ducks must have been slaughtered in an abattoir that meets standards at least equivalent to those contained in the Australia New Zealand Food Standards Code Standard 4.2.2 Primary Production and Processing Standard for Poultry Meat (Australia Only) (FSANZ 2010).
- The ducks must have been slaughtered and processed in an abattoir and processing facility approved by the Veterinary Authority of Thailand.
- The duck meat must have been cooked in a commercial process using moist heat to a minimum core temperature of at least 70°C for a minimum of 8.2 minutes (or time/temperature equivalent approved by the department).

These measures reduce animal biosecurity risks to a level that is consistent with Australia's ALOP.

This draft review was developed as an extension of the *Generic Import Risk Analysis Report for Chicken Meat* (Biosecurity Australia 2008). Disease information relevant to duck meat was reviewed and updated and diseases that occur in ducks but not chickens were also reviewed. Disease agents that were clearly identified as susceptible to the cooking parameters described above, such as avian influenza viruses and Newcastle disease virus, were removed from further assessment at the hazard identification stage.

This draft review contains details of the risk assessments and the biosecurity measures for the identified hazards. Interested stakeholders are invited to provide comments and submissions to the department within the 60 day consultation period.

References

FSANZ 2010, 'Standard 4.2.2: primary production and processing standard for poultry meat (Australia only)', *Australia New Zealand Food Standards Code*, Food Standards Australia New Zealand, Canberra, available at <u>https://www.legislation.gov.au/Details/F2012L00292</u> accessed 28 June 2016.

Introduction

1.1 Australia's biosecurity framework

Australia's biosecurity framework aims to protect Australia from the entry, establishment and spread of exotic pests and diseases that would threaten Australia's agricultural industries and unique flora and fauna that are relatively free from serious pests and diseases.

The risk analysis process is an important part of developing and reviewing Australia's biosecurity system. It enables the Australian Government to formally assess the risks associated with proposals to import new products into Australia. If the risks are found to exceed Australia's appropriate level of protection (ALOP), risk management measures are recommended to reduce the risks to an acceptable level. If it is not possible to reduce the risks to an acceptable level, import of the goods into Australia will not be permitted until suitable risk management measures are identified.

Successive Australian Governments have maintained a conservative, but not zero risk approach to the management of biosecurity risks. This approach is expressed in terms of Australia's ALOP, which reflects community expectations through import conditions and is described as providing a high level of protection aimed at reducing risk to a very low level, but not to zero.

The department's science-based risk assessment process is consistent with Australian Government policy as well as Australia's rights and obligations under the World Trade Organization's Agreement on the Application of Sanitary and Phytosanitary Measures. This assessment may take the form of an import risk analysis, a non-regulated review of existing conditions or technical advice.

Further information about Australia's biosecurity framework is provided in the *Biosecurity Import Risk Analysis Guidelines 2016* located on the <u>Australian Government Department of Agriculture</u> website.

The department recognises that there might be new scientific information and technologies, or other measures that may provide an equivalent level of biosecurity protection from the disease agents identified as requiring risk management. Submissions supporting equivalence measures will be considered case-by-case.

1.2 Background

Currently, only foie gras from France, canned or retorted duck meat products and meat-based flavours that meet specific temperature and time requirements during the manufacturing process are permitted into Australia.

This draft review of the animal biosecurity risks associated with the importation of cooked duck meat from Thailand has been undertaken in response to a request from Thailand's National Bureau of Agricultural Commodity and Food Standards and the Department of Livestock Development for access to Australian markets for cooked duck meat.

Many diseases of ducks also occur in chickens, therefore, this draft review was developed as an extension of the *Generic Import Risk Analysis Report for Chicken Meat* (Biosecurity Australia 2008) (chicken meat IRA (2008)). The disease information that was relevant to duck meat was reviewed and

updated to include any advances in scientific knowledge that have occurred since the release of the chicken meat IRA in 2008. Specific diseases that occur in ducks but not chickens were also reviewed.

1.3 Scope

The scope of this draft review is limited to an assessment of the animal biosecurity risks posed by the importation of cooked duck meat from Thailand. The definition of cooking is taken from the chicken meat IRA (2008) for control of Newcastle disease virus, which requires heating the meat in a commercial process using moist heat to a minimum core temperature of 70°C for a minimum of 8.2 minutes (or time/temperature equivalent).

In this draft review the definition of duck meat is limited to muscle tissue from domestic ducks (*Anas platyrhynchos domesticus* and *Cairina moschata*, or mule ducks, which are a hybrid of *Anas platyrhynchos domesticus* and *Cairina moschata*). It includes soft tissues contained within the muscle body, such as nerves and blood contained within muscle vasculature. It also includes skin and fat overlying the muscle. This draft review does not include an assessment of the risks associated with duck offal, product containing bone or cartilage, or whole duck. In addition, the duck must have been slaughtered in an abattoir that meets standards at least equivalent to those contained in the *Australia New Zealand Food Standards Code – Standard 4.2.2 – Primary Production and Processing Standard for Poultry Meat (Australia Only)* (FSANZ 2010).

This draft review is limited to diseases covered in the chicken meat IRA (2008) that infect ducks, as well as diseases that were not considered in the chicken meat IRA (2008) because they infect ducks but not chickens. It also takes into account relevant changes in scientific knowledge since the release of the chicken meat IRA (2008). Disease agents in the chicken meat IRA (2008) that were clearly identified as susceptible to the cooking parameters described above were removed from further assessment at the hazard identification stage, as described in Chapter 3.

1.4 Existing conditions

International arrangements

Import conditions exist for foie gras from France, canned or retorted duck meat products and meatbased flavours from all countries, for human consumption.

The <u>import requirements</u> for these commodities can be found on the department's website (<u>agriculture.gov.au</u>).

Australia takes into account the following when considering the approval of, and conditions for, the export of animals and their products to Australia from any country:

- the animal health status of the country
- the capacity of veterinary services and other relevant certifying authorities
- legislative controls over animal health, including biosecurity
- the standard of reporting of major contagious disease outbreaks to the World Organisation for Animal Health (OIE)
- the veterinary laboratory services in the country, including compliance with relevant international standards

• systems in place to maintain the integrity of the certification/documentation of products intended for export to Australia.

Domestic arrangements

The Australian Government is responsible for regulating the movement of animals and animal products into and out of Australia. However, the state and territory governments are responsible for animal health and environmental controls within their individual jurisdictions. Legislation relating to resource management or animal health may be used by state and territory government agencies to control interstate movement of animals and their products. Once animals and animal products have been cleared by Australian biosecurity officers, they may be subject to interstate movement conditions. It is the importer's responsibility to identify and ensure compliance with all requirements.

Trade movements of all types of duck meat can occur freely between all states and territories in Australia. Restrictions have existed from time to time due to outbreaks of exotic disease such as virulent Newcastle disease or highly pathogenic avian influenza, during eradication of these diseases.

1.5 Duck meat industry

Australia's duck processing industry is highly concentrated with two companies located in New South Wales and Victoria accounting for the bulk of production. There are also several smaller duck farmers, largely located in Victoria and Western Australia, who process their own birds for local markets.

A small number of backyard ducks are kept in Australia. Most backyard poultry flocks are chickens that are maintained for table egg production and while ducks may be used for egg production they are not as common.

1.6 Next Steps

This draft review is released for 60 days public consultation to give stakeholders the opportunity to provide technical comment. Stakeholder submissions will be considered when finalising the draft review.

The final review will be published on the department's website along with a notice advising stakeholders of the release. The department will also notify the proposer, the registered stakeholders and the World Trade Organization Secretariat about the release of the final report. Publication of the final report represents the end of the review process. The conditions recommended in the final report will be the basis of any imports permitted.

Should the final review recommend that the importation of cooked duck meat to Australia from Thailand be permitted, subject to biosecurity measures, the department will verify the systems in place in Thailand prior to issuing any import permits. This will ensure there is official oversight of the processes and that the biosecurity integrity of the processing is maintained from when the duck is slaughtered to when the final product is packaged ready for export. A favourable report will be required before import permits can be issued and trade can commence.

References

Biosecurity Australia 2008, *Generic import risk analysis report for chicken meat: final report. Part C - detailed assessments*, Biosecurity Australia, Canberra, available at http://agriculture.gov.au/biosecurity/risk-analysis/animal/chicken-meat.

FSANZ 2010, 'Standard 4.2.2: primary production and processing standard for poultry meat (Australia only)', *Australia New Zealand Food Standards Code*, Food Standards Australia New Zealand, Canberra, available at <u>https://www.legislation.gov.au/Details/F2012L00292</u> accessed 28 June 2016.

2 Method

The World Organisation for Animal Health (OIE), in its *Terrestrial Animal Health Code* (OIE 2018), describes the components of risk analysis in Chapter 2.1 as:

- hazard identification
- risk assessment (entry assessment, exposure assessment, consequence assessment and risk estimation)
- risk management
- risk communication.

Hazard identification, risk assessment and risk management are sequential steps within a risk analysis. Risk communication is conducted as an ongoing process, and includes both formal and informal consultation with stakeholders.

This draft review has drawn on the following sources of information (this list is not exhaustive):

- the Generic import risk analysis report for chicken meat (Biosecurity Australia 2008)
- the Terrestrial Animal Health Code 2018 (OIE 2018)
- a review of relevant scientific literature.

The department considered that the comprehensive chicken meat IRA (2008) was still relevant and applicable, and provided a strong framework for a review of the animal biosecurity risks for the importation of cooked duck meat from Thailand. Assessments made in the chicken meat IRA (2008) which was completed as a semi quantitative (as a numerical estimate) assessment, were accepted in the draft review. Where an agent was not covered or new information that may affect the final outcome of an assessment was available, the department applied the qualitative (in words) assessment method described in this review.

The chicken meat IRA (2008) was completed to cover raw product from any country. The draft review for the importation of cooked duck meat from Thailand considered only cooked duck meat from Thailand.

2.1 Review of hazard identification

Hazard identification is described in the Code (Article 2.1.2) as a classification step that is undertaken to identify potential hazards that may be associated with the importation of a commodity (OIE 2018).

In accordance with the Code, a disease agent was considered to be a potential hazard relevant to the importation of cooked duck meat if it was assessed to be:

- a disease of ducks
- emerging or capable of producing adverse consequences.

A hazard was retained for further review (hazard refinement) if:

• it was not present in Australia, or present in Australia and a notifiable disease or subject to official control or eradication

• it was present in the country of export (Thailand).

However, as this draft review is for the hazards associated with cooked duck meat, disease agents that were identified in the chicken meat IRA (2008) as being susceptible to inactivation by cooking (definition in Section 1.3 of this document) were not retained for risk assessment.

2.2 Risk assessment

Details of the risk assessment process relevant to animals and animal products are provided in Chapter 2.1 of the Code (OIE 2018).

In accordance with the Code, the entry assessment describes the probability of the entry of each of the potential hazards in an importing country and the exposure assessment describes the biological pathways necessary for exposure of animals and estimates the probability of exposure occurring. The consequence assessment describes the potential consequences of a given exposure and estimates the probability of them occurring. The unrestricted risk estimate is the combination of the likelihood of entry, exposure, establishment and/or spread, and the overall effect of establishment and/or spread.

Steps in determining the unrestricted risk estimate are illustrated diagrammatically in Figure 1.

Figure 1. Components of the unrestricted risk estimate



A review of peer-reviewed scientific literature was conducted and contact with relevant experts sought, where necessary, for each hazard retained for risk assessment. Based on this information, a decision was made whether or not to continue with the risk assessment as outlined below.

The risk assessment concluded with an unrestricted risk for each hazard. If the unrestricted risk did not achieve Australia's ALOP, then risk management measures were recommended to reduce the risk to achieve the ALOP.

Evaluating and reporting likelihood

For those hazards retained for risk assessment, the assessment was conducted using a qualitative approach based on the same qualitative descriptive definitions described in the chicken meat IRA (2008).

Entry assessment

The entry assessment considered a single entry scenario defined as the period from slaughter, cooking and export up to arrival in Australia. A number of factors were taken into account in determining the likelihood of a disease agent entering Australia in cooked duck meat, such as:

• prevalence of the hazard in ducks in Thailand

- ante-mortem inspection
- post-mortem inspection
- tissue distribution of the disease agent
- cross-contamination at slaughter
- the effect of cooking, including inactivation temperature (and duration required) for the disease agent
- the effect of storage and transport.

For each disease agent, a qualitative likelihood was then assigned to describe the likelihood of the disease agent being present in the imported cooked duck meat products arriving at the Australian border. The final outcome of the entry assessment was the likelihood of entry of a potential hazard into Australia.

Exposure assessment

The exposure assessment describes the process that was used to estimate the likelihood that a susceptible bird in Australia will be exposed to the cooked duck meat contaminated with a disease agent. It takes into account the groups of birds most likely to be affected as well as the possible pathways by which exposure of these groups could occur.

Exposure groups

The term 'exposure group' categorises a group of animals that may be susceptible to one or more of the potential hazards/pathogens considered in risk assessments. The chicken meat IRA (2008) identified the four most likely exposure groups to imported poultry meat:

- 1) wild birds
- 2) low biosecurity poultry—backyard poultry and free-range commercial poultry including ratites
- 3) medium biosecurity poultry—non-genetic stock commercial poultry
- 4) non-avian species, where applicable.

The chicken meat IRA (2008) analysed the sequence of steps for imported infected chicken meat to cause infection in susceptible animals (the exposure pathways). The exposure pathways for imported duck meat are similar. As only cooked products would be imported, it is likely that there would be less on-shore processing than what was assumed in the chicken meat IRA (2008). Therefore, most imported product would move from retailer/distributor direct to household consumers or to the food service industry. Product not consumed would be either dumped (where it may or may not be exposed to wild birds or to non-avian species) or, more likely in product sold directly to households, it could be exposed to backyard (low biosecurity) poultry. There are restrictions on feeding poultry meat or the by-products of poultry processing to ruminants, however, this material may be fed to birds or poultry. If raw or semi-cooked product were allowed entry under an approved arrangement for further processing, disposal of all waste in a manner that manages the animal biosecurity risks would be required

Given there would probably be minimal on-shore processing, the likelihood of waste product from processing being made into rendered product, and then into poultry feed, would be lower than described in the chicken meat IRA (2008).

In addition to the distribution variables, summarised above and described in detail in the chicken meat IRA (2008), there are a number of exposure group dependent variables:

- the likelihood that birds/animals in each exposure group will ingest duck meat material should they be exposed to it
- the likelihood that feed containing rendered duck meat may be fed to birds/animals in each exposure group

and pathogen-dependent variables, including:

- the hardiness of the pathogen and the likelihood it will remain viable in the environment for the period before it is exposed to the susceptible animals
- the likelihood that an infective dose is consumed.

These variables are discussed in detail in the chicken meat IRA (2008).

For each agent, the final outcome of the exposure assessment was an estimate of the likelihood that susceptible birds in each exposure group would be exposed to the disease agent via the contaminated imported product (i.e. the likelihood of exposure).

Estimation of likelihood of entry and exposure

The likelihood of entry and exposure for each exposure group was estimated by combining the likelihood of entry and the corresponding likelihood of exposure using the matrix described in Table 1. The basis for combining qualitative likelihoods using a matrix is described by Standards Australia and Standards New Zealand (Standards Australia 2005).

	High	Negligible	Extremely low	Very low	Low	Moderate	High
Likelihood of entry	Moderate	Negligible	Extremely low	Very low	Low	Low	Moderate
	Low	Negligible	Extremely low	Very low	Very low	Low	Low
	Very low	Negligible	Extremely low	Extremely low	Very low	Very low	Very low
	Extremely low	Negligible	Negligible	Extremely low	Extremely low	Extremely low	Extremely low
	Negligible	Negligible	Negligible	Negligible	Negligible	Negligible	Negligible
		Negligible	Extremely low	Very Low	Low	Moderate	High

Table 1. Matrix for combining qualitative likelihoods

Likelihood of exposure

Likelihood of establishment and/or spread associated with the outbreak scenario

Once exposure of a susceptible population has occurred, a number of possible outbreak scenarios could follow. These represent a continuum ranging from no spread to widespread establishment. The chicken meat IRA (2008) grouped all likely outbreak scenarios into four categories:

- the disease agent does not establish or is not recognised within the directly exposed population
- the disease agent establishes within the directly exposed population only, is identified and is eradicated

- the disease agent establishes within the directly exposed population, spreads to other populations, including other exposure groups if applicable, but is eradicated
- the disease agent establishes within the directly exposed population and spreads to other populations.

In this draft review, for each exposure group, all categories of outbreak scenarios were evaluated for plausibility based on the epidemiology of each disease agent. The most likely outbreak scenario for each hazard, resulting from the exposure of susceptible animals, was considered (described in the relevant disease chapter).

The likelihood of the outbreak scenario occurring was then estimated to obtain a likelihood of establishment and/or spread.

When estimating the effects associated with the outbreak scenario, qualitative descriptors were used as described in the chicken meat IRA (2008).

Determination of the overall effects of establishment and/or spread associated with the outbreak scenario

The overall effects of establishment and/or spread were addressed in terms of direct and indirect effects on the life and health of susceptible animals on a national scale, including adverse human health, environmental and socioeconomic effects. Impacts on human life and health are the responsibility of the Australian Government Department of Health and Food Standards Australia New Zealand (FSANZ). The department consults with these agencies on assessments for zoonotic agents.

Direct effects:

- life or health (including production effects) of susceptible animals, including public health consequences
- the living environment, including life and health of wildlife, and any effects on the non-living environment.

Indirect effects:

- new or modified eradication, control, monitoring or surveillance and compensation strategies or programs
- domestic trade or industry, including changes in consumer demand and effects on other industries supplying inputs to, or using outputs from, directly affected industries
- international trade, including loss of markets, meeting new technical requirements to enter or maintain markets and changes in international consumer demand
- the environment, including biodiversity, endangered species and the integrity of ecosystems
- communities, including reduced tourism, reduced rural and regional economic viability and loss of social amenity, and any 'side effects' of control measures.

An effect was not assessed more than once and direct effects were considered separately from indirect effects.

The overall effect of establishment and/or spread associated with the outbreak scenario took into account the geographic level of these effects:

- local—restricted to a single locality or town
- regional—a recognised geographic area such as far north Queensland
- state or territory
- national

and the magnitude of these effects:

- indiscernible—not usually distinguishable from normal day-to-day variation
- minor significance—recognisable, but minor and reversible
- significant—serious and substantive, but reversible and unlikely to have permanent economic effects
- highly significant—extremely serious and irreversible and likely to have permanent economic effects.

An outbreak may occur on a small geographical level but have significant national effects, and vice versa. Based on the geographic level and magnitude of effects, the overall effect of establishment and/or spread was determined using the rules described in Table 2.

Extreme	The effect is likely to be highly significant at the national level. Implies that economic stability, societal values or social well-being would be seriously affected.
High	The effect is likely to be significant at the national level and highly significant within affected zones. Implies that the effect would be of national concern. However, serious effects on economic stability, societal values or social well-being would be limited to a given zone.
Moderate	The effect is likely to be recognised on a national level and significant within affected zones. The effect is likely to be highly significant to directly affected parties.
Low	The effect is likely to be recognised within affected zones and significant to directly affected parties. It is not likely that the effect will be recognised at the national level.
Very low	The effect is likely to be minor to directly affected parties. The effect is unlikely to be recognised at any other level.
Negligible	The effect is unlikely to be recognised at any level within Australia.

Table 2. Rules for determining the overall effect of establishment and/or spread

Consequence assessment

The likely consequences were determined by combining the likelihood of establishment and/or spread (associated with the outbreak scenario) with the overall effect of establishment and/or spread using the matrix shown in Table 3.

Table 3. Likely consequences: a combination of the likelihood and overall effect of establishment and/or spread

lent	nent	High	Negligible	Very low	Low	Moderate	High	Extreme
istablishn spread	Moderate	Negligible	Very low	Low	Moderate	High	Extreme	
	Low	Negligible	Negligible	Very low	Low	Moderate	High	
d of e	Likelihood of e and/or	Very low	Negligible	Negligible	Negligible	Very low	Low	Moderate
lihoo		Extremely low	Negligible	Negligible	Negligible	Negligible	Very low	Low
Like		Negligible	Negligible	Negligible	Negligible	Negligible	Negligible	Very low
			Negligible	Very low	Low	Moderate	High	Extreme

Overall effect of establishment and spread

Risk estimation and evaluation

Risk estimation is the integration of likelihood of entry and exposure, and likely consequences of a hazard introduced by the importation of cooked duck meat from Thailand into Australia.

The risk is estimated by:

- determining the likelihood of entry and exposure
- determining the likelihood of establishment and/or spread among susceptible populations and the overall effect of establishment and/or spread to estimate the likely consequences
- combining the likelihood of entry and exposure with the estimate of likely consequences.

Combining the likelihood of entry and exposure and likely consequences was undertaken using the rules shown in the risk estimation matrix in Table 4.

	,						
Likelihood of entry and exposure	High likelihood	Negligible risk	Very low risk	Low risk	Moderate risk	High risk	Extreme risk
	Moderate likelihood	Negligible risk	Very low risk	Low risk	Moderate risk	High risk	Extreme risk
	Low likelihood	Negligible risk	Negligible risk	Very low risk	Low risk	Moderate risk	High risk
	Very low likelihood	Negligible risk	Negligible risk	Negligible risk	Very low risk	Low risk	Moderate risk
	Extremely low likelihood	Negligible risk	Negligible risk	Negligible risk	Negligible risk	Very low risk	Low risk
	Negligible likelihood	Negligible risk	Negligible risk	Negligible risk	Negligible risk	Negligible risk	Very low risk
		Negligible effect	Very low effect	Low effect	Moderate effect	High effect	Extreme effect

Table 4. Risk estimation matrix

Risk evaluation is described in the OIE Code as the process of comparing the estimated risk with a country's ALOP.

A risk estimation that was either 'very low' or 'negligible' was considered sufficient to achieve Australia's ALOP. This provided a benchmark for evaluating risk and determining whether risk management was required.

The use of a benchmark for evaluating risks for each disease agent is illustrated in the process outlined below:

- if the unrestricted risk was 'negligible' or 'very low', then it achieved Australia's ALOP and risk management was not required
- if the unrestricted risk was 'low', 'moderate', 'high' or 'extreme', risk management measures were required.

This was considered the final output of the risk assessment.

2.3 Risk management

Once the unrestricted risk for a particular hazard has been assessed and evaluated as exceeding Australia's ALOP, measures to manage and reduce that risk are considered and proposed. The imposition of a particular risk management measure or a combination of measures results in the derivation of the restricted risk. The aim of risk management measures being to meet Australia's ALOP by reducing the restricted risk to 'very low' or 'negligible'.

Risk management options may be imposed pre-border with the purpose of reducing the likelihood of hazards entering Australia, or post-arrival aiming to prevent the exposure and/or establishment and spread of the hazard in susceptible local populations.

2.4 Risk communication

Risk communication is defined in the Code as 'the interactive transmission and exchange of information and opinions throughout the risk analysis process concerning risk, risk-related factors and risk perceptions among risk assessors, risk managers, risk communicators, the general public and other interested parties' (OIE 2018).

In conducting import risk analyses and reviews, the department consults with the Australian Government Department of Health to ensure that public health considerations are included in the development of Australia's animal biosecurity measures. Furthermore, a formal process of consultation with external stakeholders is a standard procedure for all import risk analyses and reviews to enable stakeholder assessment and feedback on draft conclusions and recommendations about Australia's animal biosecurity measures.

References

Biosecurity Australia 2008, *Generic import risk analysis report for chicken meat: final report. Part C* - *detailed assessments*, Biosecurity Australia, Canberra, available at <u>http://agriculture.gov.au/biosecurity/risk-analysis/animal/chicken-meat</u>.

OIE 2018, 'Terrestrial Animal Health Code', World Organisation for Animal Health, France, available at <u>http://www.oie.int/standard-setting/terrestrial-code/access-online/</u> accessed 31 October 2018.

Standards Australia 2005, *Risk management guidelines: companion to AS/NZS 4360:2004*, Standards Australia International and Standards New Zealand, Sydney and Wellington.

3 Hazard identification

The list of diseases (hazards) of potential animal biosecurity concern was compiled from:

- diseases identified in the chicken meat IRA (2008), Conditions for the importation from approved countries of fertile eggs (domestic duck) (1999) and Importation of hatching (fertile) duck eggs from approved countries – policy review (2009) (Biosecurity Australia 1999, 2009)
- other diseases identified in the literature as occurring in ducks.

The method of hazard identification and refinement is described in Section 2.1. The hazard identification decision making process is shown in Figure 2. The preliminary list of diseases/disease agents is shown in Table 5. This table summarises the results of the hazard refinement process, including the reason for removal or retention of each identified hazard.

Ubiquitous or common commensals which may be present in Australia in addition to those that are opportunistic, not reported to be pathogenic, or of uncertain relevance in the commodity due to limited or insufficient information were included in the hazard refinement process.

The initial hazard list was taken from the chicken meat IRA (2008). It was updated to include agents that are specific to ducks and considered relevant to this draft review.

Hazards that were assessed in the chicken meat IRA (2008) as having an unrestricted risk below Australia's ALOP were not further assessed. Similarly, hazards that were assessed in the chicken meat IRA (2008) as being susceptible to the cooking parameters as described in the scope (Section 1.3), were not further assessed in this draft review.

However, an exception to the above was when the department determined there was evidence of significantly different epidemiology or adverse effects of an agent between chickens and ducks. These hazards were reassessed as primary pathogens of ducks.

The following diseases were retained for risk assessment on the basis of the information provided in Table 5:

- 1) Duck hepatitis A virus
- 2) Duck Tembusu virus
- 3) Duck virus enteritis
- 4) Waterfowl parvovirus.





Table 5. Hazard identification

Disease (disease agent)	Disease affects or is spread by domestic ducks (Anas platyrhynchos domesticus and Cairina moschata)	Emerging and/or capable of producing adverse consequences	Present in Australia	Present in Thailand	Assessed in CM IRA as either not requiring risk assessment or not requiring risk management	Agent assessed in CM IRA and found to be inactivated by cooking as defined in scope	Retained for risk review
Avian infectious laryngotrachitis virus	No	Yes	Yes	Yes	Yes	No	No
Avian nephritis virus	Yes	Yes	Yes	No	Yes	No	No
Avian paramyxovirus 2 and avian paramyxovirus 3	No	Yes	No	Yes	No	Yes	No
Borrelia anserina	Yes	Yes	Yes	No	Yes	No	No
Brachyspira spp	Yes	Yes	Yes	Yes	No	No	No
Campylobacter jejuni	Yes	Yes	Yes	Yes	Yes	No	No
Chlamydophila psittaci	Yes	Yes	Yes	Yes	Yes	No	No
Duck adenovirus-1 (Egg Drop Syndrome Virus/Atadenovirus)	Yes	Yes	Yes	No	No	No	No
Duck adenovirus-2 (Adenovirus Group 1)	Yes	Yes	Yes	No	No	Νο	No
Duck hepatitis A virus (Duck virus hepatitis type 1/ picornaviridae)	Yes	Yes	No	Yes	No	No	Yes
Duck Tembusu virus	Yes	Yes	No	Yes	No	No	Yes
Duck virus hepatitis type 2 (Duck astrovirus 1/ DAstV-1)	Yes	Yes	No	Noª	No	Νο	No ^b
Duck virus hepatitis type 3 (Duck astrovirus -2/DAstV- 2)	Yes	Yes	No	Noc	No	No	No ^b

Disease (disease agent)	Disease affects or is spread by domestic ducks (Anas platyrhynchos domesticus and Cairina moschata)	Emerging and/or capable of producing adverse consequences	Present in Australia	Present in Thailand	Assessed in CM IRA as either not requiring risk assessment or not requiring risk management	Agent assessed in CM IRA and found to be inactivated by cooking as defined in scope	Retained for risk review
Duck virus enteritis	Yes	Yes	No	Yes	No	No	Yes
Eastern equine encephalitis/Western equine encephalitis viruses	Yes	Yes	No	EEE Yes WEE No	Yes	No	No No
Enterohaemorrhagic Escherichia coli	Yes	No	Yes	Yes	Yes	No	No
Fowl pox	Yes	Yes	Yes	Yes	Yes	No	No
Haemophilus paragallinarum/ Avibacterium paragallinarum	No	Yes	Yes	Yes	Yes	No	No
Highly pathogenic avian influenza virus	Yes	Yes	No (Controls)	No	Νο	Yes	No
Infectious bursal disease virus serotype 1 (very virulent and antigenic variant strains)	No ^d	Yes	No	Yes	No	No	No
Japanese Encephalitis virus	Yes	Yes	Yes	Yes	Yes	No	No
Low pathogenicity avian influenza virus	Yes	Yes	No (Controls)	No	Νο	Yes	No
Mareks disease	No	Yes	Yes	Yes	Yes	No	No
Muscovy duck reovirus	Yes	Yes	No	No	No	No	No
Mycobacterium avium	Yes	Yes	Yes (Controls)	Yes	Yes	No	No
Mycoplasma gallisepticum	Yes	Yes	Yes	Yes	Yes	No	No
Mycoplasma synoviae	Yes	Yes	Yes	Yes	Yes	No	No
Mycoplasma anatis	Yes	Yes	No	No	No	No	No

Disease (disease agent)	Disease affects or is spread by domestic ducks (Anas platyrhynchos domesticus and Cairina moschata)	Emerging and/or capable of producing adverse consequences	Present in Australia	Present in Thailand	Assessed in CM IRA as either not requiring risk assessment or not requiring risk management	Agent assessed in CM IRA and found to be inactivated by cooking as defined in scope	Retained for risk review
Newcastle disease	Yes	Yes	No (Controls)	Yes	No	Yes	No
Ornithobacterium rhinotracheale	Yes	Yes	Yes	Yes	Yes	No	No
Pasteurella multocida	Yes	Yes	Yes	Yes	Yes	No	No
Riemerella anatipestifer	Yes	Yes	Yes	Yes	Yes	No	No
Salmonella enteritidis	Yes	Yes	Yes (controls)	Yes	No	Yes	No
Salmonella Gallinarum	Yes	Yes	No ^e (controls)	No	No	Yes	No
Salmonella Pullorum	Yes	Yes	No ^e (controls)	Yes	No	Yes	No
Salmonella Typhimurium, antibiotic resistant strains	Yes	Yes	No	Yes	No	Yes	No
Waterfowl parvoviruses	Yes	Yes	No	Yes	No	No	Yes
West Nile virus	Yes	Yes	No	Yes	Yes	No	No

a Only ducks appear to be affected by duck virus hepatitis type 2 (duck astrovirus type 1 [DAstV-1]). Outbreaks only appear to have occurred in England and potentially China. The earliest described outbreaks occurred in ducks kept on open fields in eastern England, initially reported in 1965 and up to 1969, then again in 1983/84, which were the last reported outbreaks in England (Gough et al. 1985; OIE 2018; Woolcock & Tsai 2013). In 2008 there was a severe outbreak of duck hepatitis in China. Virus present in liver extracts had very high amino acid sequence identity (Fu et al. 2008) to the DAstV-1 sequenced by Todd et al. (2009)and very different to DAstV-2 (Todd et al. 2009).

b Due to the similarities between the three types of duck virus hepatitis, any risk management that would be required for Duck hepatitis A virus (Duck virus hepatitis type 1/ picornaviridae) would also apply to duck virus hepatitis type 2 (duck astrovirus type 1 [DAstV-1]) and duck virus hepatitis type 3 (duck astrovirus type 2 [DAstV-2])

c Duck virus hepatitis type 3 (duck astrovirus type 2 [DAstV-2]) has only been described in the United States of America (OIE 2018; Woolcock 2008).

d Available literature suggests that IBDV serotype 1 is capable of infecting ducks and producing a limited antibody response, probably due to poor virus replication. No clinical or microscopic signs have been observed in ducks to date. Virus has been isolated from faeces and caecal tonsils of wild ducks in one study. Experimental infection studies in ducks have failed to isolate virus. There is no available literature to support the conclusion that significant replication of pathogenic IBDV serotype 1 occurs in ducks (Eddy 1990; Eterradossi & Saif 2013; Hollmén et al. 2000; McFerran et al. 1980; McNulty, Allan & McFerran 1979; Okoye, Iguomu & Nwosuh 1990; Oladele 2010; Oluwayelu et al. 2007; Tsai et al. 1983; Woo-Jin et al. 2008; Yamada, Matsuo & Uchinuno 1982).

e Salmonella Gallinarum and Salmonella Pullorum are not present in commercial poultry.

References

Biosecurity Australia 1999, *Conditions for the importation from approved countries of fertile eggs (domestic duck)*, Department of Agriculture, Fisheries and Forestry, Canberra, available at <u>http://agriculture.gov.au/biosecurity/risk-analysis/animal/hatchingeggs-ducks</u> accessed 9 January 2019.

---- 2009, Importation of hatching (fertile) duck eggs from approved countries – policy review, Canberra, available at http://agriculture.gov.au/biosecurity/risk-analysis/animal/fertile-duck-eggs.

Eddy, RK 1990, 'Antibody responses to infectious bursal disease virus serotypes 1 and 2 in ducks', Veterinary Record, vol. 127, no. 15, p. 382 (Abstract only)

Eterradossi, N & Saif, YM 2013, 'Infectious bursal disease', in Swayne, DE, Glisson, JR, McDougald, LR, Nolan, LK, Suarez, DL & Nair, VL (eds), *Diseases of poultry*, 13th edn, Wiley-Blackwell, Ames, Iowa, USA, pp. 219-46.

Fu, Y, Pan, M, Wang, X, Xu, Y, Yang, H & Zhang, D 2008, 'Molecular detection and typing of duck hepatitis A virus directly from clinical specimens', *Veterinary Microbiology*, vol. 131, no. 3, pp. 247-57, available at <u>https://doi.org/10.1016/j.vetmic.2008.03.011</u>, accessed 8 March 2016.

Gough, RE, Borland, ED, Keymer, IF & Stuart, JC 1985, 'An outbreak of duck hepatitis type II in commercial ducks', *Avian Pathology*, vol. 14, no. 2, pp. 227-36, available at https://doi.org/10.1080/03079458508436224, accessed 9 January 2019.

Hollmén, T, Franson, JC, Docherty, DE, Kilpi, M, Hario, M, L.H., C & Petersen, MR 2000, 'Infectious bursal disease virus antibodies in eider ducks and herring gulls', *The Condor*, vol. 102, no. 3, pp. 688-91, available at http://dx.doi.org/10.1650/0010-5422(2000)102[0688:IBDVAI]2.0.CO;2, accessed 9 January 2019.

Langevin, SA, Bunning, M, Davis, B & Komar, N 2001, 'Experimental infection of chickens as candidate sentinels for West Nile virus', *Emerging Infectious Diseases*, vol. 7, no. 4, pp. 726-9

McFerran, JB, McNulty, MS, McKillop, ER, Connor, TJ, McCracken, RM, Collins, DS & Allan, GM 1980, 'Isolation and serological studies with infectious bursal disease viruses from fowl, turkeys and ducks: demonstration of a second serotype', *Avian Pathology*, vol. 9, pp. 395-404

McNulty, MS, Allan, GM & McFerran, JB 1979, 'Isolation of infectious bursal disease virus from Turkeys', *Avian Pathology*, vol. 8, no. 3, pp. 205-12, available at http://www.informaworld.com/10.1080/03079457908418346

OIE 2018, 'Duck virus hepatitis', in *Manual of diagnostic tests and vaccines for terrestrial animals 2018*, World Organisation for Animal Health, Paris, accessed 9 January 2019.

Okoye, JOA, Iguomu, EP & Nwosuh, C 1990, 'Pathogenicity of an isolate of infectious bursal disease virus in local Nigerian ducks', *Tropical Animal Health and Production*, vol. 22, no. 3, pp. 160-2, available at <u>https://doi.org/10.1007/BF02241008</u>, accessed 9 January 2019.

Oladele, OA 2010, 'Chronological studies on experimental infectious bursal disease virus infections in chickens, turkeys and ducks', *Journal of Commonwealth Veterinary Association*, vol. 26, no. 1, pp. 23-8, available at http://www.commonwealthvetassoc.org/pdf/Jan2010.pdf#page=25

Oluwayelu, DO, Emikpe, BO, Oladele, OA, Ohore, OG & Fagbohum, OA 2007, 'Seroprevalence of infectious bursal disease in flocks of indigenous Nigerian ducks (Anas platyrhynchos)', *Journal of Animal and Veterinary Advances*, vol. 6, no. 1, pp. 64-7, available at https://www.medwelljournals.com/abstract/?doi=javaa.2007.64.67, accessed 9 January 2019.

Todd, D, Smyth, VJ, Ball, NW, Donnelly, BM, Wylie, M, Knowles, NJ & Adair, BM 2009, 'Identification of chicken enterovirus-like viruses, duck hepatitis virus type 2 and duck hepatitis virus type 3 as astroviruses', *Avian Pathology*, vol. 38, no. 1, pp. 21-9, available at https://doi.org/10.1080/03079450802632056, accessed 9 January 2019.

Tsai, HJ, Lu, YS, Lin, DF & Lee, SH 1996, 'Pathogenicity of duck and chicken isolates of infectious bursal disease virus', *Taiwan Journal of Veterinary Medicine* and Animal Husbandry, vol. 66, no. 1, pp. 19-26

Wilcox, GE, Flower, RLP, Baxendale, W & Mackenzie, JS 1983, 'Serological survey of wild birds in Australia for the prevalence of antibodies to egg drop syndrome 1976 (EDS-76) and infectious Bursal disease viruses', *Avian Pathology*, vol. 12, no. 1, pp. 135-9, available at http://dx.doi.org/10.1080/03079458308436155

Woo-Jin, J, Eun-Kyoung, L, Seong-Joon, J, Jun-hun, K, Chang-Bum, Y, Yeo-Sung, Y & Kang-Seuk, C 2008, 'Very virulent infectious bursal disease virus isolated from wild birds in Korea: epidemiological implications', *Virus Research*, vol. 137, no. 1, pp. 153-6, available at https://doi.org/10.1016/j.virusres.2008.06.013, accessed 9 January 2019.

Woolcock, PR 2008, 'Duck hepatitis', in Saif, YM (ed) *Diseases of Poultry*, 12th edn, Blackwell Publishing, Ames, Iowa, USA.

Woolcock, PR & Tsai, H-J 2013, 'Duck hepatitis', in Swayne, DE, Glisson, JR, McDougald, LR, Nolan, LK, Suarez, DL & Nair, VL (eds), *Diseases of poultry*, 13th edn, Wiley-Blackwell, Ames, Iowa, USA, pp. 422-31.

Yamada, S, Matsuo, K & Uchinuno, Y 1982, 'Susceptibility of ducks and duck-origin cell cultures to infectious bursal disease virus', *Avian Diseases*, vol. 26, no. 3, pp. 596-601

4 Risk reviews

4.1 Duck hepatitis A virus

4.1.1 Background

Duck hepatitis A virus (DHAV) includes three antigenically unrelated genotypes, 1, 2 and 3. Duck hepatitis A virus type 1 (DHAV-1) is the most prevalent and widespread of the three types. There is limited information available on types 2 and 3, however, the clinical presentation is reported to be similar to type 1. Therefore, this draft risk review will focus on DHAV-1.

DHAV-1 is a highly contagious, acute and lethal pathogen that affects young ducklings. Normally, after three weeks of age, ducklings begin to develop age-related resistance, and at six weeks of age are no longer susceptible. However, DHAV-1 was determined to be the cause of egg drop, reduced feed consumption and ovary-oviduct infection in laying ducks in China (Zhang et al. 2018). DHAV-1 is one of five viruses that causes liver disease in ducks, however DHAV-1 is considered to be the most pathogenic and causes significant losses in the duck industry (Lin et al. 2016; Song et al. 2014). This type can cause high mortality and morbidity levels with a mortality rate greater than 80 per cent in ducklings under three weeks of age (Lin et al. 2016).

The first recorded identification of duck hepatitis was in 1945, on Long Island, the United States of America (Fabricant, Rickard & Levine 1957). It has now been officially reported in many countries in Europe, Asia and North America. Outbreak investigations have been reported for cases in Canada, Egypt, Japan, Korea, Poland, Taiwan, the United States of America and Vietnam (Doan et al. 2016; Erfan et al. 2015; Friend & Trainer 1974; Hassaan, Shahin & Eid 2018; Kamomae et al. 2017; Kozdru, Czekaj & Lorek 2014; MacPherson & Avery 1957; Soliman et al. 2015; Tseng, Knowles & Tsai 2007). DHAV-2 has been identified in Taiwan and DHAV-3 in China, Vietnam and South Korea (Doan et al. 2016; Li et al. 2013).

Duck virus hepatitis is an OIE-listed disease and nationally notifiable in Australia (Department of Agriculture 2017; OIE 2018).

4.1.2 Technical information

Epidemiology

Susceptible species

Ducklings are the primary hosts, and mature ducks are not affected. Field observations indicate chickens and turkey are resistant (Woolcock & Tsai 2013). A serological survey of bean and white fronted geese in Germany showed none had antibodies to DHAV (Hlinak et al. 1998).

In an unpublished study investigating species sensitivity to the virus, Cherry Valley and Tsiay ducklings were found to be highly sensitive species, Pekin ducklings and hybrid Pekin and Tsiay ducklings were relatively sensitive to disease while mule ducklings showed intermediate sensitivity and Muscovy ducklings were resistant (Woolcock & Tsai 2013).

Modes of transmission

In the field, DHAV-1 can spread quickly through a flock, infecting all susceptible ducklings. Although this virus is considered extremely contagious, there have been exceptions within farms with adjoining duck pens reporting both negligible mortality and 65 per cent mortality (Woolcock & Fabricant 1991).

The natural routes of transmission are airborne or via contaminated feed or water. The infection route is the pharynx or upper respiratory tract (Toth & Norcross 1981). DHAV-1 was found in one study to be lethal when ducklings were infected with an aerosol form of the virus (Priz, cited inPriz, cited inWoolcock & Tsai 2013). Oral vaccination of ducklings has resulted in successful immunisation (Tripathy & Hanson 1986).

Incubation period

The incubation period is generally one to two days with all mortalities occurring within three to four days. Generally ducklings die within an hour of clinical signs appearing (Woolcock & Tsai 2013).

Ducklings that have recovered may excrete the virus for up to eight weeks in faeces, and have immunity with the serum containing virus neutralising antibodies (Kamomae et al. 2017; Reuss, cited Reuss, cited Woolcock & Tsai 2013).

Distribution and prevalence

There have been no official reports of DHAV-1 in Thailand; however, a duck hepatitis virus has been isolated from 9 to 14 day old ducklings from Chonburi and Rayong provinces (Sirivan, Tantaswasdi & Chuenchai 1997). Experimental vaccination of ducks with a DHV vaccine resulted in protection but the unvaccinated ducklings were susceptible to infection with 60 to 80 per cent mortality (Sirivan, Tantaswasdi & Chuenchai 1997).

There is variation amongst strains, and reports of the pathogen in China have indicated that the diversity of strains is dependent upon where they are isolated (Li et al. 2013). The virus strains can be allocated into three regional groups dependent upon the phylogenetic analysis of the VP1 gene. Interestingly, Asian DHAV-1 clusters with Egyptian DHAV-1, with the closest nucleotide homology to the Asian strains being 98 per cent. The difference in virulence between the strains was identified when embryonated eggs were inoculated with the Egyptian strain of DHAV-1. These eggs did not display lesions, while reports of inoculations with the Asian strain of DHAV-1 indicate that lesions are usually observed (Erfan et al. 2015).

Australian Status

DHAV-1 has not been reported in Australia and there is no active surveillance. However, duck virus hepatitis is nationally notifiable.

Agent properties

DHAV-1 is a single-stranded, positive-sense RNA virus and is classified as the type strain of the genus *Avihepatovirus* in the family *Picornaviridae* (Tseng, Knowles & Tsai 2007; Yang et al. 2008). The pathogen is one of three genotypes and was originally identified in 1950 in chicken embryos after which it was classified as an enterovirus. It was not until after 2006 when the complete genome was determined that the virus was classified in the new genus *Avihepatovirus* (ICTV 2018).

DHAV-1 can remain viable in the environment for an extended period of time, particularly in cold climates. In laboratory conditions the virus has been able to survive for nine years at -20°C and for longer than two years at 4°C (Hanson, Rhoades & Schricker 1964; Dvorakova, Asplin, cited in Dvorakova, Asplin, cited in Woolcock 2003). In warmer conditions, the virus has been able to survive at 37°C for seven days (Hanson, Rhoades & Schricker 1964; Pollard, cited inPollard, cited inWoolcock & Tsai 2013).

In more natural conditions, the virus has been recorded to survive greater than 37 days in moist faeces in a cool shed (Asplin, cited in Asplin, cited in Woolcock & Tsai 2013). Furthermore, it has been found to survive in infected feeders for at least 10 weeks and withstand heating to 50 to 56°C for 60 minutes or more (Trefilov et al. 2018). However, another study showed that DHAV was inactivated rapidly at 56°C with 99 per cent of activity lost in 30 minutes and none detectable after 90 minutes, as measured by chicken embryo ID_{50} (Hanson, Rhoades & Schricker 1964). Further studies on the virus showed the half-life of DHAV at 56°C to be 1.26 minutes but this could be extended by the addition of salts to the virus solution (Davis 1987).

Several other members of the family *Picornaviridae*, namely foot and mouth disease virus (FMDV), hepatitis A virus and poliovirus, are heat resistant to effective inactivation up to 60°C (Kamolsiripichaiporn et al. 2007). This is supported by a study by Gubbins and colleagues (2016) in the inactivation of FMDV by heat treatments used in extruded pet food manufacture. They observed that there was no detectable virus following treatment of FMDV-spiked meat slurry at 68°C for 5 minutes; treatment of FMDV-spiked slurry and meal mix at 79°C for 10 or 30 seconds, or treatment of homogenised bovine tongue epithelium, taken from an FMDV-infected animal, at 79°C for 10 seconds.

Exposure to 3 per cent chloramine, 2 per cent caustic soda or 1 per cent formaldehyde at 15°C to 20°C for two hours will also inactivate DHAV due to pH sensitivity (Dvorakova, Pollard, cited in Dvorakova, Pollard, cited in Woolcock 2003). The infectivity of DHAV was lost at pH less than or equal to 2 and pH greater than or equal to 12 within 30 minutes of exposure (Davis 1987).

4.1.3 Presence in duck meat

Song et al (2014) researched the invasive properties of a virulent strain (VS) and an attenuated strain (AS) in different organs in ducklings and the levels of the two strains in the different organs were compared. RNA extracts from the heart, spleen, lung, kidney, brain, intestine and muscle of infected ducklings at 36 hours post infection were subjected to quantitative reverse transcriptase polymerase chain reaction (RT-PCR). The test showed high levels of VS and AS RNA in the liver, spleen and intestine, moderate levels in the kidney, muscle, lung and heart, and low levels in the brain. VS and AS showed similar tissue tropism in the liver, spleen and intestine, suggesting that the three organs are targeted by these two DHAV-1 strains, whilst the intestine might be the initial infection and proliferation site during natural infection. Moreover, although there was a higher number of AS virus copies inoculated than VS, significantly more VS virus RNA copies than AS were detected in each organ. These results demonstrated that VS replicated more efficiently than AS in ducklings.

The findings by Lin et al (2016) were similar to those of Song et al (2014). The viral loads of DHAV-1 in liver and spleen were significantly higher than those in pancreas, heart, kidney, brain, bursa of Fabricius and the thymus, which indicated that the liver and spleen of ducklings are the major target organs for the two viruses.

4.1.4 Pathogenesis

Infection in ducklings occurs mainly through the mucous membranes of the digestive and respiratory organs. The virus spreads via the blood circulation to several organs, especially the liver and brain. The liver becomes enlarged with petechial and ecchymotic hemorrhages throughout and there is extensive hepatocyte necrosis, hyperplasia and haemorrhages throughout (Trefilov et al. 2018).

Within the first hours of infection, virus titre in the blood circulation rises rapidly, then gradually decreases by 48 to 72 hours. The virus titre in the liver and brain also increases, usually within 48 to 72 hours of infection (Trefilov et al. 2018). Liver function decreases, allowing toxic products to spread throughout the body. Ducklings die as a result of irreversible changes in the liver and other organs. The toxins produced typically cause opisthotonus, especially in fat ducklings (Trefilov et al. 2018).

DHAV-1 has been isolated predominately in the allantoic liquids of duck embryos and the liver (Chen et al. 2013). The spleen, kidneys and bursa of Fabricius are also reported to show signs of disease (Jin et al. 2008; Sheng et al. 2014).

Lin et al (2016) determined that the age of the ducklings did not correlate with DHAV-1 viral loads in tested organs. Ducklings tested between 3 and 19 days old died regardless of the specific age when they reached a critical level of DHAV-1 (Lin et al. 2016).

Although wild aquatic fowl are considered to be mechanical carriers of the virus, studies have not confirmed this hypothesis. Testing 520 birds of six species found no serological confirmation of DHAV-1 infection (Asplin, cited in Asplin, cited in Woolcock & Tsai 2013). Furthermore, none of the 36 wild ducks of four species demonstrated evidence of DHAV-1 even though the virus had been found in domestic ducks from the same pond (Ulbrich, cited in Ulbrich, cited in Woolcock & Tsai 2013). However, wild duck embryonating eggs, when taken from an infected area, have been found to be susceptible to infection, confirming that although the species is susceptible, they do not appear to be extensive carriers of the virus (Woolcock & Tsai 2013).

4.1.5 Diagnosis

Clinical signs

DHAV-1 causes acute and highly contagious hepatitis that usually results in mortality within three to four days of infection. Soon after infection ducklings become lethargic and stop moving, have partially closed eyes and squat before falling on their sides. They then kick spasmodically, draw their heads back, and usually die within an hour of clinical signs starting (Fabricant, Rickard & Levine 1957; Kim et al. 2007).

Ducklings less than one week of age are affected most severely with mortality reaching up to 95 per cent. One to three week old ducklings usually suffer up to 50 per cent mortality, while ducklings four to five weeks of age have low or negligible mortality (Kim et al. 2007).

Pathology

The liver contains ecchymotic haemorrhages, is enlarged, and has the primary lesions. The kidney and spleen are often swollen, with the former having congested renal blood vessels and the latter showing signs of necrosis (Woolcock & Tsai 2013). As the disease progresses, the organs will appear differently to those in the acute stage of infection. For example, the liver will no longer be enlarged, but have irregular, pale mottling (Fabricant, Rickard & Levine 1957; Woolcock 2003).

Histological changes seen from DHAV-1 are necrosis of hepatic cells, subcutaneous oedema and bile duct hyperplasia with different degrees of haemorrhaging and inflammatory cell response. Ducklings that survive have been observed to have regeneration of the liver parenchyma (Fitzgerald, Hanson & Simon 1969; Woolcock 2003).

Testing

DHAV-1 can be isolated from the liver or allantoic liquids of duck embryos (Chen et al. 2013). The pathogen may then be confirmed by inoculation of the isolate into susceptible ducklings, into allantoic sacs of susceptible embryonating duck eggs or primary cultures of duck embryo liver cells (Woolcock 1986; Woolcock & Tsai 2013). Other detection techniques include RT-PCR, multiplex polymerase chain reaction and VP1 ELISA (Enzyme-linked immunosorbent assay) (Chen et al. 2013; Liu et al. 2010; Wen et al. 2014).

4.1.6 Treatment

There is no effective treatment for DHAV-1 infection.

4.1.7 Control

Both inactivated vaccines and live attenuated vaccines are available to protect ducks against DHAV-1.

Live attenuated DHAV-1 vaccines are used before the start of egg laying to immunise breeder ducks or to actively immunise susceptible ducklings. Breeder ducks, primed with live attenuated vaccine or previously exposed to live DHAV-1, may also be vaccinated using an inactivated DHAV-1 vaccine. Well-vaccinated breeder ducks may transmit passive immunity in eggs for eight to nine months (Woolcock & Tsai 2013).

4.1.8 Current biosecurity measures

For imports of fertile duck eggs into Australia, the flock from which the eggs were derived are required to be sample tested free from DHAV-1. The eggs are hatched under quarantine in Australia and the hatched ducklings undergo further testing.

4.1.9 Conclusion

- DHAV-1 is an OIE-listed disease and is nationally notifiable in Australia.
- DHAV-1 can cause high mortality and morbidity levels with a mortality rate greater than 80 per cent in ducklings under three weeks of age. Slaughter age ducks (7-8 weeks) are resistant to infection.
- Moderate levels of DHAV-1 have been demonstrated in the skeletal muscle of infected ducklings three days post infection.
- Ducklings that have recovered may excrete the virus for up to 8 weeks in faeces, and have immunity with the serum containing virus neutralising antibodies.
- There has been no official report of DHAV-1 in Thailand; however, a duck hepatitis virus (DHV) was isolated from 9 to14 day old ducklings from Chonburi and Rayong provinces.
- Several viruses of the family *Picornaviridae* demonstrate resistance to inactivation up to around 60°C. DHAV-1 can withstand heating to 50 to 56°C for 60 minutes but is likely to be completely

inactivated when heated to 70°C for 8.2 minutes, based on inactivation studies for other viruses of the same family.

Based on the above information cooking as defined in the scope at section 1.3 of this document (70°C for 8.2 minutes) will inactivate any DHAV-1 that may be present in meat of ducks slaughtered at seven to eight weeks of age. Therefore the department concluded that further risk assessment of DHAV-1 was not required.

References

Chen, L-L, Xu, Q, Zhang, R-H, Yang, L, Li, J-X, Xie, Z-J, Zhu, Y-L, Jiang, S-J & Si, X-K 2013, 'Improved duplex RT-PCR assay for differential diagnosis of mixed infection of duck hepatitis A virus type 1 and type 3 in ducklings', *Journal of Virological Methods*, vol. 192, no. 1-2, pp. 12-7, available at https://doi.org/10.1016/j.jviromet.2013.04.012, accessed 3 March 2016.

Davis, D 1987, 'Temperature and pH stability of duck hepatitis virus', *Avian Pathology*, vol. 16, no. 1, pp. 21-30, available at <u>http://dx.doi.org/10.1080/03079458708436349</u>, accessed 21 March 2016.

Department of Agriculture 2017, 'National list of notifiable animal diseases', Canberra, available at <u>http://www.agriculture.gov.au/pests-diseases-weeds/animal/notifiable</u> accessed 5 June 2019.

Doan, HTT, Le, XTK, Do, RT, Hoang, CTM, Nguyen, KT & Le, TH 2016, 'Molecular genotyping of duck hepatitis A viruses (DHAV) in Vietnam', *Journal of Infection in Developing Countries*, vol. 10, no. 9, pp. 988-95, available at <u>https://doi.org/10.3855/jidc.7239</u>, accessed 5 September 2018.

Erfan, AM, Selim, AA, Moursi, MK, Nasef, SA & Abdelwhab, EM 2015, 'Epidemiology and molecular characterisation of duck hepatitis A virus from different duck breeds in Egypt', *Veterinary Microbiology*, vol. 177, no. 3, pp. 347-52, available at <u>https://doi.org/10.1016/j.vetmic.2015.03.020</u>, accessed 21 March 2016.

Fabricant, J, Rickard, CG & Levine, PP 1957, 'The pathology of duck virus hepatitis', *Avian Diseases*, vol. 1, no. 3, pp. 256-75, available at <u>http://dx.doi.org/10.2307/1587740</u>, accessed 17 March 2016.

Fitzgerald, JE, Hanson, LE & Simon, J 1969, 'Histopathologic changes induced with duck hepatitis virus in the developing chicken embryo', *Avian Diseases*, vol. 13, no. 1, pp. 147-57, available at <u>http://dx.doi.org/10.2307/1588423</u>, accessed 7 March 2016.

Friend, M & Trainer, DO 1974, 'Experimental DDT-duck hepatitis virus interaction studies', *The Journal of Wildlife Management*, vol. 38, no. 4, pp. 887-95, available at http://dx.doi.org/10.2307/3800062, accessed 8 March 2016.

Gubbins, S, Forster, J, Clive, S, Schley, D, Zuber, S, Schaaff, J & Corley, D 2016, 'Thermal inactivation of foot and mouth disease virus in extruded pet food', *Revue Scientifique et Technique de l'Office International des Epizooties*, vol. 35, no. 3, pp. 965-72, available at https://doi.org/10.20506/rst.35.3.2582, accessed 3 January 2018.

Hanson, LE, Rhoades, HE & Schricker, RL 1964, 'Properties of duck hepatitis virus', *Avian Diseases*, vol. 8, no. 2, pp. 196-202, available at <u>http://dx.doi.org/10.2307/1588060</u>, accessed 3 March 2016.

Hassaan, MN, Shahin, AM & Eid, AAM 2018, 'Isolation and molecular identification of duck hepatitis A virus in Sharkia Governorate', *Zagazig Veterinary Journal*, vol. 46, no. 1, pp. 88-95, available at https://doi.org/10.5281/zenodo.1246279, accessed 5 September 2018.

Hlinak, A, Müller, T, Kramer, M, Mühle, RU, Liebherr, H & Ziedler, K 1998, 'Serological survey of viral pathogens in bean and white-fronted geese from Germany', *Journal of Wildlife Diseases*, vol. 34, no. 3, pp. 479-86, available at <u>https://doi.org/10.7589/0090-3558-34.3.479</u>, accessed 1 May 2018.

ICTV 2018, 'Genus: Avihepatovirus', International Committee on Taxonomy of Viruses, Washington, available at https://talk.ictvonline.org/ictv-reports/ictv_online_report/positive-sense-rna-viruses/picornavirales/w/picornaviridae/677/genus-avihepatovirus accessed 9 January 2019.

Jin, X, Zhang, W, Zhang, W, Gu, C, Cheng, G & Hu, X 2008, 'Identification and molecular analysis of the highly pathogenic duck hepatitis virus type 1 in Hubei province of China', *Research in Veterinary Science*, vol. 85, no. 3, pp. 595-8, available at <u>https://doi.org/10.1016/j.rvsc.2008.01.001</u>, accessed 21 March 2016.

Kamolsiripichaiporn, S, Subharat, S, Udon, R, Thongtha, P & Nuanualsuwan, S 2007, 'Thermal inactivation of foot-and-mouth disease viruses in suspension', *Applied and Environmental Microbiology*, vol. 73, no. 22, pp. 7177-84, available at http://aem.asm.org/cgi/content/abstract/73/22/7177

Kamomae, M, Kameyama, M, Ishii, J, Nabe, M, Ogura, Y, Iseki, H, Yamamoto, Y & Mase, M 2017, 'An outbreak of duck hepatitis A virus type 1 infection in Japan', *Journal of Veterinary Medical Science*, vol. 79, no. 5, pp. 917-20, available at <u>https://doi.org/10.1292/jvms.16-0646</u>, accessed 4 September 2018.

Kim, MC, Kwon, YK, Joh, SJ, Kwon, JH, Kim, JH & Kim, SJ 2007, 'Development of one-step reverse transcriptase-polymerase chain reaction to detect duck hepatitis virus type 1', *Avian Diseases*, vol. 51, no. 2, pp. 540-5, available at <u>http://avdi.allenpress.com/avdionline/?request=get-</u>document&doi=10.1637%2F0005-2086(2007)51%5B540:DOORTC%5D2.0.CO%3B2

Kozdru, W, Czekaj, H & Lorek, M 2014, 'Outbreak of duck viral hepatitis in duckling flocks in Poland - case report', *Bulletin of the Veterinary Institute in Puawy*, vol. 58, no. 4, pp. 513-5, available at <u>https://doi.org/10.2478/bvip-2014-0078</u>, accessed 5 September 2018.

Li, J, Bi, Y, Chen, C, Yang, L, Ding, C & Liu, W 2013, 'Genetic characterization of duck hepatitis A Viruses isolated in China', *Virus Research*, vol. 178, no. 2, pp. 211-6, available at https://doi.org/10.1016/j.virusres.2013.10.007, accessed 21 March 2016.

Lin, S-L, Cong, R-C, Zhang, R-H, Chen, J-H, Xia, L-L, Xie, Z-J, Wang, Y, Zhu, Y-L & Jiang, S-J 2016, 'Circulation and in vivo distribution of duck hepatitis A virus types 1 and 3 in infected ducklings', *Archives of Virology*, vol. 161, no. 2, pp. 405-16, available at <u>https://doi.org/10.1007/s00705-015-</u> <u>2648-z</u>, accessed 3 March 2016.

Liu, M, Zhang, T, Zhang, Y, Meng, F, Li, X, Hou, Z, Feng, X & Kong, X 2010, 'Development and evaluation of a VP1-ELISA for detection of antibodies to duck hepatitis type 1 virus', *Journal of Virological Methods*, vol. 169, no. 1, pp. 66-9, available at https://doi.org/10.1016/j.jviromet.2010.06.018, accessed 17 March 2016.

MacPherson, LW & Avery, RJ 1957, 'Duck virus hepatitis in Canada', *Canadian Journal of Comparative Medicine and Veterinary Science*, vol. 21, no. 2, pp. 26-31, available at http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1614372/, accessed 5 September 2018.

OIE 2018, 'OIE-listed diseases, infections and infestations in force in 2018', World Organisation for Animal Health, Paris, available at <u>http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-</u>2018/ accessed 31 October 2018.

Sheng, XD, Zhang, WP, Zhang, QR, Gu, CQ, Hu, XY & Cheng, GF 2014, 'Apoptosis induction in duck tissues during duck hepatitis A virus type 1 infection', *Poultry Science*, vol. 93, no. 3, pp. 527-34, available at <u>https://doi.org/10.3382/ps.2013-03510</u>, accessed 21 March 2016.

Sirivan, P, Tantaswasdi, U & Chuenchai, S 1997, 'Isolation and identification of duck hepatitis virus in ducklings', *Abstracts from The Journal of Thai Veterinary Medical Association: 1997 issues*, available at <u>http://tvma.tripod.com/news/abstracts97.html</u> accessed 7 November 2018.

Soliman, M, Alfajaro, MM, Lee, MH, Jeong, YJ, Kim, DS, Son, KY, Kwon, J, Choi, JS, Lee, TU, Cho, KO & Kang, MI 2015, 'The prevalence of duck hepatitis A virus types 1 and 3 on Korean duck farms', *Archives of Virology*, vol. 160, no. 2, pp. 493-8, available at <u>https://doi.org/10.1007/s00705-014-</u>2264-3, accessed 4 September 2018.

Song, C, Liao, Y, Gao, W, Yu, S, Sun, Y, Qiu, X, Tan, L, Cheng, A, Wang, M, Ma, Z & Ding, C 2014, 'Virulent and attenuated strains of duck hepatitis A virus elicit discordant innate immune responses in vivo', *Journal of General Virology*, vol. 95, no. 12, pp. 2716-26, available at <u>https://dx.doi.org/10.1099/vir.0.070011-0</u>

Toth, TE & Norcross, NL 1981, 'Humoral immune response of the duck to duck hepatitis virus: virusneutralizing vs. virus-precipitating antibodies', *Avian Diseases*, vol. 25, no. 1, pp. 17-28

Trefilov, BB, Nikitina, NV, Yavdoshak, LI, Dmitriev, KY & Trubitsyn, MM 2018, 'Duck hepatitis virus type I', *European Journal of Natural History*, vol. 2018, no. 1, pp. 3-6, available at <u>https://worldscience.ru/en/article/view?id=33830</u>, accessed 5 September 2018.

Tripathy, DN & Hanson, LE 1986, 'Impact of oral immunization against duck viral hepatitis in passively immune ducklings', *Preventive Veterinary Medicine*, vol. 4, no. 4, pp. 355-60, available at https://doi.org/10.1016/0167-5877(86)90016-4, accessed 23 March 2016.
Tseng, C-H, Knowles, NJ & Tsai, H-J 2007, 'Molecular analysis of duck hepatitis virus type 1 indicates that it should be assigned to a new genus', *Virus Research*, vol. 123, no. 2, pp. 190-203, available at https://doi.org/10.1016/j.virusres.2006.09.007, accessed 3 March 2016.

Wen, XJ, Cheng, AC, Wang, MS, Jia, RY, Zhu, DK, Chen, S, Liu, MF, Liu, F & Chen, XY 2014, 'Detection, differentiation, and VP1 sequencing of duck hepatitis A virus type 1 and type 3 by a 1-step duplex reverse-transcription PCR assay', *Poultry Science*, vol. 93, no. 9, pp. 2184-92, available at https://doi.org/10.3382/ps.2014-04024, accessed 21 March 2016.

Woolcock, PR 1986, 'An assay for duck hepatitis virus type I in duck embryo liver cells and a comparison with other assays', *Avian Pathology*, vol. 15, no. 1, pp. 75-82, available at <u>http://dx.doi.org/10.1080/03079458608436267</u>, accessed 1 July 2009.

-- -- 2003, 'Duck hepatitis', in Saif, YM, Barnes, HJ, Glisson, JR, Fadly, AM, McDougald, LR & Swayne, DE (eds), *Diseases of poultry*, 11th edn, Iowa State Press, Ames, pp. 343-54.

Woolcock, PR & Fabricant, J 1991, 'Duck virus hepatitis', in Calnek, BW, Barnes, HJ, Beard, CW, Reid, WM & Yoder, HW, Jr. (eds), *Diseases of poultry*, 9th edn, Iowa State University Press, Ames, pp. 597-607.

Woolcock, PR & Tsai, H-J 2013, 'Duck hepatitis', in Swayne, DE, Glisson, JR, McDougald, LR, Nolan, LK, Suarez, DL & Nair, VL (eds), *Diseases of poultry*, 13th edn, Wiley-Blackwell, Ames, Iowa, USA, pp. 422-31.

Yang, M, Cheng, A, Wang, M & Xing, H 2008, 'Development and application of a one-step real-time Taqman RT-PCR assay for detection of duck hepatitis virus type1', *Journal of Virological Methods*, vol. 153, no. 1, pp. 55-60, available at <u>https://doi.org/10.1016/j.jviromet.2008.06.012</u>, accessed 8 March 2016.

Zhang, R, Chen, J, Zhang, J, Yang, Y, Li, P, Lan, J, Xie, Z & Jiang, S 2018, 'Novel duck hepatitis A virus type 1 isolates from adult ducks showing egg drop syndrome', *Veterinary Microbiology*, vol. 221, pp. 33-7, available at <u>https://doi.org/10.1016/j.vetmic.2018.05.023</u>, accessed 4 September 2018.

4.2 Duck Tembusu virus

4.2.1 Background

Duck Tembusu virus (DTMUV) is a flavivirus, a genus within the family *Flaviviridae*, which causes a highly infectious disease in ducks. It was first identified in April 2010 in Shanghai, China and spread rapidly to most of the duck producing regions of China, especially in the south-east provinces (Su et al. 2011; Yan et al. 2011). The disease is characterised by a dramatic fall in egg production in both egg laying and breeding ducks, and by diarrhoea, neurological disorders and death in ducklings under two weeks of age (Su et al. 2011; Sun et al. 2014; Thontiravong et al. 2015). Since 2010, the disease has been reported in ducks in Malaysia and Thailand (Homonnay et al. 2014; Thontiravong et al. 2015).

Flaviviruses are positive sense, single-stranded RNA viruses that occur throughout the world. Over 70 different members in the genus Flavivirus have been discovered. Many flaviviruses are transmitted to humans and animals by infected mosquitoes or ticks (Liu et al. 2013).

Tembusu viruses (TMUV) have been reported to infect mosquitoes, chickens, ducks, geese (family *Anatidae*, genera *Anser*, *Branta* and *Chen*), pigeons (*Columba livia*) and sparrows (*Passeridae*) with DTMUV being reported only in a variety of duck breeds, especially the Pekin ducks, Muscovy ducks and domesticated Mallards (Zhang et al. 2017). TMUV is not known to be a zoonotic disease although TMUV antibodies have been detected in workers involved with the duck industry (Tang et al. 2013; Wang et al. 2016).

Infection due to DTMUV is not OIE-listed nor is it nationally notifiable or subject to official control or eradication in Australia.

4.2.2 Technical information

Epidemiology

DTMUV can spread very rapidly within and between flocks and there is clear evidence supporting both direct transmission and spread via a mosquito vector. Direct transmission and airborne transmission were observed to be the primary method of spread of DTMUV (Homonnay et al. 2014; Li et al. 2015b). This is unusual in that this group of flavivirus usually relies on a mosquito vector for spread (Liu et al. 2013). However, evidence of direct transmission is very strong. In China outbreaks have occurred during times when mosquitoes are not active (Yan et al. 2011). Further, high levels of DTMUV have been identified in tracheal cells, enabling direct transmission of virus between birds in close proximity to each other (Cha et al. 2013).

There have also been reports of faecal-oral transmission of DTMUV (Thontiravong et al. 2015).

Mosquito vectors do have a role in transmission as DTMUV genes have been detected in mosquitoes and DTMUV outbreaks in Thailand have been most frequent at times of high mosquito activity (Tang et al. 2015; Thontiravong et al. 2015). TMUV is experimentally transmissible to chickens through at least one member of the *Culex* genus, *C. vishnui* (O'Guinn et al. 2013). *C. vishnui* has not been identified in Australia.

The identification of the virus in other avian species may indicate an ability of the virus to adapt quickly to new hosts (Liu et al. 2013). As flaviviruses are RNA viruses, they are prone to transcription errors that make them more likely than DNA viruses to be modified by the process of replication (Carter & Saunders 2007).

Infection and morbidity rates are very high (up to 90 per cent) and, depending on the management conditions of flocks, mortality typically ranges from 5 to 15 per cent and occasionally up to 30 per cent, especially when there is secondary bacterial infection (Zhang et al. 2017).

Distribution and prevalence

Infection due to DTMUV was first reported in China in 2010. The cause of the outbreaks in China was initially identified as a new flavivirus, Baiyangdian virus, and the disease was called Duck Egg Drop Syndrome (DEDS) (Su et al. 2011). DEDS has been responsible for economic losses through decreased egg production and the deaths of laying ducks in mainland China (Jiang et al. 2012; Liu et al. 2013; Yan et al. 2011). In 2011, DEDS also occurred in several chicken and goose farms in China and a Tembusu-like virus, having homology to the DTMUV, was identified as the cause (Liu et al. 2012).

Outbreaks characterised by neurological signs and significant losses in young Pekin ducks were reported in Malaysia in 2012 with the strain of DTMUV identified being named Perak virus (Homonnay et al. 2014). This strain is around 90 per cent homologous to the strain affecting ducks in Thailand, indicating that Perak virus is the same species as the virus affecting ducks in Thailand and China (Thontiravong et al. 2015). Flaviviruses are defined as being of the same species if they are more than 84 per cent homologous (Kuno et al. 1998). There were earlier reports of a TMUV named Sitiawan virus causing encephalitis in chickens in Malaysia (Kono et al. 2000).

In Thailand, outbreaks of disease due to DTMUV were first reported in layer and broiler duck farms in 2013 with investigations indicating infection occurring as far back as 2010 (Chakritbudsabong et al. 2015; Thontiravong et al. 2015). A strain of DTMUV, KPS54A61, was later identified as causing duck mortalities and reduced egg production in some provinces in Thailand. More recently, it has been proposed that this virus is the same as the DTMUV found in the ducks in China, to which it is 98 per cent homologous (Chakritbudsabong et al. 2015).

Australian Status

Tembusu viruses, including DTMUV, have not been reported in Australia, there is no active surveillance, and tembusu viruses of ducks are not nationally notifiable.

Agent properties

There is no information on the chemo-physical properties of the TMUV. However, flaviviruses share remarkable similarities in virion properties, structure, and genome organisation but vary in biological and antigenic properties (Goo et al. 2016; Kuhn et al. 2002). As the flaviviruses are enveloped viruses and composed of 17 per cent lipids by weight, they are stable in an alkaline environment of pH 8 under laboratory (in vitro) conditions but are sensitive to treatment with organic solvents and detergents, and to high temperature.

In general, flaviviruses are typically inactivated by temperatures above 56°C for at least 30 minutes, in solutions of pH less than or equal to 6, by ultraviolet light and by gamma-radiation, and are known to be susceptible to a range of disinfectants (Charrel et al. 2016). Because it was thought that Zika virus, also a flavivirus, was more stable than other flaviviruses, Müller *et al* (2016) assessed the environmental stability of Zika virus to heat and change in pH. The virus was stable at temperatures up to 50°C but lost all infectivity at temperatures of greater than 60°C (Figure 3). They also observed that Zika virus infectivity was highest after adjusting the stock to a pH of approximately 9 but adjusting to pH 12 or to less than pH 4 abrogated infectivity (Figure 3).





(from Müller et al 2016)

Left graph - Zika virus was incubated for 5 min at indicated temperatures. Temperatures >60°C inactivated the virus. Right graph - Stocks were adjusted to indicated pH values and incubated for 10 min. pH levels <4 or >11 inactivated the virus.

Bovine viral diarrhoea virus, a pestivirus which is also a genus within the family Flaviviridae, is enzootic in most cattle populations throughout the world and is present throughout the body of persistently infected cattle. The virus in whole and ground muscle meat was consistently inactivated when cooked to temperatures greater than or equal to 75°C (Bratcher et al. 2012).

4.2.3 Presence in duck meat

There is no specific information available on DTMUV in duck meat.

Diseased meat ducks, especially 20–40 day old ducklings, typically develop nervous system disorders including unsteady standing, falling and quivering. Older ducks show less severe clinical signs, except for breeder or egg-laying ducks which show severe drop in egg production. Histology has revealed evidence of viral encephalitis. However, necropsy of female infected ducks consistently showed severe ovarian haemorrhage, ovaritis, with occasional ruptured ovarian follicles and peritonitis. Enlarged spleens and leg muscle hemorrhage were occasionally noted (Han et al. 2016). High levels of DTMUV have also been identified in tracheal cells (Cha et al. 2013).

Wu et al (2014) observed that DTMUV in adult male and female ducks was detectable in most of the parenchymatous organs as well as the oviduct and intestinal tract from days one to seven after inoculation. Viral titres were maintained at high levels for at least nine days in the spleen, kidney, bursa of Fabricius, brain and ovary. However, no virus was detected in any of these organs or tissues 18 days after inoculation. Wu et al (2014) concluded that DTMUV causes systemic infection in male and female ducks, and is capable of replication in most organs, with the exception of the ovaries and testes.

4.2.4 Pathogenesis

Flaviviruses enter cells by receptor-mediated endocytosis after attaching to the cell surface. The E protein involved in this process also plays a role in the immune response, including as a target for antibodies (Liu et al. 2013; Murphy et al. 1999). The virus replicates rapidly in cells with cytopathic effects (Murphy et al. 1999). Cells favoured by flaviviruses include cells involved in immunity such as macrophages, monocytes and dendritic cells (Liu et al. 2013). In particular, the spleen has been identified as one of the main target organs for virus replication, based on the appearance at post

mortem (Jiang et al. 2012). There is also a post mortem reduction in lymphocyte numbers in the spleen (Wang et al. 2014). In the case of the ovary, the virus mainly affects the follicle cells, causing degeneration and necrosis of these cells and haemorrhage and congestion in the follicular cavity itself (Wang et al. 2014).

DTMUV is present in the blood four hours after infection and ducks begin shedding the virus in the faeces around eight hours after infection (Wang et al. 2014). Clinical signs may be evident two to three days later (Yan et al. 2011).

4.2.5 Diagnosis

Clinical signs

Sudden onset and rapid spread are features of the disease caused by DTMUV (Zhang et al. 2017). Anorexia is usually the first clinical sign, becoming evident within two to three days of experimental inoculation, or five days following contact with infected ducks (Yan et al. 2011). Within 10 days, egg production drops to lower than 10 per cent (Liu et al. 2013). Other clinical signs include rhinorrhoea, diarrhoea, pyrexia and reduced growth. Neurological signs such as paralysis, ataxia and behavioural changes are also common (Jiang et al. 2012; Liu et al. 2013; Zhang et al. 2017). In drakes, the pathological effects on the testes can lead to an absence of spermatogenesis (Wu et al. 2014).

The age of the infected duck can influence the severity of signs, with very young ducks showing more severe clinical signs and reduced clearance of the virus (Li et al. 2015a). In the outbreaks in Thailand, clinical signs were evident in broiler ducks over three weeks of age and in laying ducks when in lay (Thontiravong et al. 2015). The Malaysian Perak virus outbreak was characterised by neurological signs including lameness, ataxia and progressive paralysis, which were most likely to be observed in young ducks (four to seven weeks of age) (Homonnay et al. 2014).

Morbidity can approach 90 per cent or even 100 per cent in infected populations (Liu et al. 2013; Zhang et al. 2017). Mortality in the Chinese outbreaks was reported to range from 5 per cent to 30 per cent, depending on the supportive treatment provided, with seroconversion occurring in the surviving birds (Liu et al. 2013). A mortality rate of 7.5 per cent was observed in an experimentally infected flock of 80 nine-week-old shelducks (most ducks in genus *Tadorna*) (with 10 per cent mortality in a smaller experiment carried out earlier) (Yan et al. 2011). In the outbreaks in Thailand, mortality was as high as 50 per cent in some groups, with a strong correlation with secondary bacterial infection (Thontiravong et al. 2015).

Pathology

Post-mortem examinations of ducks in the Chinese outbreaks revealed ovarian pathology such as ovaritis, ovarian haemorrhage and regression, with or without ruptured follicles and peritonitis (Liu et al. 2013). Lymphocyte infiltration in the ovaries and portal area interstitial inflammation were also observed (Tang et al. 2012). However, there were often lesions affecting other organs, including severe splenic necrosis or enlargement, pulmonary haemorrhage, moderate necrosis of the brain tissue, and nephrosis (Liu et al. 2013; Yan et al. 2011). Similar ovarian and splenic pathology was seen in ducks in Thailand, with multifocal gliosis and perivascular cuffing identified on histopathological examination of the cerebellum and spinal cord (Thontiravong et al. 2015). In the Perak virus outbreaks in Malaysia, non-purulent panencephalomyelitis was the main feature of the microscopic pathological lesions found in the brain and spinal cord of infected Pekin ducks (Homonnay et al. 2014).

Chicken embryos died three to five days after experimental inoculation of eggs with the virus. Severe cutaneous haemorrhage was evident in the embryos on examination (Thontiravong et al. 2015).

Testing

In endemic regions, presumptive diagnosis can often be made based on the clinical signs, rapid spread through the flock and post-mortem findings (Zhang et al. 2017).

Real-time RT-PCR and reverse transcription loop-mediated isothermal amplification assays can be used to definitively diagnose the virus (Liu et al. 2013). Both methods can detect DTMUV within 30 minutes, with the latter having the advantage of being inexpensive and usable in the field compared to RT-PCR, which is more expensive and requires a higher level of skill to perform (Jiang et al. 2012). Both the RT-PCR and reverse transcription loop-mediated isothermal amplification assay tests have high sensitivity and specificity, with the specificity determined through analysis with other flaviviruses with no evidence of a cross-reaction (Jiang et al. 2012). ELISA tests have also been used, but like RT-PCR, these have the disadvantage of being expensive and requiring laboratory facilities (Wang et al. 2014). A latex-agglutination assay has also been employed and is reported to be specific for DTMUV (Wang et al. 2014).

Duck blood remains positive for viral nucleic acid for up to four weeks following infection, while faeces can be positive for up to 13 days, suggesting they might be used for fluid sampling (Wang et al. 2014). The spleen has been identified as one of the preferred targets for replication of the virus and therefore a good site for post-mortem tissue sampling due to the high viral loads present in infected birds, although the virus can no longer be detected in the spleen by 18 days after infection (Jiang et al. 2012; Wu et al. 2014). The kidneys, brain, ovaries and bursa of Fabricius can also be used, with similar constraints on the sampling period (Wu et al. 2014).

4.2.6 Treatment

There is no effective treatment for DTMUV infection.

4.2.7 Control

Currently there is no commercial vaccine available. In recent years there has been promising research in the development and testing of vaccines for protection against DTMUV. These include an inactivated vaccine injected subcutaneously twice in the neck of ducks, a live attenuated vaccine and an oral DNA vaccine (Huang et al. 2018; Li et al. 2014; Zhang et al. 2017). Combination vaccines are also being developed and trialled, including a trivalent vaccine to prevent avian influenza (H5N1), DTMUV, and duck enteritis virus infections in ducks using the CRISPR/Cas9 system (Zou et al. 2017).

4.2.8 Current biosecurity measures

There is no biosecurity measure against DTMUV for the importation of fertile duck eggs, however the virus has not been identified in countries approved for sourcing.

4.2.9 Conclusion

- DTMUV is not an OIE-listed disease nor is it nationally notifiable in Australia.
- DTMUV is an emerging disease, with outbreaks resulting in significant morbidity and mortality first reported in China in 2010. Since then outbreaks have been reported in Thailand and Malaysia.

- DTMUV is present in Thailand but has not been identified in Australia.
- DTMUV can spread rapidly within and between duck flocks, causing up to 100 per cent morbidity and 30 per cent mortality. It spreads via mosquito vectors, direct transmission and/or airborne transmission.
- There is no treatment or commercially produced vaccine currently available.
- Prevention and control rely mainly on early diagnosis using both serologic and molecular methods, disinfection of premises, removal of birds from the infected environment and depopulation.
- DTMUV causes systemic infection in male and female ducks, and is capable of replication in most organs. Cells favoured by DTMUV include cells involved in immunity such as macrophages, monocytes and dendritic cells.
- DTMUV is no longer present in ducks that survive within 18 days of onset of infection.
- Given that flaviviruses are typically inactivated by temperatures above 56°C for at least 30 minutes, Zika virus is inactivated after five minutes at 60°C and bovine viral diarrhoea virus is inactivated when core temperature reaches 75°C, it is expected that DTMUV would be inactivated when heating duck meat to a minimum core temperature of 70°C for a minimum of 8.2 minutes.

Based on the above information cooking as defined in the scope at section 1.3 of this document (70°C for 8.2 minutes) will inactivate any DTMUV that may be present in meat of ducks. Therefore the department concluded that further risk assessment of duck Tembusu virus was not required.

References

Bratcher, CL, Wilborn, BS, Finegan, HM, Rodning, SP, Galik, PK, Riddell, KP, Marley, MS, Zhang, Y, Bell, LN & Givens, MD 2012, 'Inactivation at various temperatures of bovine viral diarrhea virus in beef derived from persistently infected cattle', *Journal of Animal Science*, vol. 90, no. 2, pp. 635-41, available at doi: 10.2527/jas.2011-4232, accessed 29 January 2016.

Carter, J & Saunders, V 2007, Virology: principles and applications, Wiley, Hoboken, New Jersey, USA.

Cha, S-Y, Kang, M, Moon, O-K, Park, C-K & Jang, H-K 2013, 'Respiratory disease due to current egg drop syndrome virus in Pekin ducks', *Veterinary Microbiology*, vol. 165, no. 3, pp. 305-11, available at https://doi.org/10.1016/j.vetmic.2013.04.004, accessed 15 August 2018.

Chakritbudsabong, W, Taowan, J, Lertwatcharasarakul, P, Phattanakunanan, S, Munkhong, A, Songserm, T & Chaichoun, K 2015, 'Genomic characterization of a new Tembusu flavivirus from domestic ducks in Thailand', *Thai Journal of Veterinary Medicine*, vol. 45, no. 3, pp. 419-25, available at <u>https://tci-thaijo.org/index.php/tjvm/article/view/39819</u>, accessed 20 January 2016.

Charrel, RN, Leparc-Goffart, I, Pas, S, de Lamballerie, X, Koopmans, M & Reusken, C 2016, 'Background review for diagnostic test development for Zika virus infection', *Bulletin of the World Health Organization*, vol. 94, no. 8, pp. 574-84, available at <u>https://dx.doi.org/10.2471%2FBLT.16.171207</u>, accessed 15 August 2018. Goo, L, Dowd, KA, Smith, ARY, Pelc, RS, DeMaso, CR & Pierson, TC 2016, 'Zika virus is not uniquely stable at physiological temperatures compared to other flaviviruses', *mBio*, vol. 7, no. 5, pp. e01396-16, available at <u>https://doi.org/10.1128/mBio.01396-16</u>, accessed 15 August 2018.

Han, K, Zhao, D, Liu, Y, Liu, Q, Huang, X, Yang, J, An, F & Li, Y 2016, 'Quantitative proteomic analysis of duck ovarian follicles infected with duck Tembusu virus by label-free LC-MS', *Frontiers in Microbiology*, vol. 7, pp. 463, available at <u>https://doi.org/10.3389/fmicb.2016.00463</u>, accessed 15 August 2018.

Homonnay, ZG, Kovács, EW, Bányai, K, Albert, M, Fehér, E, Mató, T, Tatár-Kis, T & Palya, V 2014, 'Tembusu-like flavivirus (Perak virus) as the cause of neurological disease outbreaks in young Pekin ducks', *Avian Pathology*, vol. 43, no. 6, pp. 552-60, available at <u>https://doi.org/10.1080/03079457.2014.973832</u>, accessed 14 August 2018.

Huang, J, Shen, H, Jia, R, Wang, M, Chen, S, Zhu, D, Liu, M, Zhao, X, Yang, Q, Wu, Y, Liu, Y, Zhang, L, Yin, Z, Jing, B & Cheng, A 2018, 'Oral vaccination with a DNA vaccine encoding capsid protein of duck Tembusu virus induces protection immunity', *Viruses*, vol. 10, no. 4, pp. 180, available at https://doi.org/10.3390/v10040180, accessed 14 August 2018.

Jiang, T, Liu, J, Deng, Y-Q, Su, J-L, Xu, L-J, Liu, Z-H, Li, X-F, Yu, X-D, Zhu, S-Y, Gao, GF, Qin, E-D & Qin, C-F 2012, 'Development of RT-LAMP and real-time RT-PCR assays for the rapid detection of the new duck Tembusu-like BYD virus', *Archives of Virology*, vol. 157, no. 12, pp. 2273-80, available at <u>https://doi.org/10.1007/s00705-012-1431-7</u>, accessed 19 January 2016.

Kono, Y, Tsukamoto, K, Abd Hamid, M, Darus, A, Lian, TC, Sam, LS, Yok, CN, Di, KB, Lim, KT, Yamaguchi, S & Narita, M 2000, 'Encephalitis and retarded growth of chicks caused by Sitiawan virus, a new isolate belonging to the genus Flavivirus', *The American Journal of Tropical Medicine and Hygiene*, vol. 63, no. 1, pp. 94-101, available at <u>https://doi.org/10.4269/ajtmh.2000.63.94</u>, accessed 27 January 2016.

Kuhn, RJ, Zhang, W, Rossmann, MG, Pletnev, SV, Corver, J, Lenches, E, Jones, CT, Mukhopadhyay, S, Chipman, PR, Strauss, EG, Baker, TS & Strauss, JH 2002, 'Structure of dengue virus: Implications for flavivirus organization, maturation, and fusion', *Cell*, vol. 108, no. 5, pp. 717-25, available at https://doi.org/10.1016/S0092-8674(02)00660-8, accessed 30 October 2018.

Kuno, G, Chang, G-JJ, Tsuchiya, KR, Karabatsos, N & Cropp, CB 1998, 'Phylogeny of the genus Flavivirus', *Journal of Virology*, vol. 72, no. 1, pp. 73-83, available at http://jvi.asm.org/cgi/content/abstract/72/1/73, accessed 19 January 2016.

Li, G, Gao, X, Xiao, Y, Liu, S, Peng, S, Li, X, Shi, Y, Zhang, Y, Yu, L, Wu, X, Yan, P, Yan, L, Teng, Q, Tong, G & Li, Z 2014, 'Development of a live attenuated vaccine candidate against duck Tembusu viral disease', *Virology*, vol. 450-451, pp. 233-42, available at <u>https://doi.org/10.1016/j.virol.2013.12.028</u>, accessed 19 January 2016.

Li, N, Wang, Y, Li, R, Liu, J, Zhang, J, Cai, Y, Liu, S, Chai, T & Wei, L 2015a, 'Immune responses of ducks infected with duck Tembusu virus', *Frontiers in Microbiology*, vol. 6, pp. 425, available at https://doi.org/10.3389/fmicb.2015.00425, accessed 14 August 2018.

Li, X, Shi, Y, Liu, Q, Wang, Y, Li, G, Teng, Q, Zhang, Y, Liu, S & Li, Z 2015b, 'Airborne transmission of a novel Tembusu virus in ducks', *Journal of Clinical Microbiology*, vol. 53, no. 8, pp. 2734-6, available at doi: 10.1128/jcm.00770-15, accessed 14 Augsut 2018.

Liu, M, Chen, S, Chen, Y, Liu, C, Chen, S, Yin, X, Li, G & Zhang, Y 2012, 'Adapted Tembusu-like virus in chickens and geese in China', *Journal of Clinical Microbiology*, vol. 50, no. 8, pp. 2807-9, available at doi: 10.1128/jcm.00655-12, accessed 28 August 2018.

Liu, P, Lu, H, Li, S, Wu, Y, Gao, GF & Su, J 2013, 'Duck egg drop syndrome virus: an emerging Tembusu-related flavivirus in China', *Science China Life Sciences*, vol. 56, no. 8, pp. 701-10, available at <u>https://doi.org/10.1007/s11427-013-4515-z</u>, accessed 15 January 2016.

Müller, JA, Harms, M, Schubert, A, Jansen, S, Michel, D, Mertens, T, Schmidt-Chanasit, J & Münch, J 2016, 'Inactivation and environmental stability of Zika Virus', *Emerging Infectious Diseases*, vol. 22, no. 9, pp. 1685-7, available at <u>https://dx.doi.org/10.3201/eid2209.160664</u>, accessed 15 August 2018.

Murphy, FA, Gibbs, EPJ, Horzinek, MC & Studdert, MJ 1999, *Veterinary virology*, 3rd edn, Academic Press, San Diego.

O'Guinn, ML, Turell, MJ, Kengluecha, A, Jaichapor, B, Kankaew, P, Miller, RS, Endy, TP, Jones, JW, Coleman, RE & Lee, JS 2013, 'Field detection of tembusu virus in western Thailand by RT-PCR and vector competence determination of select culex mosquitoes for transmission of the virus', *The American Journal of Tropical Medicine and Hygiene*, vol. 89, no. 5, pp. 1023-8, available at https://doi.org/10.4269/ajtmh.13-0160, accessed 30 October 2018.

Su, J, Li, S, Hu, X, Yu, X, Wang, Y, Liu, P, Lu, X, Zhang, G, Hu, X, Liu, D, Li, X, Su, W, Lu, H, Mok, NS, Wang, P, Wang, M, Tian, K & Gao, G, F. 2011, 'Duck egg-drop syndrome caused by BYD virus, a new Tembusu-related flavivirus', *PLos One*, vol. 6, no. 3, pp. e18106, available at https://doi.org/10.1371/journal.pone.0018106, accessed 19 January 2016.

Sun, XY, Diao, YX, Wang, J, Liu, X, Lu, AL, Zhang, L, Ge, PP & Hao, DM 2014, 'Tembusu virus infection in Cherry Valley ducks: the effect of age at infection', *Veterinary Microbiology*, vol. 168, no. 1, pp. 16-24, available at <u>https://doi.org/10.1016/j.vetmic.2013.10.003</u>, accessed 14 August 2018.

Tang, Y, Diao, Y, Chen, H, Ou, Q, Liu, X, Gao, X, Yu, C & Wang, L 2015, 'Isolation and genetic characterization of a Tembusu virus strain isolated from mosquitoes in Shandong, China', *Transboundary and Emerging Diseases*, vol. 62, no. 2, pp. 209-16, available at https://doi.org/10.1111/tbed.12111, accessed 28 January 2016.

Tang, Y, Diao, Y, Gao, X, Yu, C, Chen, L & Zhang, D 2012, 'Analysis of the complete genome of Tembusu virus, a flavivirus isolated from ducks in China', *Transboundary and Emerging Diseases*, vol.

59, no. 4, pp. 336-43, available at <u>https://doi.org/10.1111/j.1865-1682.2011.01275.x</u>, accessed 28 January 2016.

Tang, Y, Gao, X, Diao, Y, Feng, Q, Chen, H, Liu, X, Ge, P & Yu, C 2013, 'Tembusu virus in human, China', *Transboundary and Emerging Diseases*, vol. 60, no. 3, pp. 193-6, available at https://doi.org/10.1111/tbed.12085, accessed 14 August 2018.

Thontiravong, A, Ninvilai, P, Tunterak, W, Nonthabenjawan, N, Chaiyavong, S, Angkabkingkaew, K, Mungkundar, C, Phuengpho, W, Oraveerakul, K & Amonsin, A 2015, 'Tembusu-related flavivirus in ducks, Thailand', *Emerging Infectious Diseases*, vol. 21, no. 12, pp. 2164-7, available at https://doi.org/10.3201/eid2112.150600, accessed 15 January 2016.

Wang, H-J, Li, X-F, Liu, L, Xu, Y-P, Ye, Q, Deng, Y-Q, Huang, X-Y, Zhao, H, Qin, ED, Shi, P-Y, Gao, GF & Qin, C-F 2016, 'The emerging duck flavivirus is not pathogenic for primates and is highly sensitive to mammalian interferon antiviral signaling', *Journal of Virology*, vol. 90, no. 14, pp. 6538-48, available at <u>https://dx.doi.org/10.1128%2FJVI.00197-16</u>, accessed 14 August 2018.

Wang, Q, Wen, Y, Huang, Y, Wu, Y, Cai, Y, Xu, L, Wang, C, Li, A, Wu, B & Chen, J 2014, 'Isolation and identification of duck Tembusu virus strain LH and development of latex-agglutination diagnostic method for rapid detection of antibodies', *Avian Diseases*, vol. 58, no. 4, pp. 616-22, available at <u>https://doi.org/10.1637/10795-021114-Reg</u>, accessed 30 October 2018.

Wu, L, Liu, J, Chen, P, Jiang, Y, Ding, L, Lin, Y, Li, Q, He, X, Chen, Q & Chen, H 2014, 'The sequential tissue distribution of duck Tembusu virus in adult ducks', *BioMed Research International*, vol. 2014, pp. 703930, available at <u>http://dx.doi.org/10.1155/2014/703930</u>, accessed 19 January 2016.

Yan, P, Zhao, Y, Zhang, X, Xu, D, Dai, X, Teng, Q, Yan, L, Zhou, J, Ji, X, Zhang, s, Liu, G, Zhou, Y, Kawaoka, Y, Tong, G & Li, Z 2011, 'An infectious disease of ducks caused by a newly emerged Tembusu virus strain in mainland China', *Virology*, vol. 417, no. 1, pp. 1-8, available at https://doi.org/10.1016/j.virol.2011.06.003, accessed 15 January 2016.

Zhang, W, Chen, S, Mahalingam, S, Wang, M & Cheng, A 2017, 'An updated review of avian-origin Tembusu virus: a newly emerging avian Flavivirus', *Journal of General Virology*, vol. 98, no. 10, pp. 2413-20, available at https://dx.doi.org/10.1099/jgv.0.000908, accessed 14 August 2018.

Zou, Z, Huang, K, Wei, Y, Chen, H, Liu, Z & Jin, M 2017, 'Construction of a highly efficient CRISPR/Cas9-mediated duck enteritis virus-based vaccine against H5N1 avian influenza virus and duck Tembusu virus infection', *Scientific Reports*, vol. 7, pp. 1478, available at https://dx.doi.org/10.1038%2Fs41598-017-01554-1

4.3 Duck virus enteritis (duck plague)

4.3.1 Background

Duck virus enteritis (DVE) or duck plague is caused by a contagious herpesvirus, the duck virus enteritis virus (DVEV). DVE is an acute, though sometimes chronic disease of waterfowl of the family Anatidae (ducks, swans and geese). Natural infections occur in ducks from seven days of age to adults, and causes significant economic losses (Hansen & Gough 2007). Ducklings infected with a low virulence strain often suffer from secondary bacterial infections due to the virus having an immunosuppressive effect (Campagnolo et al. 2001; Shawky, Sandhu & Shivaprasad 2000). Laying ducks can suffer from significant reductions in egg production and adult males may suffer from prolapse of the penis (Woolcock & Tsai 2013).

DVE was first identified in 1923 in the Netherlands, and in the 1940s was termed 'duck plague' when the disease could not be reproduced in species other than Anatidae (Woolcock & Tsai 2013). The virus has subsequently been confirmed in Austria, Belgium, Canada, China, Denmark, England, France, Germany, Hungary, India, Poland, Thailand, the United States of America and Vietnam (Dhama et al. 2017).

DVE is not an OIE-listed disease, having been removed from the list in 2010 due to its limited spread across country borders. DVE is not nationally notifiable nor subject to official control or eradication in Australia.

4.3.2 Technical information

Epidemiology

Susceptible species

With natural infection, members of the Anatidae family are the only species susceptible to DVEV; however, susceptibility varies greatly among the different species (Friend 1999). High mortalities have been reported in a variety of domestic ducks (*Anas platyrhynchos*) and Muscovy ducks (*Cairina moschata*). Despite the widespread distribution and frequent occurrence of duck plague in captive and wild waterfowl in North America, wild waterfowl have only been infrequently infected.

Mallards are more resistant to the lethal effects yet are considered to be a possible natural reservoir of infection (Dhama et al. 2017). In the northern hemisphere, Mallards are usually migratory, but in the southern hemisphere, where they have been introduced, they are not known to be migratory.

Other types of aquatic birds do not become infected. Outbreaks in domestic ducks generally occur when they have access to aquatic environments co-habited by wild ducks, geese and/or swans (Woźniakowski & Samorek-Salamonowicz 2014).

The virus only replicates in cell cultures obtained from the Anseriformes and Galliformes orders. The virus can be modified to grow in chickens less than two weeks of age as well as embryonating chicken eggs (Campagnolo et al. 2001; Hansen & Gough 2007).

Modes of transmission

Transmission occurs by direct contact between susceptible and infected birds, with oral transmission being the most common route. Transmission via indirect contact can also occur with contaminated

water and movement of infected birds being the most likely cause of exposure in both domestic and wild populations. Wild ducks have been implicated as the carriers, thus playing a crucial role in DVEV transmission within and between continents. Both domestic and wild populations have been found to shed the virus periodically when recovering from disease (Campagnolo et al. 2001; Dhama et al. 2017; Woolcock & Tsai 2013).

Incubation period

Ducks as young as one week old can be infected by DVEV (Dhama et al. 2017). The birds can become infected as a result of direct exposure to infected birds or indirectly from contaminated environments, including water. The incubation period of DVEV ranges from three to seven days, with a maximum of 14 days (Campagnolo et al. 2001; Woolcock & Tsai 2013). After overt clinical signs appear, death occurs within one to five days. The mortality and morbidity of the flock can range between 5 and 100 per cent depending on the health of the flock and the virulence of the strain. Those that survive can become carriers of the infection for several years, with infection latent in the trigeminal ganglion, where upon re-activation, shedding occurs periodically (Burgess & Yuill 1983).

A study by Burgess et al (1979) found that all species tested (including Mallard, black, pintail and gadwall ducks) had a high frequency of DVEV shedding. This study also determined that oral erosions did not signify the occurrence of shedding, rather that they are an indicator of persistent infection, which may occur latent in the trigeminal ganglion. Long term persistence of the virus has been suggested with reports of DVEV being isolated from Pekin ducks 45 days after initial infection. Butterfield and Dardiri (1969) also recovered DVEV 17 days after infection from the cloacae of clinically healthy waterfowl (Burgess, Ossa & Yuill 1979; Converse & Kidd 2001).

Distribution and prevalence

DVEV is considered endemic in domestic waterfowl in many Asian countries including Thailand and Vietnam (Hansen & Gough 2007; Woolcock & Tsai 2013).

Australian Status

DVEV has not been reported in Australia and there is no active surveillance. However, infection with DVEV is nationally notifiable.

Agent properties

The disease is caused by DVEV, or Anatid herpesvirus 1, which is a double-stranded DNA virus. It is classified as the type strain of the genus *Mardivirus*, in the subfamily Alphaherpesvirinae and family Herpesviridae (Converse & Kidd 2001). Differences in virulence occur amongst strains, though immunologically they are identical (Akter et al. 2004; Dhama et al. 2017).

Herpesviruses are enveloped virus, with the envelope containing considerable amounts of lipid. Because of this structure, herpesviruses are susceptible to chemical agents, especially those that are lipid solvents (Maris 1995).

Salts and proteins affect the thermostability of herpesviruses, being thermolabile in tissue cultures, extremely thermosensitive in isotonic solutions, but thermostable at 50°C by simple dilution in distilled water, and by 1 M Na₂SO4 or Na₂HPO4 (Wallis & Melnick 1965). Strains of herpesvirus hominis have been reported to resist thermal inactivation in the presence of sodium ions (Wallis & Melnick 1965).

Using heat during inactivation experiments, 10 minutes at 56°C or 90 to 120 minutes at 50°C, nullifies the infectivity of DVEV (Sandhu 2003). Laboratories handling live viruses generally recognise that herpesviruses are inactivated within 30 minutes at 56°C (Public Health Agency of Canada 2011).

A live attenuated DVEV strain used in a vaccine was observed to be non-infective above 42°C within two hours and had become unstable within two hours above 65°C (Makhija & Kumar 2017).

Short-term pasteurisation of human breast milk, that is, five seconds at 72°C, was able to completely destroy the infectivity of cytomegalovirus, a virus in the order Herpesvirales (Hamprecht et al. 2004). Furthermore a cell culture supernatant containing koi herpesvirus was inactivated completely when heated to temperatures above 50°C for one minute (Kasai, Muto & Yoshimizu 2005).

4.3.3 Presence in duck meat

The virus replicates mainly in the mucosa of the digestive tract, then spreads to the bursa of Fabricius, thymus, spleen and liver where it continues to replicate. More specifically, DVEV replicates in the epithelial cells, lymphocytes and macrophages in these organs (Dhama et al. 2017).

Immunoperoxidase staining of a DVEV protein, UL51, enabled the identification of DVEV antigen in the bursa of Fabricius, thymus, spleen, liver, oesophagus, small intestine and large intestine of DVEV-infected ducks, and less so in the Harder's glands, glandularis ventriculus, cerebrum, kidney, lung, pancreas and myocardium. The DVEV protein was also expressed in the cytoplasm of various types of cells, most abundantly in the cytoplasm of lymphocytes, reticulum cells, macrophages, epithelial cells and hepatocytes. However, throughout the infection process UL51 was not seen in the muscle (Shen et al. 2010).

4.3.4 Pathogenesis

The pathologic response is dependent upon age, stage of infection, sex, species affected as well as virus virulence and intensity (Leibovitz 1969; Leibovitz & Hwang 1968; OIE 2018). Mortality rates differ between isolates of DVEV as well as between and amongst species. The original O strain of the virus studied by Jansen (1961) had 100 per cent mortality, though a virus recovered from a 1959 outbreak in the Netherlands caused less than 80 per cent mortality, which could suggest that virus virulence can decrease over time (Hansen & Gough 2007).

Replication of DVEV occurs within the host cell's nucleus. During the late stages of infection, the replicated DNA is packaged into capsids so that it can move into extracellular spaces (Johnson & Baines 2011).

Initial lesions observed in ducks were in the stratified squamous mucosa of the gastrointestinal tract (Proctor 1975). Lesions then develop in the submucosal lymphoid tissue, intestinal villous lamina propria and intestinal crypt epithelium. Infection then proceeds towards the thymus, bursa of Fabricius, spleen and liver. Lesions in these tissues have also been described (Dhama et al. 2017). After viral replication, the epithelial cells containing the viral antigens begin to degenerate (Proctor 1975).

Degenerative changes of parenchymatous organs and generalised haemorrhages are the result of vascular damage induced by DVEV. It has been suggested that lymphoid depletion caused by the necrosis and apoptosis of lymphocytes may lead to the virus causing immunosuppression, which would explain the high presence of secondary infections seen in natural outbreaks (Dhama et al. 2017).

4.3.5 Diagnosis

Diagnosis is based on a combination of clinical signs, gross pathology and histopathology, supported by identification of the virus by either isolation or polymerase chain reaction (PCR) (OIE 2018).

Clinical signs

High mortality is commonly the first observation in domestic flocks as there are few clinical signs. Sometimes carcasses are found floating on the water surface (Dhama et al. 2017).

Further signs of infection are extreme thirst, anorexia, ataxia, nasal discharge, ruffled feathers, watery diarrhoea, half-closed and swollen eyelids, droopiness, solid vents, inability to stand and tremors of head, neck and body. Clinical signs may develop as the infection progresses (Akter et al. 2004; OIE 2018; Woolcock & Tsai 2013). Sick birds may maintain an upright stance by using their wings for support, but their overall appearance is one of weakness and depression (OIE 2018).

Young ducklings may develop conjunctivitis, nasal exudate, lacrimation, blood-staining of the vent, dehydration and weight loss (Akter et al. 2004; OIE 2018; Woolcock & Tsai 2013).

Pathology

Internally, infection is characterised by lesions of lymphoid organs, tissue haemorrhages, vascular damage, digestive mucosal eruptions and degenerative changes in parenchymatous organs (Woolcock & Tsai 2013; Zhang et al. 2010).

Lesions can range from 1 to 10mm in length and are found through the entire digestive tract from the oral cavity to the cloaca. A haemorrhagic surface covers the wall of the lumen of bursa of Fabricius, and there are haemorrhages in the liver in the early stages before the liver becomes a dark bronze colour later in the infection process (OIE 2018; Woolcock & Tsai 2013).

DVEV affects ducklings slightly differently to mature birds, with prominent lymphoid lesions and less tissue haemorrhage, while mature domestic ducks have the primary lesions on the reproductive tract as well as tissue haemorrhages (Woolcock & Tsai 2013). This difference may account for the higher rate of secondary bacterial infections in ducklings.

Parenchymatous and lymphoid organs as well as the gastrointestinal tract will show mucosa with necrotic degenerative changes and disseminated intravascular coagulopathy. Petechiae may be found over the surface of the intestines, kidney, pancreas, liver and lungs, with extravasations of blood found in visceral organs and the myocardium. The ovarian follicles of laying females may be discoloured and deformed, and haemorrhaging of the ovary which fills the abdominal cavity may be present (Woolcock & Tsai 2013).

The vascular damage caused by DVEV results in degenerative changes for the affected tissues. Those without gross lesions may have microscopic changes such as haemorrhages in capillaries within the intestinal villi as well as in venules near the liver and lungs (Woolcock & Tsai 2013).

Testing

The OIE Terrestrial Manual provides recommendations for diagnostic tests for DVEV in ducks (OIE 2018).

Clinical and gross findings may provide a presumptive diagnosis; however, detection of viral antigen or virus isolation is necessary for confirmation (Woolcock & Tsai 2013). The real-time PCR and conventional PCR are sensitive tests for identifying DVEV antigen.

The virus isolation test, along with PCR, is best for confirmation of clinical cases using samples of liver, spleen or kidney tissue. However, the usefulness of the virus isolation test is severely limited for confirming population freedom from infection.

Antigen tests not described in the OIE Manual include the antigen-capture ELISA which is reported to be a sensitive and reliable method for detection of the DVEV antigen (Dhama et al. 2017; Jia et al. 2009).

Another method of detecting DVEV is a loop-mediated isothermal amplification assay. It is a simple yet specific method that is superior to the PCR method and can be used for on-farm diagnosis (Ji et al. 2009).

The recommended tests for determining population freedom are the microtitre plate virus neutralisation assay and neutralisation assay in duck embryos. These are also the recommended tests for determining the post vaccination immune status in individual animals or populations (OIE 2018).

4.3.6 Treatment

There is no effective treatment for DVEV infection.

4.3.7 Control

In most European countries and the United States of America, both live attenuated and killed vaccines are used in broiler and breeder ducks over two weeks old (Dhama et al. 2017). The live attenuated vaccines are not ideal because of the risk of latency; however, there is no evidence of spread by contact from vaccinated to unvaccinated ducklings.

4.3.8 Current biosecurity measures

There are current biosecurity measures in place for this disease in the *Policy review: importation of hatching (fertile) duck eggs from approved countries* (Biosecurity Australia 2009). However, as the duck egg policy review required that the youngest bird not be less than 40 weeks old when eggs are collected for export, these measures are not relevant for meat ducks which are slaughtered at seven to eight weeks of age.

4.3.9 Conclusion

- DVE is present in ducks in Thailand but has not been identified in Australia. It is not an OIE-listed disease nor is it nationally notifiable in Australia.
- DVEV is transmissible by direct contact with oral transmission being the most common route. Transmission via indirect contact can also occur through contaminated water.
- Whilst DVEV is not found in muscle tissue it is found in most organs within the abdominal and thoracic cavities, head and in some white blood cells. There is the potential for infected white blood cells to be present in muscle tissue.
- Using heat during inactivation experiments, 10 minutes at 56°C or 90 to 120 minutes at 50°C, renders DVEV inactive.

• In general, laboratories handling live viruses recognise that herpesviruses are inactivated within 30 minutes at 56°C.

Based on the above information cooking as defined in the scope at section 1.3 of this document (70°C for 8.2 minutes) will inactivate any DVEV that may be present in meat of ducks. Therefore the department concluded that further risk assessment of duck virus enteritis was not required.

References

Akter, S, Islam, MA, Hossain, MT, Begum, MIA, Amin, MM & Sadekuzzaman, M 2004, 'Characterization and pathogenicity of duck plague virus isolated from natural outbreaks in ducks of Bangladesh', *Bangladesh Journal of Veterinary Medicine*, vol. 2, no. 2, pp. 107-11, available at <u>http://dx.doi.org/10.3329/bjvm.v2i2.2540</u>, accessed 24 February 2016.

Biosecurity Australia 2009, *Importation of hatching (fertile) duck eggs from approved countries – policy review*, Canberra, available at <u>http://agriculture.gov.au/biosecurity/risk-analysis/animal/fertile-duck-eggs</u>.

Burgess, EC, Ossa, J & Yuill, TM 1979, 'Duck plague: a carrier state in waterfowl', *Avian Diseases*, vol. 23, no. 4, pp. 940-9, available at doi: 10.2307/1589610, accessed 29 February 2016.

Burgess, EC & Yuill, TM 1983, 'The influence of seven environmental and physiological factors on duck plague virus shedding by carrier mallards', *Journal of Wildlife Diseases*, vol. 19, no. 2, pp. 77-81, available at http://www.jwildlifedis.org/cgi/content/abstract/19/2/77

Butterfield, WK & Dardiri, AH 1969, 'Serologic and immunologic response of ducks to inactivated and attenuated duck plague virus', *Avian Diseases*, vol. 13, no. 4, pp. 876-87

Campagnolo, ER, Banerjee, M, Panigrahy, B & Jones, RL 2001, 'An outbreak of duck viral enteritis (duck plague) in domestic Muscovy ducks (Cairina moschata domesticus) in Illinois', *Avian Diseases*, vol. 45, no. 2, pp. 522-8, available at doi: 10.2307/1592999, accessed 29 February 2016.

Converse, KA & Kidd, GA 2001, 'Duck plague epizootics in the United States, 1967–1995', *Journal of Wildlife Diseases*, vol. 37, no. 2, pp. 347-57, available at <u>https://doi.org/10.7589/0090-3558-37.2.347</u>, accessed 29 February 2016.

Dhama, K, Kumar, N, Saminathan, M, Tiwari, R, Karthik, K, Kumar, MA, Palanivelu, M, Shabbir, MZ, Malik, YS & Singh, RK 2017, 'Duck virus enteritis (duck plague) – a comprehensive update', *Veterinary Quarterly*, vol. 37, no. 1, pp. 57-80, available at <u>https://doi.org/10.1080/01652176.2017.1298885</u>, accessed 8 August 2018.

Friend, M 1999, 'Duck plague', in Friend, M & Franson, JC (eds), *Field manual of wildlife diseases—general field procedures and diseases of birds*, Information and Technology Report 1999-0001 edn, US Geological Survey, Washington, DC, available at

https://pubs.er.usgs.gov/publication/itr19990001, accessed 23 August 2018.

Hamprecht, K, Maschmann, J, Müller, D, Dietz, K, Besenthal, I, Goelz, R, Middleldorp, JM, Speer, CP & Jahn, G 2004, 'Cytomegalovirus (CMV) inactivation in breast milk: reassessment of pasteurization and freeze-thawing', *Pediatric Research*, vol. 56, no. 4, pp. 529-35, available at https://doi.org/10.1203/01.PDR.0000139483.35087.BE, accessed 31 October 2018.

Hansen, WR & Gough, RE 2007, 'Duck plague (duck virus enteritis)', in Thomas, NJ, Hunter, DB & Atkinson, CT (eds), *Infectious diseases of wild birds*, Blackwell Publishing, Ames, Iowa, USA, pp. 87-107.

Jansen, J 1961, 'Duck plague', *British Veterinary Journal*, vol. 117, no. 8, pp. 349-56, available at <u>https://doi.org/10.1016/S0007-1935(17)43502-0</u>, accessed 31 October 2018.

Ji, J, Du, LQ, Xie, QM, Cao, YC, Zuo, KJ, Xue, CY, Ma, JY, Chen, F & Bee, YZ 2009, 'Rapid diagnosis of duck plagues virus infection by loop-mediated isothermal amplification', *Research in Veterinary Science*, vol. 87, no. 1, pp. 53-8, available at <u>https://doi.org/10.1016/j.rvsc.2008.11.003</u>, accessed 1 March 2016.

Jia, R, Cheng, A, Wang, M, Qi, X, Zhu, D, Ge, H, Luo, Q, Liu, F, Guo, Y & Chen, X 2009, 'Development and evaluation of an antigen-capture ELISA for detection of the UL24 antigen of the duck enteritis virus, based on a polyclonal antibody against the UL24 expression protein', *Journal of Virological Methods*, vol. 161, no. 1, pp. 38-43, available at https://doi.org/10.1016/j.jviromet.2009.05.011, accessed 1 March 2016.

Johnson, DC & Baines, JD 2011, 'Herpesviruses remodel host membranes for virus egress', *Nature Reviews Microbiology*, vol. 9, pp. 382-94, available at <u>http://dx.doi.org/10.1038/nrmicro2559</u>, accessed 1 March 2016.

Kasai, H, Muto, Y & Yoshimizu, M 2005, 'Virucidal effects of ultraviolet, heat treatment and disinfectants against koi herpesvirus (KHV)', *Fish Pathology*, vol. 40, no. 3, pp. 137-8, available at <u>https://doi.org/10.3147/jsfp.40.137</u>, accessed 31 October 2018.

Leibovitz, L 1969, 'The comparative pathology of duck plague in wild Anseriformes', *The Journal of Wildlife Management*, vol. 33, no. 2, pp. 294-303, available at doi: 10.2307/3799829, accessed 29 February 2016.

Leibovitz, L & Hwang, J 1968, 'Duck plague on the American continent', *Avian Diseases*, vol. 12, no. 2, pp. 361-78, available at doi: 10.2307/1588237, accessed 25 February 2018.

Makhija, A & Kumar, S 2017, 'Characterization of duck plague virus stability at extreme conditions of temperature, pH and salt concentration', *Biologicals*, vol. 45, pp. 102-5, available at <u>https://doi.org/10.1016/j.biologicals.2016.09.009</u>, accessed 22 August 2018.

Maris, P 1995, 'Modes of action of disinfectants', *Revue Scientifique et Technique de l'Office International des Epizooties*, vol. 14, no. 1, pp. 47-55

OIE 2018, 'Duck virus enteritis', in *Manual of diagnostic tests and vaccines for terrestrial animals 2018*, World Organisation for Animal Health, Paris, available at <u>http://www.oie.int/standard-setting/terrestrial-manual/access-online/</u>, accessed 17 September 2018.

Proctor, SJ 1975, 'Pathogenesis of digestive tract lesions in duck plague', *Veterinary Pathology*, vol. 12, no. 5-6, pp. 349-61, available at <u>https://doi.org/10.1177/0300985875012005-00602</u>, accessed 29 February 2016.

Public Health Agency of Canada 2011, 'Pathogen safety data sheets: infectious substances – herpes simplex virus', Government of Canada, Ottawa accessed 12 December 2018.

Sandhu, T 2003, 'Diseases of ducks', available at http://soyasa.org/documents/1/Sadhu_P_Diseases_of_Ducks_Apr03.htm.

Shawky, S, Sandhu, T & Shivaprasad, HL 2000, 'Pathogenicity of a low-virulence duck virus enteritis isolate with apparent immunosuppressive ability', *Avian Diseases*, vol. 44, no. 3, pp. 590-9, available at doi: 10.2307/1593098, accessed 29 February 2018.

Shen, C, Cheng, A, Wang, M, Xu, C, Jia, R, Chen, X, Zhu, D, Luo, Q, Cui, H, Zhou, Y, Wang, Y, Xu, Z, Chen, Z & Wang, X 2010, 'Expression and distribution of the duck enteritis virus UL51 protein in experimentally infected ducks', *Avian Diseases*, vol. 54, no. 2, pp. 939-47, available at https://doi.org/10.1637/9172-112109-ResNote.1, accessed 23 August 2018.

Wallis, C & Melnick, JL 1965, 'Thermostabilization and thermosensitization of herpesvirus', *Journal of Bacteriology*, vol. 90, no. 6, pp. 1632-7, available at <u>https://jb.asm.org/content/90/6/1632</u>, accessed 22 August 2018.

Woolcock, PR & Tsai, H-J 2013, 'Duck hepatitis', in Swayne, DE, Glisson, JR, McDougald, LR, Nolan, LK, Suarez, DL & Nair, VL (eds), *Diseases of poultry*, 13th edn, Wiley-Blackwell, Ames, Iowa, USA, pp. 422-31.

Woźniakowski, G & Samorek-Salamonowicz, G 2014, 'First survey of the occurrence of duck enteritis virus (DEV) in free-ranging Polish water birds', *Archives of Virology*, vol. 159, no. 6, pp. 1439-44, available at <u>https://doi.org/10.1007/s00705-013-1936-8</u>, accessed 23 August 2018.

Zhang, S, Xiang, J, Cheng, A, Wang, M, Li, X, Li, L, Chen, X, Zhu, D, Luo, Q & Chen, X 2010, 'Production, purification and characterization of polyclonal antibody against the truncated gK of the duck enteritis virus', *Virology Journal*, vol. 7, no. 1, pp. 241, available at <u>https://doi.org/10.1186/1743-422X-7-241</u>, accessed 25 February 2016.

4.4 Waterfowl parvoviruses

4.4.1 Background

Waterfowl parvoviruses are members of the Parvoviridae family. The International Committee on Taxonomy of Viruses (ICTV) currently classifies the waterfowl parvoviruses, Muscovy duck parvovirus (MDPV) and goose parvovirus (GPV), as a single species (namely, Anseriform dependoparvovirus 1), which is classified into the genus Dependoparvovirus in subfamily Parvovirinae due to similar genetic properties and evolutionary origins (ICTV 2014).

Variants of both GPV and MDPV have appeared in waterfowl and include the duck parvovirus (DPV), a recently identified variant of the GPV (Jordan et al. 2016; Li et al. 2018; Woźniakowski, Kozdruń & Samorek-Salamonowicz 2009; Zádori et al. 2006). A variant of the MDPV was also recently identified in Muscovy ducks in China and found to be a recombination between classical MDPV and GPV strains (Wang et al. 2017).

The GPV and MDPV are antigenically related (Mahardika et al. 2016; Yu et al. 2016) and possess approximately 85 per cent homology in overall protein sequences. Nonetheless, GPV and MDPV are distinct, and as a result, any antibody-driven cross protection between these two related viruses is largely limited (Yu et al. 2016).

Waterfowl parvoviruses are highly contagious with high mortalities in young birds, often causing between 70 and 100 per cent mortality in goslings and ducklings less than four weeks old (Zsak 2016). GPV causes Derzsy's disease, also known as goose hepatitis, goose plague or gosling plague, a disease characterised by watery diarrhoea, lethargy, anorexia, prostration, and weight loss in young Muscovy ducklings and goslings (Zsak 2016). MDVP affects only young Muscovy ducklings and mule ducks (Woolcock et al. 2000) and causes ascites, enteritis, myocarditis and hepatitis with high morbidity and mortality (Woźniakowski, Kozdruń & Samorek-Salamonowicz 2009). DPV causes similar clinical signs in mule ducks and Pekin ducks (Cherry Valley breed) (Chen et al. 2016). Young birds that survive infection and infected older birds show degenerative skeletal muscle myopathy and growth retardation characterised by the beak atrophy and dwarfism syndrome (Zsak 2016).

Vaccinating breeding flocks provides a high level of maternal antibody protection in the progeny until approximately two weeks of age (Zsak 2016).

Infection due to waterfowl parvoviruses is not OIE-listed nor is it nationally notifiable or subject to official control or eradication in Australia.

4.4.2 Technical information

Epidemiology

Susceptible species

Waterfowl parvovirus infects birds within the Anatidae family, namely geese, Muscovy ducks, swans (*Cygnus* spp), mule ducks and more recently Pekin ducks, especially those of the Cherry Valley breed (Li et al. 2018). GPV primarily infects geese and Muscovy ducks, and can be highly contagious in goslings under the age of four weeks (Yu et al. 2016). However, other breeds of ducks have shown susceptibility to variants of GPV. In 1989/1990 outbreaks, Muscovy ducks, mule ducks and Tsaiya ducks were susceptible to GPV (Chang et al. 2000; Liao et al. 1991). In 2014, a Malaysian study revealed that GPV was detected in Pekin ducks from Sarawak (Syamsiah et al. 2017). GPV antibodies

have been detected in the sera of wild geese (Hlinak et al. 1998). Since 2015, a number of DPV strains have been identified in Pekin and mule ducks in northern China (Chen et al. 2016; Li et al. 2018).

Geese are resistant to MDPV infection; however, both GPV and MDPV can cause severe disease in Muscovy ducks and several hybrid duck breeds (Yu et al. 2016).

Evidence suggests that wild geese are implicated in the epidemiology of the disease as a high prevalence of GPV antibodies were detected in the sera of wild geese (bean geese and white-fronted geese) in Germany (Hlinak et al. 1998). In addition, in 2004, epidemiological investigations of the outbreaks in the United Kingdom and Sweden led to a concern that some migratory wild geese species (including the Canada goose (*Branta canadensis*)) may have been the source of introduction of GPV (Gough et al. 2005; Jansson et al. 2007). Although over 50 taxa of waterbirds have been identified in Australia (Kingsford & Porter 2009), the susceptibility of Australian wild waterfowl species to waterfowl parvoviruses is not known. However, the Canada goose (*Branta canadensis*), suspected to be a source of GPV in the UK and Sweden and originally introduced into Australia as an exotic waterfowl species, have been occasionally sighted in the wild (DPIRD 2017).

Modes of transmission

Waterfowl parvoviruses are excreted in large amounts in the faeces of infected waterfowl and can subsequently spread rapidly to susceptible birds by both direct and indirect routes (Yu et al. 2016). Goslings and ducklings can be infected through vertical transmission via eggs laid by infected breeder geese and ducks (Chen et al. 2016; Zádori et al. 2006; Zsak 2016). Infected ducklings hatching from eggs where the virus was vertically transmitted, are often the source of the virus when outbreaks of GPV occur in countries free of the disease (Zsak 2016).

GPV may cause subclinical infection in older geese which pass on the virus to their progeny. Recovered birds are thought to act as carriers, and latency may be established with vertical (egg) transmission of the virus and potentially further horizontal transmission (Irvine & Holmes 2010; Palya 2010).

Horizontal infection is faecal-oral and occurs between flock-mates through the digestive tract via virus contaminated water and feed (Zádori et al. 2006). There is no reliable scientific information about transmission via the respiratory system or the conjunctiva.

Due to its resistance in the environment, parvovirus can persist in buildings and on poorly cleaned and disinfected surfaces. This can result in transmission between subsequent flocks (Palya 2010). Machinery and other farm equipment can also spread the disease as the virus is very difficult to remove using conventional cleaning agents.

Incubation period

In unprotected goslings infected with GPV in their first 10 days of life, clinical signs develop five to seven days post infection. The signs include inappetance, listlessness, anorexia, polydipsia, and diarrhoea. Deaths peak around 11 to 12 days post infection (Zádori et al. 2006). In susceptible neonatal birds, death may occur in two to five days (Irvine & Holmes 2010). Older geese and Muscovy ducks may develop subclinical infection (Irvine & Holmes 2010).

In three day old Muscovy ducklings experimentally infected with GPV, mortality reached 100 per cent within 10 days and the virus was distributed to many tissues within two days after inoculation by mouth (Takehara et al. 1998). GPV DNA was detected by PCR in the duodenum and rectum from day

3 to day 28 post inoculation. This suggests that birds may excrete virus in their faeces for a long period of time (Takehara et al. 1998).

Egg transmission will lead to embryo death during incubation or hatching in an infected state, depending on the level of yolk-sac antibody (Zádori et al. 2006). There are reports that latency may be established with vertical transmission of the virus (Irvine & Holmes 2010).

Distribution and prevalence

Currently, the majority of northern hemisphere countries with intensively farmed waterfowl have GPV (Zádori et al. 2006). It has been reported to cause disease in geese and Muscovy ducks in China, France, Germany, Hungary, North America, Poland, Thailand, Taiwan and Vietnam (Chang et al. 2000; Deemagarn & Tangdee 2015; Irvine & Holmes 2010; Jordan et al. 2016; Mahardika et al. 2016; Syamsiah et al. 2017; Yu et al. 2016)

It was suspected that imports of large numbers of goslings and goose eggs from Thailand might have been involved in a 1982 outbreak of GPV in Taiwan (Chang et al. 2000).

MDPV was first isolated from Muscovy ducks in the west of France in 1989. The United States of America also experienced an outbreak of MDPV in 1997 (Woolcock et al. 2000). The disease has been reported in Europe and Asia including China, France, Germany, Indonesia, Japan, Malaysia and Thailand (Li et al. 2018; Mahardika et al. 2016; Sirivan, Tantaswasdi & Chuenchai 1997; Takehara et al. 1994; Woolcock et al. 2000).

Australian Status

Waterfowl parvovirus has not been reported in Australia, there is no active surveillance, and waterfowl parvovirus is not nationally notifiable.

Agent properties

Parvoviruses can survive for long periods in the environment and show marked resistance to environmental and thermal stress, and pH extremes (Jordan et al. 2016; Ridpath & Mengeling 1988).

Parvoviruses, being non-enveloped and hydrophilic, have high resistance to disinfection. The GPV particles are resistant to various organic solvents and disinfectants (including chloroform, ether, fluorocarbon, 1:1000 formalin, 0.5 percent phenol, and sodium deoxycholate) and they cannot be inactivated by low pH (pH 3) or heat treatment typically used for disinfection applications (56°C for 30 minutes or 65°C for 30 minutes) (Schettler 1973; Takehara et al. 1994). The department is not aware of any inactivation studies specifically for GPV or MDPV.

4.4.3 Presence in duck meat

Transmission of GPV or MDPV by feeding infected duck meat or by feed contaminated with infected duck meat has not been reported. However, virus has been demonstrated in the organs and muscles of Muscovy ducks that were experimentally inoculated with attenuated GPV by mouth (Takehara et al. 1998). In this experiment, GPV DNA was detected by PCR at 2, 6, 10, 14 and 28 days post inoculation and viral titre in cardiac muscle reached 850 plaque forming units (PFU)/ml and 1316 PFU/ml at days 10 and 14 post inoculation respectively. GPV DNA was detected in the skeletal muscle up to 28 days post inoculation. The infective dose of GPV and MDPV has not been established.

4.4.4 Pathogenesis

Once the virus enters the gastrointestinal tract it replicates in the cells of the intestinal wall. From there the virus spreads through the body to other organs including the heart, liver and skeletal muscle (Zádori et al. 2006; Zsak 2016).

GPV genome has been detected by PCR in tissue samples from the respiratory tract, digestive tract, muscle (musculus longus colli and major adductor magnus), bone marrow and brain of Muscovy ducklings which were inoculated with the virus. GPV in these tissues was detected from day 4 to 10 post inoculation (Limn et al. 1996). A study to measure the distribution of GPV within tissues after oral infection found that virus was consistently detected in the blood, heart, liver, spleen, kidney, bursa of Fabricius, thymus and Harder's glands by eight hours post inoculation, and titres had significantly decreased at six and nine days (Yang et al. 2009).

In a later study in 2016, 40 liver and spleen tissue samples from ducklings retrieved from farms affected with GPV outbreaks in China were all positive for the virus (Yu et al. 2016).

MDPV was isolated in allantoic fluid after the inoculation of a pool of liver, heart and skeletal muscle tissue samples from ducks found dead during an outbreak in California (January 1998) into the chorioallantoic sacs of embryonating Muscovy duck eggs (Woolcock et al. 2000).

A study showed DPV DNA present in hatching eggs, duck embryos and newly hatched ducklings. Of the newly hatched ducklings, 58.33 per cent (21/36) were seropositive. Further, isolates were obtained from a 12 day old duck embryo and a newly hatched duckling. DPV infection did not reduce the fertilisation rate and hatchability (Chen et al. 2016).

4.4.5 Diagnosis

Clinical signs

Infection with GPV can result in 100 per cent mortality in its acute form and geese remain susceptible to GPV throughout their lives (FAO 2002; Zádori et al. 2006). However, clinical sensitivity to GPV in geese is usually less than 30 days of age, compared to 40 to 50 days of age in Muscovy ducklings infected with MDPV (Zádori et al. 2006). If birds are infected during the first week of life, high mortalities (up to 100 per cent) often occur, but once the animals are four to five weeks old it is likely infection will be subclinical (FAO 2002; Jordan et al. 2016; Zádori et al. 2006). In one outbreak in China, 5 per cent of total tissue samples and 39 per cent of total faecal swabs collected from ducklings with no clinical signs were positive for GPV (Yu et al. 2016). Clinical findings in MDPV infected ducklings resemble those of GPV-infected goslings (Mahardika et al. 2016).

During an outbreak of MDPV in California in October 1997, weakness, lateral recumbency and inability to walk were observed in Muscovy ducklings one to four weeks old. There was also increased mortality (10 to 50 per cent) and morbidity (30 to 80 per cent) in these ducklings (Woolcock et al. 2000).

There have been recent reports that GPV has also caused duck beak atrophy and dwarfism syndrome in ducklings (Palya 2010; Yu et al. 2016).

Pathology

Post-mortem findings include pale thigh and leg muscles, pale myocardium, fibrinous exudate on the liver capsule, pulmonary oedema and congestion, and ascites. Older survivors can have pathological

changes such as rounded hearts, and chronic passive congestion of the liver, and ascites (Mahardika et al. 2016; Poonia et al. 2006)

Testing

Clinical diagnosis is not sufficient to confirm disease and laboratory confirmation is necessary to diagnose GPV and MDPV (Xiao et al. 2017).

As none of the waterfowl parvoviruses is OIE-listed, there is no recommended diagnostic test. PCR and serology are the most commonly used laboratory tests (Palya 2010).

Immunoelectron microscopy, ELISA, immunofluorescence antibody testing and agar gel precipitin have been used for serology to detect GPV or MDPV antibodies (Irvine & Holmes 2010; Woolcock et al. 2000). In one study, sera from older ducklings (35 day old and 63 day old) were positive on ELISA, but no antibodies were detected in 10 day old ducklings. In the same study three of ten sera taken from 21 day old ducklings had low titres of antibody which may indicate the start of a serological response (Woolcock et al. 2000). Antibodies may persist for years in older Muscovy ducks that have subclinical infection (Irvine & Holmes 2010).

There are no tests available that can differentiate between GPV and MDPV antibodies because the two viruses share many antigenic similarities (Woolcock et al. 2000).

Electron microscopy, PCR or RT-PCR can detect the virus (Yu et al. 2016).

4.4.6 Treatment

There is no effective treatment for waterfowl parvovirus infection.

4.4.7 Control

Control for GPV infection relies on effective flock biosecurity, hygiene and eliminating carrier birds (Irvine & Holmes 2010).

Many outbreaks have been attributed to birds hatched from eggs contaminated with the virus from GPV infected breeding flocks. Therefore, eggs should be imported only from countries that can guarantee breeder flock freedom from GPV. Geese that have survived an outbreak should not be used for breeding purposes as recovered birds may act as carriers (Irvine & Holmes 2010; Zsak 2016).

Live and inactivated vaccines are used in some countries to protect goslings during the first few weeks of life. The preferred vaccines are based on live attenuated viruses (Jordan et al. 2016). Vaccine used in breeder geese results in the transfer of immunity to newly hatched goslings for two to three weeks (FAO 2002).

There are conflicting reports of whether GPV vaccine is effective against both GPV and MDPV. Whilst some of the literature suggests that GPV strain vaccines are ineffective against MDPV (Zádori et al. 2006); others report that 'vaccinating Muscovy ducklings with strains of GPV provided them with a high level of protection against a challenge with MDPV' (Woolcock et al. 2000).

Studies on a recently developed vaccine based on a live attenuated Muscovy duck parvovirus, produced by Merial, showed that vaccination with a live attenuated MDPV strain provided immunity to Muscovy ducks against GPV as well as MDPV (Maurin-Bernaud et al. 2014). This vaccine is now registered for use within the European Union (EMA 2015).

4.4.8 Current biosecurity measures

Australia currently requires that for foie gras for human consumption from France, during the 12 to 21 day stuffing period prior to slaughter, certification that there has been no outbreak of Derzsy's disease on the properties of origin of the ducks and/or geese from which the livers were derived.

With regards to importing fertile duck eggs, Australia requires:

- Either the country of origin is free from Derszy's disease or serological testing of the parent flock within 21 days before the first day of collection of eggs for export to Australia has been performed, with negative results;
- The source flock is free from clinical signs of GPV and MDPV (Muscovies only) for the 90 day period prior to collection of the eggs for export to Australia and has not come into contact with any birds showing evidence of these diseases;
- Any clinical evidence of disease in the source flock in the 21 days since the last day of collection of the eggs exported to Australia has been investigated and the results indicate disease caused by GPV or MDPV has not occurred; and
- Serology of imported birds in quarantine for goose parvovirus has been performed 6 weeks post hatching, with negative results.

4.4.9 Conclusion

- Waterfowl parvoviruses have been reported to cause disease in geese and ducks in Thailand.
- Waterfowl parvoviruses have not been reported in Australia.
- Transmission of GPV or MDPV by feeding infected duck meat or by feed contaminated with infected duck meat has not been reported.
- Virus has been demonstrated in the organs and muscles of Muscovy ducks that were experimentally inoculated with attenuated GPV by mouth.
- The virus is markedly resistant to environmental and thermal inactivation and may survive for long periods of time in infected duck carcasses. It is likely that the virus will resist cooking at the time and temperature defined for this review.
- GPV and MDPV are not OIE-listed disease agents and there are no recommendations in the OIE Code on disease testing and measures for safe trade.
- Waterfowl parvovirus is not nationally notifiable.
- Vaccines are available to manage the disease.

Based on the above information cooking as defined in the scope at section 1.3 of this document (70°C for 8.2 minutes) is likely to reduce the viral load but not fully inactivate the waterfowl parvoviruses that may potentially be present in meat of ducks. Therefore the department concluded that further risk assessment of the waterfowl parvoviruses was required.

4.4.10 Risk Assessment

Entry assessment

- Both MDPV and GPV have been reported in Thailand. Whilst there are no reports of MDPV affecting species other than Muscovy ducks and their hybrid breeds, GPV has been reported in Pekin ducks.
- Although GPV causes severe disease in young ducklings, older ducklings greater than four to five weeks of age may not show clinical signs but may be shedding the virus. Older birds may also be subclinically infected. Furthermore recovered birds may act as carriers and be a source of infection. Clinical signs will usually be observed in young ducklings and as the disease is likely to cause severe morbidity and mortality it would almost certainly be identified on affected farms and limit the ability to supply meat.
- Infection of slaughter age ducks (greater than seven weeks) are likely to be subclinical and may not be identified on ante-mortem inspection. Gross lesions may not be readily visible on post-mortem inspection of these ducks.
- The virus spreads through many organs including skeletal muscle. The department is not aware
 of any inactivation studies specifically for GPV or MDPV and it has been reported that parvovirus
 is resistant to thermal inactivation (56°C for 30 minutes or 65°C for 30 minutes) and cannot be
 inactivated by low pH. Therefore, it is likely that while the viral load would be reduced, some
 virus would resist cooking at the time and temperature defined for this review, 70°C for 8.2
 minutes.

Conclusion: Considering the above factors, the likelihood of entry in imported cooked duck meat from Thailand was assessed to be **moderate.**

Exposure assessment

- The virus is markedly resistant and could remain viable for a long period of time in the environment.
- Vertical transmission through infected eggs is the most commonly reported source of outbreaks in disease-free countries. There have been no reports of oral transmission through infected meat; however, transmission by mouth cannot be discounted given that oral transmission of GPV has been demonstrated experimentally. The infective dose of GPV and MDPV has not been described.
- The impact of the viruses in Australian wild waterfowl is not known. Evidence suggests that some Northern Hemisphere species of wild geese were implicated in the epidemiology of the disease.
- As only cooked products consisting of muscle meat and overlying fat and skin would be imported, most imported product would move from the retailer/distributor to household consumers or to the food service industry. Product not consumed would be disposed of, which may result in exposure to wild birds and non-avian species. More likely in products sold directly to households, backyard (low biosecurity) poultry could also be exposed to waste. Limiting the scope to muscle meat and overlying fat and skin that has been cooked reduces the type and volume of waste that may be created. Bone, trimmings and offal would not be imported, reducing the risk of exposure to waste.

 A possible exposure pathway for some waste to Australian commercial ducks and geese is via feed for ducks containing meat meal made from waste from imported cooked duck meat. It is unknown whether rendering, especially the continuous low temperature type (70 to 95°C) will totally inactivate any parvovirus if present. However the Australian Standard for Hygienic Rendering of Animal Products (AS 5008:2001) requires processing plants to verify that their process inactivates spore forming bacteria (Primary Industries Standing Committee & Primary Industries Ministerial Council 2007). In poultry diets, meat and bone meal is typically limited to less than five per cent of the diet because of the high calcium, phosphorus, and lysine content of the meal. The possibility of exposure via this pathway is further reduced by the restriction on the products that can be imported as outlined in the scope.

Conclusion: Based on this information, the likelihood of exposure of domestically kept geese and ducks to waterfowl parvovirus infected duck meat was assessed to be **low**.

Estimation of likelihood of entry and exposure

The likelihood of entry of this agent to Australia and the corresponding likelihood of exposure to the Australian duck and goose population was estimated using the matrix of rules described in Table 1. As the assessment of the likelihood of entry is moderate and the likelihood of exposure is low the estimation of the likelihood of entry and exposure of a waterfowl parvovirus was assessed to be **low**.

Likelihood of establishment and/or spread associated with the outbreak scenario

The most likely outbreak scenario following exposure to a waterfowl parvovirus was considered to be establishment in susceptible wild and low biosecurity geese and duck populations with vertical and horizontal spread to more than one state or territory.

The following factors were considered relevant to an estimate of the likelihood of establishment and/or spread associated with exposure of susceptible duck and geese to a waterfowl parvovirus:

- Vertical transmission through infected eggs is the most commonly reported source of an outbreak. GPV may cause subclinical infection in older geese and pass on the virus to their progeny. Recovered birds may be carriers.
- Waterfowl parvoviruses are excreted in large amounts in the faeces of infected waterfowl and can subsequently spread rapidly to susceptible birds by both direct and indirect routes.
- Due to its resistance in the environment, parvovirus can persist in buildings and on poorly cleaned and disinfected surfaces which results in transmission between subsequent flocks.
- The role of wild geese and ducks in the spread of the disease is undefined. Evidence suggests that two species of wild geese in the northern hemisphere may be implicated in the epidemiology of the disease. Although over 50 taxa of waterbirds have been identified in Australia (Kingsford & Porter 2009), the susceptibility of Australian wild waterfowl species to waterfowl parvoviruses is not known. However, the Canada goose (*Branta canadensis*), suspected to be a source of GPV in the UK and Sweden and originally introduced into Australia as an exotic waterfowl species, have been occasionally sighted in the wild (DPIRD 2017).
- Waterfowl parvoviruses are highly contagious and fatal in goslings, Muscovy ducklings, and Pekin ducklings, often causing between 70 and 100 per cent mortality in goslings less than four weeks old.

Conclusion: Considering the above factors, the likelihood of establishment and spread of a waterfowl parvovirus through the Australian duck or goose population was assessed to be **moderate**.

Determination of the effects resulting from the outbreak scenario

Direct effects

The effect on the life or health (including production effects) of susceptible animals

- GPV is highly contagious and fatal in goslings and Muscovy ducklings, often causing between 70 and 100 per cent mortality in goslings less than four weeks old. MDPV causes similar clinical signs but causes disease only in Muscovy ducks and their hybrids.
- If the birds are four to five weeks old or older it is likely the infection will be subclinical.

The effect on the living environment, including life and health of wildlife, and any effects on the non-living environment

• There are no reports of DPV in wild ducks. Wild geese may be infected with GPV, however, there is no evidence of adverse effects on the environment.

Indirect effects

The effect on new or modified eradication, control, monitoring or surveillance and compensation strategies or programs

- GPV and MDPV are not notifiable in any Australian jurisdiction and there are no control, monitoring or surveillance programs in place.
- An outbreak would likely provoke similar control and eradication measures to those that have been introduced in other affected countries. These include active surveillance, disinfection of infected flocks and elimination of carrier birds. A vaccination program may be introduced.

The effect on domestic trade or industry, including changes in consumer demand and effects on other industries supplying inputs to, or using outputs from, directly affected industries

- The Australian duck farming industry is dominated by two vertically integrated producers, one in New South Wales and the other in Victoria. Both breed and grow ducks under intensive and biosecure conditions. There are also several smaller duck farmers, especially in Victoria and Western Australia, as well as backyard ducks scattered around the country. There are no significant producers of duck eggs for commercial purposes (PoultryHub 2018). Because of the structure of the major producers, it is unlikely they would be affected by a waterfowl parvovirus.
- Pekin and Muscovy ducks are among the major breeds used for meat production, and the Pekin duck is the predominant breed used for meat production in Australia (Stein 2012).
- Commercial customers would be unlikely to be affected by shortages of duck meat or eggs due to direct losses of stock.
- There would be no effect on consumer demand.

The effect on international trade, including loss of and restriction of markets, meeting new technical requirements to enter or maintain markets, and changes in international consumer demand

• There are unlikely to be any impacts on international trade. Only five per cent of processed duck meat produced in Australia is exported (PoultryHub 2018).

The effect on the environment, including biodiversity, endangered species and the integrity of ecosystems

• As there has not been any discernible effects on wild duck or goose species in the northern hemisphere, it is expected that there would be no discernible effects on the environment.

The effect on communities, including reduced rural and regional economic viability and loss of social amenity, and any 'side effects' of control measures

 In young ducks and goslings, GPV and MDPV may cause significant mortalities and morbidities, and it is likely that breeding flocks would be destroyed following an outbreak due to their carrier state. Loss of stock may affect the livelihood and income of those directly affected. There are only two major duck breeders in Australia employing a total of around 300 people. Therefore, affected areas may experience very minor issues while movement restrictions are in place.

Conclusion for overall direct and indirect effects: Based on the geographic level and magnitude of effects, the overall effect of establishment and/or spread for the outbreak scenario was estimated to be **low.** The effect is likely to be recognised within affected zones and significant to directly affected parties. It is not likely that the effect will be recognised at the national level.

Consequence assessment

The estimate of the overall effect associated with the outbreak scenario (low) was combined with the likelihood of establishment and/or spread for the scenario (moderate) using Table 3 to obtain an estimation of likely consequences (low).

Risk estimation

Risk estimation is the integration of likelihood of entry and exposure, and likely consequences of establishment and/or spread to derive the risk associated with entry, exposure, establishment and/or spread of waterfowl parvoviruses being introduced by cooked duck meat imported into Australia from Thailand.

Using Table 4, the likelihood of entry and exposure (**low**) was combined with the likely consequences of establishment and/or spread (**low**), resulting in a risk estimation of **very low**.

Therefore, as the unrestricted risk estimate achieves Australia's ALOP, the department concluded that specific risk management was not necessary for waterfowl parvoviruses. The detailed requirements set out in section 5 which include, but are not limited to, restrictions to the type of meat, anti and post mortem veterinary inspection, investigation of source flock disease to rule out waterfowl parvoviruses, and cooking to reduce viral load are considered by the department to be acceptable risk management measures for waterfowl parvoviruses.

References

Chang, PC, Shien, JH, Wang, MS & Shieh, HK 2000, 'Phylogenetic analysis of parvoviruses isolated in Taiwan from ducks and geese', *Avian Pathology*, vol. 29, no. 1, pp. 45-9, available at https://doi.org/10.1080/03079450094270, accessed 3 April 2018.

Chen, H, Tang, Y, Dou, Y, Zheng, X & Diao, Y 2016, 'Evidence for vertical transmission of novel duckorigin goose parvovirus-related parvovirus', *Transboundary and Emerging Diseases*, vol. 63, no. 3, pp. 243-7, available at <u>https://doi.org/10.1111/tbed.12487</u>, accessed 31 July 2018. Deemagarn, T & Tangdee, S 2015, 'Nucleotide sequences of goose parvovirus isolates from Thailand and the comparison with vaccine strains', *Proceedings of 53rd Kasetsart University annual conference, Bangkok, Thailand*, Kasetsart University, Bangkok.

DPIRD 2017, 'Canada goose: animal pest alert', Department of Primary Industries and Regional Development, Perth, available at <u>https://www.agric.wa.gov.au/birds/canada-goose-animal-pest-alert</u> accessed 12 December 2018.

EMA 2015, 'Parvoduk: Muscovy duck parvovirus (live attenuated)', *European Public Assessment Report (EPAR)*, European Medicines Agency, London, available at <u>http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/veterinary/medicines/002740/vet</u> <u>med_000290.jsp&mid=WC0b01ac058008d7a8</u> accessed 3 July 2018.

FAO 2002, 'Goose diseases', in Buckland, R & Guy, G (eds), *Goose production, FAO Animal Production and Health Paper - 154*, Food and Agriculture Organization of the United Nations, Rome, available at http://www.fao.org/docrep/005/Y4359E/y4359e0g.htm accessed 7 March 2018.

Gough, D, Ceeraz, V, Cox, B, Palya, V & Mato, T 2005, 'Isolation and identification of goose parvovirus in the UK', *Veterinary Record*, vol. 156, no. 13, p. 424, available at http://dx.doi.org/10.1136/vr.156.13.424, accessed 1 May 2018.

Hlinak, A, Müller, T, Kramer, M, Mühle, RU, Liebherr, H & Ziedler, K 1998, 'Serological survey of viral pathogens in bean and white-fronted geese from Germany', *Journal of Wildlife Diseases*, vol. 34, no. 3, pp. 479-86, available at <u>https://doi.org/10.7589/0090-3558-34.3.479</u>, accessed 1 May 2018.

ICTV 2014, 'ICTV Taxonomy history: Anseriform dependoparvovirus 1', International Committee on Taxonomy of Viruses, Washington, available at https://talk.ictvonline.org/taxonomy/p/taxonomy-history?taxnode_id=20184260 accessed 9 January 2019.

Irvine, R & Holmes, P 2010, 'Diagnosis and control of goose parvovirus', *In Practice*, vol. 32, no. 8, pp. 382-6, available at <u>http://dx.doi.org/10.1136/inpract.32.8.382</u>, accessed 1 May 2018.

Jansson, DS, Feinstein, R, Kardi, V, Mató, T & Palya, V 2007, 'Epidemiologic investigation of an outbreak of goose parvovirus infection in Sweden', *Avian Diseases*, vol. 51, no. 2, pp. 609-13, available at <u>https://doi.org/10.1637/0005-2086(2007)51[609:EIOAOO]2.0.CO;2</u>, accessed 3 July 2018.

Jordan, I, John, K, Höwing, K, Lohr, V, Penzes, Z, Gubucz-Sombor, E, Fu, Y, Gao, P, Harder, T, Zádori, Z & Sandig, V 2016, 'Continuous cell lines from the Muscovy duck as potential replacement for primary cells in the production of avian vaccines', *Avian Pathology*, vol. 45, no. 2, pp. 137-55, available at https://doi.org/10.1080/03079457.2016.1138280, accessed 21 March 2018.

Kingsford, RT & Porter, JL 2009, 'Monitoring waterbird populations with aerial surveys – what have we learnt?', *Wildlife Research*, vol. 36, no. 1, pp. 29-40, available at <u>https://doi.org/10.1071/WR08034</u>, accessed 12 December 2018.

Li, P, Lin, S, Zhang, R, Chen, J, Sun, D, Lan, J, Song, S, Xie, Z & Jiang, S 2018, 'Isolation and characterization of novel goose parvovirus-related virus reveal the evolution of waterfowl parvovirus', *Transboundary and Emerging Diseases*, vol. 65, no. 2, pp. e284-e295, available at https://doi.org/10.1111/tbed.12751, accessed 31 July 2018.

Liao, YK, Lu, YS, Lin, YL, Lee, SH & Chiu, SY 1991, *The outbreak and control of duck viral disease in Taiwan, 1989-90*, Food and Fertilizer Technology Center: extension bulletins, available at http://www.agnet.org/library/eb/338/.

Limn, C, Yamada, T, Nakamura, M & Takehara, K 1996, 'Detection of goose parvovirus genome by polymerase chain reaction: distribution of goose parvovirus in Muscovy ducklings', *Virus Research*, vol. 42, no. 1, pp. 167-72, available at <u>https://doi.org/10.1016/0168-1702(95)01310-5</u>, accessed 24 April 2018.

Mahardika, GN, Putra, MBAPA, Dewi, NPS, Dewi, NMRK & Winaya, IBO 2016, 'Muscovy duck parvovirus infection with epicarditis in Bali, Indonesia', *Journal of Veterinary Science and Technology*, vol. 7, pp. 328, available at doi: 10.4172/2157-7579.1000328, accessed 21 March 2018.

Maurin-Bernaud, L, Goutebroze, S, Merdy, O, Chanay, A, Cozette, V & Le Gros, F-X 2014, 'Efficacy of a new attenuated duck parvovirosis vaccine in Muscovy ducks', *Veterinary Record*, vol. 175, no. 11, pp. 281, available at http://dx.doi.org/10.1136/vr.102584, accessed 3 May 2018.

Palya, V 2010, 'Major viral diseases of waterfowl and their control', The British Veterinary Poultry Association, Driffield, East Yorkshire, UK, available at <u>http://www.bvpa.org.uk/library.php</u> accessed 03/21/2018.

Poonia, B, Dunn, PA, Lu, H, Jarosinski, KW & Schat, KA 2006, 'Isolation and molecular characterization of a new Muscovy duck parvovirus from Muscovy ducks in the USA', *Avian Pathology*, vol. 35, no. 6, pp. 435-41, available at <u>https://doi.org/10.1080/03079450601009563</u>, accessed 21 March 2018.

PoultryHub 2018, 'Duck', PoultryHub Australia, Armidale, NSW, available at <u>http://www.poultryhub.org/species/commercial-poultry/duck/</u> accessed 12 December 2018.

Primary Industries Standing Committee & Primary Industries Ministerial Council 2007, *Australian standard for the hygienic rendering of animal products*, PISC Report no. 87, AS 5008:2007, CSIRO Publishing, Victoria.

Ridpath, JF & Mengeling, WL 1988, 'Parvoviruses', in Darai, G (ed) *Virus diseases in laboratory and captive animals*, Martinus Nijhoff Publishing, Boston, pp. 293-320.

Schettler, CH 1973, 'Virus hepatitis of geese 3. Properties of the causal agent', *Avian Pathology*, vol. 2, no. 3, pp. 179-93, available at <u>https://doi.org/10.1080/03079457309353795</u>, accessed 3 May 2018.

Sirivan, P, Tantaswasdi, U & Chuenchai, S 1997, 'Isolation and identification of duck hepatitis virus in ducklings', *Abstracts from The Journal of Thai Veterinary Medical Association: 1997 issues*, available at <u>http://tvma.tripod.com/news/abstracts97.html</u> accessed 7 November 2018.

Stein, B 2012, 'Introduction to commercial duck farming', *Factsheet 1235*, NSW Department of Primary Industries, Orange, NSW, available at <u>https://www.dpi.nsw.gov.au/animals-and-livestock/poultry-and-birds/species/duck-raising/commercial-duck-farming</u> accessed 12 December 2018.

Syamsiah, AS, Leow, BL, Ong, GH, Muhammad, RS & Basirah, MA 2017, 'Nucleotide sequence and phylogenetic analysis of goose parvovirus detected from Pekin duck', *Malaysian Journal of Veterinary Research*, vol. 8, no. 1, pp. 131-7

Takehara, K, Hyakutake, K, Imamura, T, Mutoh, K & Yoshimura, M 1994, 'Isolation, identification, and plaque titration of parvovirus from Muscovy ducks in Japan', *Avian Diseases*, vol. 38, no. 4, pp. 810-5

Takehara, K, Saitoh, M, Kiyono, M & Nakamura, M 1998, 'Distribution of attenuated goose parvoviruses in Muscovy ducklings', *Journal of Veterinary Medical Science*, vol. 60, no. 3, pp. 341-4, available at https://doi.org/10.1292/jvms.60.341, accessed 9 May 2018.

Wang, S, Cheng, X, Yu, B, Xiao, S, Chen, S, Zhu, X, Yu, F, Lin, S, Chen, S, Lin, F, Wu, N, Wang, J, Huang, M & Zheng, M 2017, 'A simple, polymerase chain reaction and restriction fragment length polymorphism-aided diagnosis method for short beak and dwarfism syndrome in ducklings', *Infection, Genetics and Evolution*, vol. 53, pp. 85-8, available at https://doi.org/10.1016/j.meegid.2017.05.014, accessed 31 July 2018.

Woolcock, PR, Shivaprasad, HL, McFarland, MD, Jestin, V, Zwingelstein, F, Arnauld, C, Pedersen, JC & Senne, DA 2000, 'Evidence of Muscovy duck parvovirus in Muscovy ducklings in California', *The Veterinary Record*, vol. 146, no. 3, pp. 68-72, available at http://veterinaryrecord.bvapublications.com/cgi/content/abstract/146/3/68

Woźniakowski, G, Kozdruń, W & Samorek-Salamonowicz, E 2009, 'Genetic variance of Derzsy's disease strains isolated in Poland', *Journal of Molecular and Genetic Medicine*, vol. 3, no. 2, pp. 210-6, available at <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2805842/</u>, accessed 21 March 2018.

Xiao, S, Chen, S, Cheng, X, Lin, F, Wang, S, Zhu, X, Yu, B, Huang, M, Wang, J, Wu, N, Zheng, M, Chen, S & Yu, F 2017, 'The newly emerging duck-origin goose parvovirus in China exhibits a wide range of pathogenicity to main domesticated waterfowl', *Veterinary Microbiology*, vol. 203, pp. 252-6, available at https://doi.org/10.1016/j.vetmic.2017.03.012, accessed 1 August 2018.

Yang, J-L, Cheng, A-C, Wang, M-S, Pan, K-C, Li, M, Guo, Y-F, Li, C-F, Zhu, D-K & Chen, X-Y 2009, 'Development of a fluorescent quantitative real-time polymerase chain reaction assay for the detection of goose parvovirus in vivo', *Virology Journal*, vol. 6, no. 1, pp. 142, available at <u>https://doi.org/10.1186/1743-422X-6-142</u>, accessed 24 April 2018. Yu, K, Ma, X, Sheng, Z, Qi, L, Liu, C, Wang, D, Huang, B, Li, F & Song, M 2016, 'Identification of gooseorigin parvovirus as a cause of newly emerging beak atrophy and dwarfism syndrome in ducklings', *Journal of Clinical Microbiology*, vol. 54, no. 8, pp. 1999-2007, available at <u>https://doi.org/10.1128/JCM.03244-15</u>, accessed 21 March 2018.

Zádori, Z, Szelei, J, Kiss, I & Tijssen, P 2006, 'Waterfowl parvoviruses', in Kerr, JR, Cotmore, SF, Bloom, ME, Linden, RM & Parrish, CR (eds), *Parvovirus*, Hodder Arnold, London, pp. 447-56.

Zsak, L 2016, 'Overview of goose parvovirus infection', *MSD veterinary manual [online]*, Merck & Co., Inc., Kenilworth, New Jersey, USA, available at <u>https://www.msdvetmanual.com/poultry/goose-parvovirus-infection/overview-of-goose-parvovirus-infection</u> accessed 3 July 2018.

5 Requirements for the importation of cooked duck meat from Thailand

5.1 Eligibility

Importation under these conditions is restricted to cooked duck meat from Thailand only.

5.2 Documentation

A written application to import cooked duck meat must be lodged with the department before any import can occur.

Each consignment must be accompanied by:

- 1) a valid import permit
- 2) an official veterinary health certificate in accordance with 'Model veterinary certificates for international trade in live animals, hatching eggs and products of animal origin' as described in Chapter 5.10 of the Code.

The veterinary certificate must provide details of:

- the packaging of the cooked duck meat product including details of the labelling
- the addresses and Veterinary Authority of Thailand approval numbers of establishments at which the ducks from which the meat was derived were slaughtered
- the facility at which the cooked duck meat product was prepared
- the establishment at which the cooked duck meat product was stored before export
- the names and addresses of the exporter and the consignee.

An Official Government Veterinarian means a veterinarian authorised by the Veterinary Authority of Thailand to perform certain official tasks associated with animal health and/or public health, inspections of commodities, and when appropriate, to certify in conformity with the Certification Procedures of Chapter 5.2 of the Code.

Any inadequacies in certification may result in the consignment being returned to the country of origin at the importer's expense or the destruction of the duck meat without compensation.

5.3 Certification

The certificate must include the name and stamp of the Official Government Veterinarian and contain the following declarations:

 A statement that all of the establishments where the ducks from which the meat was derived were slaughtered, processed and stored have current approval from the Veterinary Authority of Thailand and are operating in accordance with the laws of Thailand.

- 2) A statement that the ducks were slaughtered and processed in accordance with the regulatory requirements of Thailand for items intended for human consumption.
- 3) A statement that the ducks from which the meat was derived passed ante- and post-mortem veterinary inspection under official veterinary supervision, and the meat is considered fit for human consumption.
- 5) A statement that the duck meat is sourced from domestic ducks (*Anas platyrhynchos domesticus* and *Cairina moschata*) only.
- 6) A statement that the duck meat is limited to muscle meat and overlying skin and fat.
- 7) A statement that the duck meat has been cooked in a commercial process using moist heat to a minimum core temperature of 70°C for at least 8.2 minutes or equivalent according to the time/temperatures specified in Annex 1 and records confirm that these time/temperature parameters were met.
- 8) A statement that the temperature recording equipment was confirmed to be in good working order prior to each day's production and was independently calibrated at least once every 12 months.
- 9) A statement that all establishments that handled, prepared, processed or stored the cooked duck meat destined for Australia have approved and documented internal processes to keep such meat isolated from any meat and meat products that are not eligible for export to Australia and to ensure the integrity and health status of the product is maintained.
- 10) A statement that the cooked duck meat was packed for transport to Australia in a clean container and sealed with an official government seal.

5.4 Annex 1

Heat treatment considered equivalent to a moist heat treatment at a core temperature of least 70 °C for not less than 8.2 minutes includes the following time/temperature combinations. The heat treatment must be applied as a minimum core temperature and minimum time at that core temperature using moist heat in a commercial heating process:

- 1) 71 °C for not less than 6.6 minutes
- 2) 72 °C for not less than 5.2 minutes
- 3) 73 °C for not less than 4.2 minutes
- 4) 74 °C for not less than 3.3 minutes
- 5) 75 °C for not less than 2.6 minutes
- 6) 76 °C for not less than 2.1 minutes
- 7) 77 °C for not less than 1.7 minutes
- 8) 78 °C for not less than 1.3 minutes
- 9) 79 °C for not less than 1.1 minutes
- 10) 80 °C for not less than 1 minute

6 Review of processes

6.1 Review of conditions

The department reserves the right to review the import conditions after the first year of trade or when there is reason to believe that the disease or sanitary status of Thailand has changed or if there is evidence of new or emerging diseases. The department may also review the import conditions if there is any change to the nature or understanding of a disease, agent or exposure pathways.

7 Meeting Australia's food standards

Imported food for human consumption must satisfy Australia's food standards. Australian law requires that all food, including imported food, meets the standards set out in the Australia New Zealand Food Standards Code. FSANZ is responsible for developing and maintaining the Code, including Standard 1.4.2, maximum residue limits (MRLs), available on the <u>ComLaw</u> website. The standards apply to all food in Australia, irrespective of whether it is grown domestically or imported.

If a specific chemical is used on imported foods to control pests and diseases, then any resulting residues must not exceed the specific MRLs in Standard 1.4.2 of the Australia New Zealand Food Standards Code for that food.

If there is no MRL listed in the Australia New Zealand Food Standards Code for a specific food (or a composite, processed food), then there must be no detectable residues in that specific food.

Where an exporting country uses a chemical for which there is no current listed Australian MRL, there are mechanisms to consider establishing an Australian MRL by harmonising with an MRL established by the Codex Alimentarius Commission (Codex) or by a regulatory authority in a recognised jurisdiction. The mechanisms include applications, submissions or consideration as part of a FSANZ proposal to vary the Australia New Zealand Food Standards Code. The application process, including the explanation of establishment of MRLs in Australia, is described at the <u>Food Standards</u> <u>Australia New Zealand</u> website.