

### T301-e Cottonseed for Food or Feed

Pest: *Fusarium oxysporum* f. sp. *vasinfectum* strains VCG 01111 and VCG 01112

Treatment: T301-e—MB at NAP—tarpaulin

T301-e has been added to this manual through the immediate process described in paragraph (b) of § 305.3. This treatment may be applied immediately, but is subject to change or removal based on public comment..

| Temperature    | Dosage Rate (lb/1,000 ft <sup>3</sup> ) | Minimum Concentration Readings (ounces) At: |       |       |        |        |        |        |
|----------------|---|---|-------|-------|--------|--------|--------|--------|
|                |   | 2 hrs                                       | 4 hrs | 6 hrs | 12 hrs | 24 hrs | 36 hrs | 48 hrs |
| 40 °F or above | 7 lbs                                   | 60  | 50    | 50    | 35     | 20     | —      | —      |
|                | 5 lbs                                   | 20  | —     | 20    | 20     | 15     | 15     | 15     |

### T302—Grains and Seeds Not Intended for Propagation



If grain and seeds are for propagation, use appropriate treatment in T203 schedules

#### T302-g-1 Acorns not intended for propagation

Two alternative treatments

Pest: *Cydia splendana* (nut fruit tortrix) and *Curculio* spp. (weevils)

Treatment: T302-g-1—MB at NAP—tarpaulin, chamber, or van container

| Temperature | Dosage Rate (lb/1,000 ft <sup>3</sup> ) | Minimum Concentration Readings (ounces) At: |       |       |       |       |       |
|-------------|---|---|-------|-------|-------|-------|-------|
|             |   | 0.5 hr                                      | 2 hrs | 3 hrs | 4 hrs | 5 hrs | 6 hrs |
| 90-95 °F    | 4 lbs                                   | 58  | 32    | 34    | —     | —     | —     |
| 80-89 °F    | 4 lbs                                   | 58  | 32    | —     | 34    | —     | —     |
| 70-79 °F    | 5 lbs                                   | 72  | 40    | —     | 42    | —     | —     |
| 60-69 °F    | 5 lbs                                   | 72  | 40    | —     | —     | 40    | —     |
| 50-59 °F    | 6 lbs                                   | 85  | 48    | —     | —     | 50    | —     |
| 40-49 °F    | 6 lbs                                   | 85  | 48    | —     | —     | —     | 48    |