

The relationships between land management practices and soil condition and the quality of ecosystem services delivered from agricultural land in Australia

STEVEN CORK (PROJECT LEADER)

LAURA EADIE

PAULINE MELE

RICHARD PRICE

DON YULE

September 2012

RESEARCH WITHOUT BOUNDS



Kiri-ganai research



About Kiri-ganai research:

Kiri-ganai Research Pty Ltd is a Canberra based company that undertakes consultancy and analytical studies concerned with environmental policy, industry performance, natural resource management and sustainable agriculture. Our strength is in turning knowledge gained from public policy, markets, business operations, science, and research into ideas, options, strategies and response plans for industries, governments, communities and businesses.

Kiri-ganai Research Pty Ltd
GPO Box 103 CANBERRA ACT 2601 AUSTRALIA
ph: +62 2 62956300 fax: +61 2 62327727
www.kiri-ganai.com.au

Funding

This project was funded by the Australian Government's Caring for our Country initiative.

Project team

This project was managed by Kiri-ganai Research Pty Ltd. The main writing team comprised Steven Cork (EcoInsights), Pauline Mele (Victorian Department of Primary Industries), Laura Eadie (Centre for Policy Development), Don Yule (CTF Solutions) and Richard Price (Kiri-ganai Research). This team was guided by four expert advisers: Anna Roberts, Neil Byron, Geoff Gorrie and Barry White.

Acknowledgements

The project team gratefully acknowledges the contribution made to the project by members of the Australian Government Land and Coasts Division, and in particular Science Adviser, Dr Michele Barson.

Disclaimer

Considerable care has been taken to ensure that the information contained in this report is reliable and that the conclusions reflect considerable professional judgment. Kiri-ganai Research Pty Ltd, however, does not guarantee that the report is without flaw or is wholly appropriate for all purposes and, therefore, disclaims all liability for any loss or other consequence which may arise from reliance on any information contained herein.

Contents

Executive summary	vii
Questions addressed.....	vii
Key conclusions.....	vii
Benefits and beneficiaries from better soil management	viii
1. Project rationale and approach	1
1.1 Rationale	1
1.2 Approach	1
2. Soils: the essential asset.....	4
2.1 Soils, life and human interaction	4
2.2 Living soils and determinants of soil condition	4
2.3 Soils and systems.....	5
3. Linking management practices, soil quality and ecosystem services	7
3.1 The concept of ecosystem services.....	7
3.2 Ecosystem services and management practice.....	8
4. Soil Carbon	9
4.1 Nature of the issues.....	9
4.2 Impacts of agriculture and measures that could build Soil Organic Carbon....	10
4.3 Evidence of the efficacy of practices to increase soil organic carbon	16
5. Soil pH.....	19
5.1 Nature of the issues.....	19
5.2 Impacts of agriculture and measures that could arrest soil acidification	20
5.3 Evidence of the efficacy of practices to increase soil pH	26
5.4 Concluding remarks.....	28
6. Wind erosion	30
6.1 Nature of the issues.....	30
6.2 Land management practices in relation to wind erosion	31

6.3 Evidence of the effectiveness of management practices for reducing wind erosion	33
7. Water erosion	36
7.1 Nature of the issues	36
7.2 Land management practices in relation to water erosion	38
7.3 Evidence of the effectiveness of management practices for reducing water erosion	40
8. Ecosystem services and resilience of soils	46
8.1 The concept of ecosystem services	46
8.2 Relating soil ecosystem processes to services and benefits	48
8.3 How better management for soil carbon, pH and erosion might affect ecosystem services	54
8.4 Resilience of soils and associated ecosystems	58
9. Private and public benefits of soils and soil management	65
9.1 Introduction	65
9.2 What is the nature of benefits from improving agricultural soil condition?	65
9.3 Who benefits from improving agricultural soil condition?	66
9.4 How significant might these benefits be?	67
9.5 How might Australia realise these benefits? Examples through case studies	73
9.6 General findings	86
10. Summary and conclusions	89
10.1 Improving the organic matter status of soils	89
10.2 Improving the pH (acid-bases balance) of soils	91
10.3 Minimising erosion of soils by wind	92
10.4 Minimising erosion of soils by water	94
10.5 improvements in the quantity and quality of ecosystem services and benefits delivered from agricultural lands	95
10.6 Summary	98
References	99

Tables

4.1. List of critical functions of soil C	9
4.2 Dairy pasture management options to conserve soil carbon	15
5.1 Options for management of soil acidity and feasibility in permanent and mixed grazing systems	25
8.1: Description of the broad groups of ecosystem services provided by soils	49
8.2: Example of the beneficiaries of soil ecosystem services	53
8.3: Conclusions from this report about the effectiveness of management practices in Australian agricultural lands	55
8.4: Ways in which actions to address soil condition are likely to affect soil processes and ecosystem services	56
9.1: Gross value of agricultural production	66
9.2: Existing estimates of the value of costs or benefits related to land management practice (footnotes explained at end of table)	69
9.3: Full range of benefits and beneficiaries – Reducing soil erosion in broadacre cropping	76
9.4: Full range of benefits and beneficiaries – Managing acid soils in broadacre cropping	79
9.5: Full range of benefits and beneficiaries – Increasing soil carbon in irrigated horticulture	82
9.6: Full range of benefits and beneficiaries – Reducing wind erosion in grazing areas	86
10.1: Ecosystem services from soils and the benefits potentially derived	96

Figures

4.1: Crop management practice and relationship with expected Soil Organic Carbon levels and benefits	11
6.1: Erosion rates in relation to ground cover when four different wind speeds were applied to lupin residues	34
7.1: Factors influencing soil erosion by water. Figure was derived from various publications cited in the text	37
7.2: Generalised relationship between ground cover and annual average soil loss from vertisol soils on the Darling Downs, Queensland	42
8.1: Conceptual relationship between land management, soil structures and processes, ecosystem services, benefits to humans and human wellbeing	47
8.2: Interrelationships between living and non-living components of soils	48
8.3: Two generalised assessments of differences in ecosystem services from 'natural' ecosystems and agricultural land	52
9.1: Who benefits, where and when?	67
9.2: Example of output from the acidity relative yield model for four plant tolerance classes within a given Al/Mn solubility class	77

Boxes

Box S1: An example of benefits from better management of soil condition	x
Box 4.1: Managing soil C through a systems approach	18
Box 5.1: Managing soil pH through a systems approach	29
Box 6.1: Managing wind erosion through a systems approach	35
Box 7.1: The Gascoyne Catchment – A Case Study of Water Erosion	41
Box 7.2: Managing water erosion through a systems approach	44

Executive summary

Questions addressed

Funded under the Caring For Our Country program by the Australian Government's Land and Coasts Division, a joint initiative between the Department of Sustainability, Environment, Water, Population and Communities and the Department of Agriculture, Fisheries and Forestry, this project addresses two key questions about relationships between land management practices, soil condition, and the quantity and quality of ecosystem services (i.e. the attributes of ecological systems that contribute to benefits for humans) delivered from agricultural land:

- What evidence exists about how improving land management practices will lead to reduced soil loss (through water and wind erosion) and improved soil condition (especially through reduced impacts of soil acidification and increased organic matter content)?
- How might reducing soil loss and improving soil condition result in improvements in the quantity and quality of ecosystem services and benefits delivered from agricultural lands, including cleaner air, improved water quality, reduced greenhouse gas emissions, and more productive soils?

Key conclusions

The project focuses on four aspects of soil condition identified in the Program Logic for Caring for our Country's Sustainable Practices target: soil carbon; soil pH; wind erosion; and water erosion. It also focuses on four broad groupings of agricultural industries: broadacre cropping; horticulture; dairy; and grazing.

In summary, evidence in the scientific and economic literature assessed and referenced in this report finds:

- Approaches to improving the soil organic carbon (SOC) content of soils, including minimising disturbance to soils from tillage and stock and increasing inputs of carbon by retaining stubble, using perennial pastures, and adding manures and other sources of carbon, have slowed the rate of loss of SOC and show potential to increase absolute SOC over time (although predicting the outcomes of interventions precisely is still difficult due to the many variables involved). Benefits in terms of better production outcomes have been demonstrated.
- Regular monitoring of soil pH and application of lime at appropriate rates has been shown to reduce acidity in surface soils, although rates of adoption of these practices are far too low to achieve widespread benefits. Net financial benefits of controlling acidity in surface soils have been demonstrated. Build-up of acid in

subsoils is of growing concern and addressing it is likely to be unaffordable for most agricultural industries in the near future.

- Maintenance of ground cover above 50-70% has been shown to be effective in reducing wind and water erosion and to yield financial benefits to farmers across all agricultural industries.
- Addressing soil carbon, acidity and susceptibility to erosion has many public and private benefits. These include better yields of agricultural products, which have private and public benefits, and better outcomes for agricultural soils, which themselves provide a range of 'ecosystem services' and benefits to both farmers and the broader public. Better soil condition generally improves the ability of soils to support benefits to the public (both urban and rural), such as clean water for drinking and recreation, protection from wind and water erosion and floods, and reduced risks from pests and diseases and reduced need to use agricultural chemicals. They can also include a range of cultural, spiritual, and intellectual benefits such as enhancing sense of place, mental wellbeing and acquisition of knowledge. Modest improvements in soil condition might only produce modest improvements in these public services and benefits, but even these modest improvements can be significant in economic terms and often greater than the private benefits.
- One of the most substantial benefits of better management of groundcover is reductions in dust storms, which have been shown to incur very large financial costs in regional and metropolitan areas across Australia. These costs relate to damage to infrastructure and health costs, as well as clean-up costs and costs of reduced water quality. There have been substantial reductions in dust indices since the 1940s, but large and damaging dust storms have occurred recently and are likely to recur in coming years during prolonged dry periods.

Benefits and beneficiaries from better soil management

Ecosystem services can be described as the attributes of ecological systems that contribute to benefits for humans. By ecological systems, we mean systems that involve interactions among multiple species of plants, animals, and other organisms and between those species and the non-living environment. To address the question of how improving soil condition might result in improvements in the quantity and quality of ecosystem services and benefits delivered from agricultural lands, a framework was developed that relates soil properties and processes to ecosystem services, benefits and beneficiaries. The framework, described fully in the main report, is a synthesis and modification of several published frameworks. It was developed because many of those available in the literature did not explicitly link

changes in soil condition to benefits to people, and because those that addressed this link were not entirely consistent with a set of principles distilled from the most recent literature in this field. The key framework principles were:

- Contributions that ecosystems make to meeting human needs (ecosystem services) should be kept separate from the contributions made by humans that are required to turn ecosystem services into benefits (for example, ecosystems generate fertile soil but for that service to become the benefit of support for crops requires humans to plant, manage and harvest those crops);
- To avoid multiple counting of benefits, it is important to distinguish between 'final ecosystem services' (ones that can be turned directly into benefits) and 'intermediate' or 'supporting' ecosystem services (ones that support other services and therefore can contribute indirectly to multiple benefits).

The living and non-living components of soil ecosystems interact to mediate a range of processes that would require engineering at an unprecedented scale to replicate. These processes transform natural resources into forms that are potentially of benefit to humans and in so doing they are said to provide 'ecosystem services'. The main report identifies 14 such services and their respective benefits from soils.

Management of land for agriculture dramatically changes the balance among ecosystem services, increasing some provisioning services, decreasing some regulating services and changing the nature of many cultural services. One aim of improved agricultural management is to adjust this balance to meet a wider range of private and public needs.

Research reviewed in this report shows that best-practice approaches to managing soil carbon, acidity and wind and water erosion are generally effective at addressing those issues and improving soil condition. Practices like minimal tillage, maintaining ground cover above 50%, adding organic matter to soil (within limits), and managing the impacts of stock and machinery on soil disturbance and compaction, have beneficial outcomes for all aspects of soil condition. These practices, therefore, potentially enhance most ecosystem services and their benefits (Box S1).

The beneficiaries include farmers, agricultural industries, communities, families and individuals in regional areas and in cities. It is possible to estimate the magnitude of these benefits under different conditions in the future, but it is not meaningful to make a single estimate of future value because of the many combinations of management practices, soil types, climatic variations, products, market opportunities, demographic changes, and demands of consumers over the coming decades.

Some general conclusions can, however, be made:

- There are achievable opportunities to address declining soil carbon and increasing acidity and reduce wind and water erosion and at the same time improve profitability of agriculture and deliver a range of public benefits (which in some cases will be worth more than the private benefits in terms of health and wellbeing outcomes);
- To do this it will be important to consider the ability of soil ecosystems to cope with ongoing and potential future shocks (i.e., their adaptive capacity and resilience), which cannot be considered in isolation from the adaptive capacity and resilience of the humans who manage agricultural landscapes;
- The resilience of soils in many parts of Australia depends strongly on building and maintaining soil carbon stocks, which affect a wide range of functions, including nutrient cycling and water infiltration and storage, and the ability of landscapes to retain topsoil;
- Another key aspect of the resilience of Australian soils is their ability to avoid passing through thresholds of change, some of which could be irreversible;
- Such thresholds include critical proportions of ground cover (50-70% depending on factors like rainfall and slope), below which erosion accelerates dramatically, carbon-content thresholds, and thresholds of acidification, especially of subsoil, which currently cannot be addressed economically by most agricultural industries.

Box S1: An example of benefits from better management of soil condition

Maintenance of 50-70% groundcover — a management practice shown to be effective at reducing wind and water erosion and contributing to increasing soil carbon content and, indirectly, to addressing soil acidity — will affect the texture of soil by retaining the small particles that would otherwise be lost due to water and wind erosion. Organic matter content and biodiversity of soil will be enhanced because of reduced losses of carbon by erosion, increased inputs of carbon as groundcover plants die and degrade, and enhanced habitat for soil species. This will affect soil structure, soil biological activity and cycling of organic matter, nutrients, gases and water within soil and between soils and the atmosphere. These processes combine in different ways to support the full range of ecosystem services and their potential benefits. The extent of the benefits and the beneficiaries from maintaining ground cover will depend on the demand for different ecosystem services and benefits, who needs these and at what scales of space and time. The benefits are likely to be increased production of food and other commodities as well as a range of public benefits to people from local to regional, national and international scales.

1. Project rationale and approach

1.1 Rationale

Soils are a national asset, the condition of which is integrally tied to the health of Australian industries, ecosystems and, ultimately, communities. However, for a country for which the vagaries of climate variability have been manifested in dust storms and land degradation on the one hand, and rich production and economic wealth on the other, soils remain very much taken for granted.

Funded under the Caring For Our Country program by the Australian Government's Land and Coasts Division, a joint initiative between the Department of Sustainability, Environment, Water, Population and Communities and the Department of Agriculture, Fisheries and Forestry, this project addresses two key questions about the relationships between land management practices, soil condition, and the quantity and quality of ecosystem services delivered from agricultural land:

What evidence exists about how improving land management practices will lead to reduced soil loss (through water and wind erosion) and improved soil condition (especially through reduced impacts of soil acidification and increased organic matter content)?

How might reducing soil loss and improving soil condition result in improvements in the quantity and quality of ecosystem services and benefits delivered from agricultural lands, including cleaner air, improved water quality, reduced greenhouse gas emissions, and more productive soils?

The project focuses on four aspects of soil condition identified in the Program Logic for Caring for our Country's Sustainable Practices target: soil carbon; soil pH; wind erosion; and water erosion. It also focuses on four broad groupings of agricultural industries: broadacre cropping; horticulture; dairy; and grazing.

1.2 Approach

Literature review

This project is largely a desktop literature review, utilising some of Australia's leading soil, agricultural systems and ecosystem service researchers.

The Program Logic for Caring for our Country's Sustainable Practices target has identified four key aspects of soil condition in Australia, including carbon and pH (which are soil conditions) and water and wind erosion (which are threatening processes). Declining soil carbon and increasing acidity (which affect both the

physical properties of soils and a number of the processes occurring in it), and continuing susceptibility to wind and water erosion (which affect both the loss of soil from some sights and its build up in others) have been identified as key concerns in recent comprehensive analyses of agricultural and other landscape processes in Australia (NLWRA 2001). This project focuses on how land management practices affect these aspects of soil, and in particular:

the extent to which land management practices are available that can reduce erosion, increase soil carbon and slow rates of acidification; and

the degree of change likely to be possible from plausible changes in land management over a range of land and farming systems and a range of future time periods.

A second component of the project addresses the extent to which soil condition affects the quality of the market and non-market benefits received by people (so-called 'ecosystem services') from agricultural land.

Valuation of benefits from better soil management

The valuation of the benefits from changed land management practices is complex and requires a wide array of data on what changes might be made, who might make them and where, how those changes might affect ecological processes, and how those processes might affect ecosystem services and the benefits that flow from them. Because of this, the valuation component of the project makes assumptions and estimates upon which the valuations are contingent. The aim is to provide indications of the size of costs and benefits that might arise from improved soil management and the types of uncertainties that still remain in those estimates.

Based in the latest thinking about valuing ecosystem goods and services, the project develops a framework that makes explicit the links between:

- soil and other landscape processes
- landscape processes and ecosystem services
- benefits that potentially flow to a range of beneficiaries
- who the beneficiaries are likely to be
- how the value to those beneficiaries can be best assessed.

Valuations are based on realistic scenarios for marginal changes in land management practices in different regions and farming systems rather than any attempt to estimate the total value of all existing soil ecosystem services across Australia. Scenarios for changes in land management practices are developed from

the literature, the researchers' experience with a range of land-use systems over many years, and selected contacts with key experts on different land-use systems. The three scenarios used, as far as possible, reflect business as usual, modest improvements to farm management and optimistic improvements.

2. Soils: the essential asset

2.1 Soils, life and human interaction

Soils underpin, literally and figuratively all of the processes that support human societies and economies and, indeed, all other terrestrial life on earth. The overwhelming focus of both ecology and agricultural sciences has been on what happens above ground, which can be seen and experienced directly by humans. Soils play physical roles in supporting plants and structures, including those created by humans. They contain a vast diversity of living organisms and non-living elements that interact to mediate processes as diverse as provision of raw materials, water filtration, breakdown of wastes, pest control, regulation of atmospheric composition, regulation of water and wind flows across landscapes, and maintenance of hydrological cycles (Bardgett *et al.* 2001; Nelson and Mele 2006; Barrios 2007; Mele and Crowley 2008; McAlpine and Wotton 2009; Colloff *et al.* 2010; Dominati *et al.* 2010; Robinson *et al.* 2012). Soils also contribute in important ways to cultural, spiritual, intellectual and other intangible aspects of landscapes that are important to humans in many different ways (Dominati *et al.* 2010).

We are entering an age that has been termed the Anthropocene: an age when the impacts of humans represent the most significant drivers of change in Earth systems (Steffen *et al.* 2011). Thus, it is timely to consider how the tools available to humans have been and might be used to improve the functioning of soils, including reversing the degradation caused by past human activities.

2.2 Living soils and determinants of soil condition

Soil condition can be defined as *the capacity of a soil to function, within land use and ecosystem boundaries, to sustain biological productivity, maintain environmental health, and promote plant, animal, and human health* (Doran and Zeiss 2000). The condition of a soil can be inferred by measuring specific soil properties (e.g., organic matter content) and by observing soil status (e.g., fertility).

Maintaining soil condition is not only important to sustaining life and ecosystems beyond the immediate physical presence of soils, but also within. Soils are the home to over a quarter of all living species on earth (Turbé *et al.* 2010). Indeed, there is a strong relationship between soil condition and the biodiversity soils support. The many organisms and micro-organisms living within soils can interact to perform three major functions required of healthy soils: chemical engineering, biological regulation and ecosystem engineering. In the case of chemical engineering, bacteria, fungi and protozoans help in the decomposition of plant organic matter into nutrients readily

available for plants. In the case of biological regulation, small invertebrates, such as nematodes, pot worms, springtails, and mites, act as predators of plants and other invertebrates or microorganisms to regulate their dynamics in space and time. Finally, in the case of ecosystem engineering, earthworms, ants, termites and some small mammals help modify or create habitats for smaller soil organisms by building resistant soil aggregates and pores, thus regulating the availability of resources for other soil organisms and supporting plant systems.

Soil biodiversity is not the only determinant of soil condition. Soil can be defined as *the weathered and fragmented outer layer of the earth's terrestrial surface* (Hillel 1980), and the physical properties of soil such as particle size and mineral composition are important in its differentiation and condition. Moreover, the chemistry and nutrient status of soils are also important. However, it is the interaction of soil physics and chemistry with soil biodiversity that influences the overall condition of soils. For example, soil pH is one of the abiotic factors susceptible to influence biology and activity of biological regulators (Turbé *et al.* 2010). In every sense, the term *living soils* is a reminder that soils too have a lifespan that can either be cut short through inappropriate interaction or sustained by appropriate nurturing or remedial attention.

2.3 Soils and systems

This report considers the relationship between soil condition and agricultural practices in four distinct sections (i.e. sections on soil carbon, acidification, wind erosion and water erosion). These aspects of soil condition do not exist in isolation, however. For example, soil carbon content also influences susceptibility to erosion as soil carbon affects soil physical and chemical properties. Similarly many soil management practices, such as ground cover maintenance, address multiple aspects of soil condition (e.g., ground cover management can increase soil carbon and decrease soil erosion).

Across Australia many farmers and graziers face more than one form of resource degradation and most will have multiple objectives they seek to achieve. Some of these objectives will be economic, but certainly environmental and social objectives also play an important part in determining agricultural practice. Because of this, taking a systems approach to agricultural practice is not only theoretically important, but it also plays an important part in the day-to-day operations of Australian farms.

The extent to which systems approaches are well practised is an altogether different question. One of the aims of any system approach is to become efficient in achieving multiple objectives, and so in the context of this report the question arises: can good

practices be combined so they are additive and multiplicative, without negative impact. An example of a systems approach in managing soil follows. The traditional response to managing soil erosion on a grain farm may be to put in contour banks to reduce the length of water flow, hence its velocity and power – this prevents rills becoming gullies. Systems thinking would suggest that erosion is caused by runoff, adding soil sediment to the runoff and then the flow moving this across the landscape. Systems practice would be to reduce runoff by increasing infiltration, hence reducing sediment concentration, and managing the flow to maintain spread across the landscape and prevent runoff concentration (where rills and gullies form). This is usually achieved by management of ground cover.

At the conclusion of each of the soil condition Sections (4-8), a box has been included to provide an example of a systems approach to managing soil C, soil pH, water erosion and wind erosion.

3. Linking management practices, soil quality and ecosystem services

3.1 The concept of ecosystem services

One key purpose of this report is to consider the links between soil condition and the benefits that soils in good condition provide for humans. There is increasing demand from the public for agricultural landscapes to be 'clean and green' and to meet a wide range of society's needs (Soils Research Development and Extension Working Group 2011). Rarely, however, have these needs been fully and clearly articulated in the past, especially with respect to soils. Soils are often seen as simply the substrate in which plants grow. This narrow view has been changing over the past decade as there has been increasing focus on the roles of soils in ecosystems and their contributions to 'ecosystem services' and the benefits that flow from those services.

The dependence of humans on ecosystems has been the focus for a body of research over the past decade and more, under the banner of 'ecosystem services'. Ecosystem services can be described as the attributes of ecological systems that contribute to benefits for humans (Fisher *et al.* 2009). In Section 8 we discuss in more detail how ecosystems services are defined and categorised, and how the concept can be put into practice with respect to soils. The essence of the concept is that the multitude of interactions among living organisms in ecological systems, and between those organisms and the non-living components of the environment, produce outcomes that not only have great value to humans but can potentially be more efficient and less costly than alternatives that involve humans and their technologies (Daily 1997).

The types of benefits that come from ecosystems broadly (i.e., including above and below ground ecosystems) include: support for production of food, fibre, fodder and other products of crops; provision of chemicals and genetic material that can have value in human health and/or industrial processes; clean air and water; natural pest control; disposal of wastes; and a range of cultural, intellectual, spiritual and other intangible benefits. Obtaining these benefits usually requires some final input from humans, which is why several recent approaches have explicitly separated the services from the benefits (see Section 8).

Soils are at the heart of virtually all processes leading to ecosystem services and subsequent benefits (Daily *et al.* 1997; Sparling 1997; Wall and Virginia 2000; Barrios 2007; Soils Research Development and Extension Working Group 2011). Hence, any changes in soil condition potentially affect a range of processes, services

and benefits to humans. The changes in benefits are not, however, always readily attributable to soils as many involve inputs from other parts of ecosystems, such as plants, animals and atmospheric processes. As such, soils often provide 'intermediate' ecosystem services (i.e., services that support other services and therefore support benefits to humans indirectly rather than directly) (Fisher *et al.* 2008). In Sections 8 and 9, we explore how changes in soil quality relate to soil ecosystem services and how the value of those services can be estimated.

3.2 Ecosystem services and management practice

A focus of this study is the relationship between ecosystem services (their quality, quantity and diversity) and agricultural practice. We know from the history of agriculture that inappropriate practices may lead to land and water degradation and potentially to the loss of the productive resources upon which agriculture depends. Examples of this are provided in Sections 4 to 8.

It is important to note that the relationships between management practices and ecosystem services provided by soils are neither linear nor homogenous; what is a sustainable practice on one soil type within one climatic zone may not be sustainable elsewhere. Moreover, some practices may result in trade-offs between different ecosystem services. For example, tree planting to manage local erosion might enhance local productive capacity but the reduction in run-off may lead to less water being made available elsewhere. From a natural resource management perspective, this example may translate into the trade-off between managing dryland salinity and environmental river flows (van Buren and Price 2004).

The heterogeneity of Australian landscapes, Australian soils and Australian production systems demands heterogeneity in agricultural practices and policy approaches across our landscapes, our soils and our production systems. This makes determining an aggregated valuation of ecosystem services resulting from changes in practice very difficult, if not impossible, as discussed in Sections 8 and 9.

4. Soil Carbon

4.1 Nature of the issues

The global soil organic carbon (SOC) pool is estimated to be $\sim 1,395 \times 10^{15}$ g (Post *et al.* 1982) which is three times more than that found in the atmosphere or in terrestrial vegetation (Schmidt *et al.* 2011). SOC refers to the diverse range of organic material that enters (e.g. plants/ manures/ herbicides) or resides (e.g. soil animals and microbes) in soil. Soil therefore contains C in diverse structural forms and with diverse residence times, encompassing living (labile), recently dead and long-dead (non-labile and recalcitrant) forms. A comprehensive list of critical functions of soil C has been developed (Lal 2004) (Table 4.1).

Table 4.1. List of critical functions of soil C (after Lal 2004)

Function
Source and sink of principal plant nutrients (e.g., N, P, S, Zn, Mo)
Source of charge density and responsible for ion exchange
Absorbent of water at low moisture potentials leading to increase in plant available water capacity
Promoter of soil aggregation that improves soil tilth
Cause of high water infiltration capacity and low losses due to surface runoff
Substrate for energy for soil biota leading to increase in soil biodiversity
Source of strength for soil aggregates leading to reduction in susceptibility to erosion
Cause of high nutrient and water use efficiency because of reduction in losses by drainage, evaporation and volatilization
Buffer against sudden fluctuations in soil reaction (pH) due to application of agricultural chemicals
Moderator of soil temperature through its effect on soil colour and albedo (reflective capacity)

These functions of SOC can be associated with provisioning, regulating and cultural ecosystem services as well as the soil processes that support these services (MA 2005). They relate to water, air and food quality, nutrient cycling and disease control (Kibblewhite *et al.* 2008). SOC is considered a 'headline' soil condition indicator nationally and internationally. It is also a key component of greenhouse accounting programs used by the Australian Greenhouse Office (AGO) through the National Carbon Accounting System (NCAS) to track changes in carbon loss and storage under alternative land-use scenarios (Wilson *et al.* 2007). Further development of NCAS is supported by the Soil Carbon and Research Program (SCaRP) which examines variations in soil organic carbon (SOC) and composition under different agricultural management practices in regional Australia using a nationally consistent methodology (Sanderman *et al.* 2011).

4.2 Impacts of agriculture and measures that could build Soil Organic Carbon

There are many ways in which agriculture impacts on the capacity to build SOC. In principal, several factors influence this process reflecting that SOC dynamics is biologically mediated by a diversity of organisms that inhabit soils (see Section 2.2). Put simply, what determines the amount of SOC that accumulates is the balance between the amount of C added to the soil, the amount lost through microbial respiration and the capacity to build the resistance of what remains (Kirkegaard *et al.* 2007; Sanderman *et al.* 2010). Climate, and specifically precipitation and temperature, exert an overriding control whilst other regulators such as soil type, particularly particle size, nitrogen inputs, and plant biomass quality and quantity, are also important because they can be managed to some degree (Parton *et al.* 1987; Paustian *et al.* 1997).

It is also well recognised that land-use change has the most profound and enduring influence on SOC stocks. A global meta-analysis indicates declines in SOC stocks after land use changes from pasture to plantation (−10%), native forest to plantation (−13%), native forest to crop (−42%), and pasture to crop (−59%). Soil C stocks increase after land use changes from native forest to pasture (+ 8%), crop to pasture (+ 19%), crop to plantation (+ 18%), and crop to secondary forest (+ 53%) (Guo and Gifford 2002; Smith *et al.* 2012).

In Australia, clearing of native vegetation for primarily agricultural purposes has caused a 40-60% decrease in SOC stocks from pre-clearing levels. Significantly, some soils are still responding to the initial land-use change with continuing declines in SOC albeit more slowly under some management regimes (Sanderman *et al.* 2011) so it is critical that management not be considered only in relative terms (e.g. stubble retention versus stubble burning) but in the broader context of land-use change.


Also noteworthy is that while there is a strong theoretical basis for management strategies that build SOC, this is supported by a limited number of field studies (Sanderman *et al.* 2010) that generally lack management history detail (e.g. past and current management including fertiliser history, rotations etc) that is critical for estimating SOC build-up (Smith *et al.* 2012). This reduces confidence in making quantitative predictions about outcomes of interventions, but there is moderately high confidence in the efficacy of many approaches (Sanderman *et al.* 2010).

The relative efficacy of management strategies to mitigate SOC losses and to potentially build SOC, evaluated below for each of the four main industry groups.

Broadacre cropping

Broadacre cropping includes cereals, oilseeds, sugar cane, legumes, hops, cotton, hay and silage, and contributes around \$13 billion or more than 50% of the gross value of agricultural production in 2009-2010 (ABS 2011b).

Figure 4.1 illustrates the crop management options that are likely to or have been shown to increase SOC.



The nearly universally observed reductions in SOC that accompany clearing of native vegetation for agriculture have been attributed to two broad categories of process changes: reduced inputs due to harvest and stubble burning; and increased loss rates of carbon due to disruption of the soil surface, leading to enhancement of decomposition rates and greater risk of water and wind erosion (Sanderman *et al.* 2010). The potential approaches to increasing SOC, therefore, focus on reversing these effects (i.e., increasing inputs and/ or reducing losses). These management options include varying planting time, sowing rates, nitrogen application, cover and crop varieties, residue management (e.g. grazing and/ or burning), tillage type and depth, and length of fallow (Ugalde *et al.* 2007; Murphy *et al.* 2011). A combination of

these options, and specifically tillage and stubble management practices, can determine the SOC levels (Sanderman *et al.* 2010; Scott *et al.* 2010; Murphy *et al.* 2011) – although Sanderman *et al.* (2010) warn that the outcomes of changed management practices is not always predictable quantitatively because of the many factors that need to be taken into account. Some of these choices affect the stability of soil, while others affect yield and, therefore, biomass potentially available to the soil carbon pool.

The amount of carbon available for addition to soils in the form of shoot and root residues/ exudates depends on how much is removed at harvest. A broadacre crop such as wheat would produce less than 2 t.ha⁻¹.yr⁻¹ compared to sugar cane which might generate inputs of 7 t.ha⁻¹.yr⁻¹ (Kirkegaard *et al.* 2007).

Based on Figure 4.1, long fallow is *likely* to be associated with lowest expected SOC levels, and pasture cropping is likely to support the highest *expected* levels of C. Expectations for enhanced SOC are now high due to improved adoption of relevant practices (Barson *et al.* 2012b). Between 2007-08 and 2009-10 there was a national 10% increase (from 49-59%) in the number of farmers using reduced tillage, or one pass sowing systems and a 3% increase in farmers using residue retention. This resulted in residue being left intact over 68% of cropped area or no cultivation apart from sowing over 76% of cropped area.

Interpreting research on the effects of soil management practices on SOC is complicated because many studies have not been able to control all variables (Sanderman *et al.* 2010). For example, rainfall, soil type, time since last cultivation, and the depth at which measurements are made all affect SOC accumulation (see review by Sanderman *et al.* 2010). How sustained these increases are is also subject to conjecture as there are limited long-term studies of these systems across the five broad agro-ecological cropping zones (summer rainfall, Mediterranean west, moist south east, dry marginal south east and high rainfall zone) and rates of accumulation are highest in surface soils, which are also most vulnerable to disturbance. These temporal and regional data are critical in determining the likelihood of increasing SOC under the proposed management options and explains the high variability in SOC levels reported for direct drilled, stubble-retained systems (Mele and Carter 1993; Sanderman *et al.* 2010; Scott *et al.* 2010; Dalal *et al.* 2011).

Apart from the options of direct drilling and stubble retention to build SOC in some regions, Sanderman *et al.* (2010) highlighted that the greatest theoretical potential for building SOC is the addition of organic materials such as manure and green waste and the inclusion of a pasture phase in a cropping sequence. Due to their

relatively recent emergence there is very little scientific evidence that associates increased SOC in Australian broadacre cropping with practices such as organic matter amendment (e.g. manure, green waste and biochar) and pasture cropping (e.g. with perennial species). There is however strong evidence supporting the feasibility of pasture cropping in broadacre cropping systems (Bruce *et al.* 2006; Millar and Badgery 2009; Dolling *et al.* 2010) and the feasibility of biochar amendments (Chan 2008; Kimetu and Lehmann 2010; Singh *et al.* 2010) as potential strategies for increasing SOC.

If management enables SOC to build up, there is also a nutrient cost reflecting the heightened demand of soil biota for these nutrients as they decompose additional C substrates. The deficit created in nitrogen (N), phosphorus (P) and sulphur (S) over and above crop requirements is 60, 12 and 9 kg respectively per tonne of humus locked up (Passioura *et al.* 2008).

Horticulture

In 2009-10 Australia's horticultural industry was the nation's third largest agricultural industry based on gross value of production (GVP) of \$8.4 billion, ranking third behind the meat and grain industries (DAFF 2012b).

Horticultural industries encompass a diverse range of fruit and vegetable industries. The total area under production in Australia is around 250,000 hectares. Generally, interest in SOC is driven by the need to mitigate greenhouse gas emissions and to improve soil health and resilience (the capacity to recover after disturbance). A survey commissioned by Horticulture Australia limited (HAL) in 2000-2003 indicated that the most important building block for healthy soil, irrespective of soil type, region, or climatic conditions was SOC.

A comparison of SOC in intensively managed vegetable production sites with 'reference sites' in Tasmania and Queensland led to the conclusion that 'good farm management practices, even for intensive land use for vegetable production, can sustain soil integrity/ soil health' (HAL 2003). A recent investigation into on-farm emissions in Bundaberg regions and in the Lockyer Valley and Bowen indicated that vegetable production was the highest emitter of C from soils ($3.50 \text{ tCO}_2\text{-e.ha}^{-1}\text{.year}^{-1}$) followed by tree crops ($2.85 \text{ tCO}_2\text{-e.ha}^{-1}\text{.year}^{-1}$), then sugar cane ($1.91 \text{ tCO}_2\text{-e. ha}^{-1}\text{.year}^{-1}$) then cane/ other crops ($1.16 \text{ tCO}_2\text{-e.ha}^{-1}\text{.year}^{-1}$). This trend was reversed when calculated as emissions per unit income (e.g. vegetables $41 \text{ tCO}_2\text{-e}/\1 million , fruit trees $221 \text{ tCO}_2\text{-e}/\1 million and cane $606 \text{ tCO}_2\text{-e}/\1 million). It was concluded that, despite the high variability in data within a production system, there was

significant scope for improvement with carbon fixed in organic matter as a recommended management option (HAL 2012b).

The vegetable industry's key management messages are to use minimum-till techniques and controlled traffic technologies and to add organic materials (such as organic mulches and biochar) to build SOC (Pattison *et al.* 2010; HAL 2010, 2011). A detailed study on the use of organic products (chicken manures, composted green wastes) for multiple benefits confirmed that additions of organic matter in these ways both offset carbon losses experienced in conventional approaches to vegetable management and increased crop productivity by up to 10% when other inputs were held constant (HAL 2011). A survey of soil management from 2007-08 to 2009-10 indicated that 28% more horticulturalists used alternate or cover crops and 33% used mulching or matting (Barson *et al.* 2012c).

Dairy

In 2010-11 the farm gate value of production for the dairy industry was \$3.9 billion (around 10% of the gross value of Australia's agricultural production) and the total area under production was 4 Mha (Barson *et al.* 2012a; Dairy Australia 2012). Generally, dairy systems have higher levels of SOC relative to other agricultural industries and therefore the focus is less on building SOC and more on maintenance or loss prevention (MacKenzie 2010). Higher levels of SOC are attributed to a number of factors such as: higher availability of water (as rainfall or irrigation); ready supply of nutrients (N and P); higher proportion of perennial species that grow continually rather than seasonally; minimal disturbance relative to cropping; and minimal erosion.

Loss of soil carbon from dairy soils does occur and has been attributed to loss of ground cover due to high stocking rates, leaching of organic acids below the root zone, and to cultivation associated with planting of annual grasses in dryer or drought prone regions such as in northern Victoria (MacKenzie (2010) reviewed experimental results from several countries as well as Australia). Management options to prevent loss of carbon in dairy pasture soils are: 1) to reduce decomposition; 2) to improve the rate of addition of organic materials; and 3) to reduce soil disturbance/ increase ground cover (Watson 2006; MacKenzie 2010; Barson *et al.* 2012a). These options are summarised in Table 4.2 together with the likelihood of adoption.

Table 4.2 Dairy pasture management options to conserve soil carbon (drawing on a research review by MacKenzie (2010) and a survey of practices by Watson (2006))

Management option	Rationale	Current likelihood of adoption
Slow the rate of decomposition of soil carbon	Clay soil tends to protect organic matter more effectively from decomposition than sandy soil.	Unlikely; on most farms, increasing clay content through techniques such as clay spreading is prohibitively expensive.
	Subsoil modification of hard pan or sodic/ Al toxic layers to encourage root penetration to deeper (cooler) layers	Unlikely; Subsoil modification costs can be high despite the likely high returns in a short timeframe (MacEwan <i>et al.</i> 1992).
	Organic materials such as biochar, waxy plant materials, and composted manure have chemical structures can potentially reduce the rate of organic carbon decomposition in soil	Likely where material is readily available and inexpensive (i.e. where financial returns are expected to exceed the costs of purchase and application). Unlikely where input material is not retained (is decomposed) and where there are other costs in terms of nutrient tie-up i.e. efficacy questionable due to scientific uncertainty (Passioura <i>et al.</i> 2008; Schmidt <i>et al.</i> 2011; Jones <i>et al.</i> 2012).
Increase the rate of addition of plant biomass	Use of ameliorants such as gypsum (for sodic soils) and lime (for acid soils) to increase plant productivity	Unlikely due to fluctuating production costs which means it is not always economically viable to correct the problems with gypsum and lime (refer Section 5); main issue is pasture utilisation rather than biomass. It should be noted that sub-soil acidity is a problem in some dairying areas (Section 5).
	Use of essential elements (e. g. N, P, S, K, Ca) to increase C transformations and optimise productivity	Unlikely to be viewed as a strategy to increase C build-up <i>per se</i> but as a means of increasing pasture biomass.
Reduce soil disturbance (pugging, tillage) increase ground-cover	Livestock management (stocking rates/ grazing intensity to protect ground cover)	Likely but requires pasture renovation as well
	Pasture renovation (increasing perennials in sward composition).	Likely but requires livestock management as well

In terms of current trends in management (2007-08 to 2009-10), dairy farmers are increasingly monitoring ground-cover (up from 72% to 88%) but fewer are setting ground-cover targets (38% to 27%) (Barson *et al.* 2012a).

Grazing

Livestock grazing is the most widespread Australian land use, covering more than 336 Mha or about 40% of the total area of Australia. Meat and wool production contribute almost 30% to the gross value of agricultural production (ABS 2011a). These enterprises encompass three broad systems; i) the native pasture dominant

systems, principally occurring in the rangelands of central and northern Australia, ii) the permanent perennial grass-based pasture zones of south-eastern Australia and iii) the more intensive mixed wheat-sheep farming systems of southern Australia that are based on improved pastures and fallow rotations (Scott *et al.* 2000; Australian State of the Environment Committee 2011).

Grazing by livestock (e.g. beef and sheep) can impact directly on SOC and nitrogen cycling by modifying plant biomass inputs into soil (shoot and root material) and by reducing ground cover and thereby exposure of SOC-rich surface layers to wind and water erosion (Earl and Jones 1996). Grazing can also impact indirectly on SOC by modifying soil structure (density and aggregate stability), moisture and temperature influencing soil faunal and microbial diversity and activity (Southon and Cattle 2004b; Teague *et al.* 2011).

Management options to increase SOC have focussed on three strategies: 1) increased productivity (irrigation and fertilisation); 2) time controlled (TC) or rotational grazing; and 3) shift to perennial species (Sanderman *et al.* 2010). Research on the impacts of these options on SOC is rare (Sanjari *et al.* 2008; Sanjari *et al.* 2009), despite the extensive research effort in sustainable grazing systems and, specifically, increasing the perenniality of pasture systems (Kemp and Dowling 2000; Mason and Kay 2000; Michalk *et al.* 2003). The emergence in the late 1980's of grazing systems referred to variously as 'cell grazing', 'savory grazing', 'short duration grazing', 'time-controlled (TC) grazing' and 'holistic management (HM) grazing' have been assessed for their impact on a range of sustainability measures including SOC (Earl and Jones 1996; McCosker 2000; Sanjari *et al.* 2008; Sanjari *et al.* 2009; Sherren *et al.* 2012). A small number of studies in south-eastern Queensland and northern NSW of TC grazing have reported increases in herbage mass, SOC, nitrogen (Sanjari *et al.* 2008), ground-litter (Earl and Jones 1996; Sanjari *et al.* 2008), and reduced runoff and soil loss (Sanjari *et al.* 2009) compared to continuous grazing. Longer monitoring periods would increase confidence in these data (Sanjari *et al.* 2008; Sanjari *et al.* 2009).

4.3 Evidence of the efficacy of practices to increase soil organic carbon

In theory, the two main ways to build soil C are to reduce gaseous loss as either CO₂ and CH₄ by reducing soil disturbance and to increase C inputs either in the form of more plant biomass (which may require measures to overcome other constraints to plant growth) or in the form of other organic materials (manures, biochar etc). In practice, only the cropping industries (broadacre and horticulture) have opted for

reducing disturbance of surface residues and increasing inputs through plant residue retention and through the addition of organic residues as strategies to increase SOC. The grazing industries (including dairy) have focussed more on maintaining SOC through indirect means such as increasing ground cover and arresting acidification.

The efficacy of practices to increase SOC is highly variable and is dependent on soil type (particle size) and climate (regional precipitation patterns) (Smith and Belvins 1987; White 1990; Mele and Carter 1993; Kirkegaard *et al.* 2007). The consensus is that, in most of the cereal cropping areas in Australia (rainfall of 250-600 mm), the potential for reduced or no-tillage (direct-drilling) and stubble-retention to store carbon and mitigate greenhouse gas emission is limited, in contrast to areas with higher rainfall and greater biomass production (Sanderman *et al.* 2010; Chan *et al.* 2003). In a review of stubble retention systems in southern Australia, the higher SOC levels under stubble retention practices (relative to stubble burnt treatments) was not attributed to the sequestering of C but rather to the slower rate of decline under stubble retention compared to burning (Scott *et al.* 2010). The higher levels of SOC in surface soils of no-till systems can be associated with other benefits such as increased infiltration, reduced disease, conservation of nutrients and increased earthworm densities (Carter and Steed 1992; Roget 1995; Simpfendorfer *et al.* 2004; Scott *et al.* 2010) which may represent a more sensitive, yet indirect measure of the benefits of SOC increases with minimum tillage and stubble retention.

For horticulture, dairy and grazing industries, evidence of the efficacy of management strategies to increase soil C is difficult to find in the primary literature. For the grazing industries, only a very small number of studies have measured changes in SOC directly (Sanjari *et al.* 2008) and the confidence in these data was low due to the relatively short time frame for monitoring differences in TC and continuous grazing systems.

The general principles that have been demonstrated in using broadacre cropping industries as the model can also be applied more broadly. Empirical data have increased confidence in the application of models to predict soil C build up (e.g. CENTURY/ROTHC), which can be useful when it is not possible or affordable to collect SOC data.

Box 4.1: Managing Soil C through a systems approach

System goal

To increase soil C or slow down its decline.

Considerations

1. Increase inputs by growing more biomass (relative to removal), adding fertiliser and ameliorants as required, growing perennials or increasing crop frequency, and adding organics (mulch, manure, compost). These practices are interactive and probably cumulative. Appropriate performance indicators would be water-use efficiency and nitrogen-use efficiency, as an optimal balance between carbon and nutrients improves water-holding capacity of soil, microbial involvement in carbon and nitrogen cycles, and efficiency of nitrogen use for growth by plants. These actions potentially apply to cropping, horticulture, grazing and dairy.

2. Reduce decomposition by: avoiding excessive soil moisture and waterlogging; eliminating tillage, burning and erosion; reducing NO_3 fertilisers, changing to NH_4 fertilisers, organics or legumes; and encouraging free-living N fixation. These actions are applicable across industries.

3. With 1 and 2, operate at a stable soil C level, not increasing. This level needs to be determined but will be higher for currently degraded soils. Maintenance inputs depend on soil C levels, lower is better. Soil C also ties up large amounts of nutrients. Should our goals be equilibrium soil C and increased C cycling of the C inputs from 1 and 2? It is difficult to increase C inputs and soil C in cropping industries with the high product removal required for viability and efficiencies.

Recommended practices

Zero tillage, increased crop frequency or perennial pastures to increase biomass production and retention, residue retention or managed grazing pressure, improved agronomy, organic fertilisers, no burning.

Performance indicators

Annual water-use efficiency and nitrogen-use efficiency, carbon and nutrient cycling (most relevant at farm scale), percentage ground cover (most relevant at farm to regional scales), and productivity (relevant at farm to regional and national scales).

Conflicts

Availability and costs of machinery for managing minimum till can be a limiting factor. Incentives may be needed to move some farmers from traditional practices. Management inputs can be high to achieve enhanced SOC.

5. Soil pH

5.1 Nature of the issues

Soil pH (potential hydrogen) is the test used to assess the concentration of hydrogen ions in soil solutions of water (pH_W) or calcium chloride (pH_{Ca}). Ideally, soil pH for crop and pasture production should be in the range of pH 5.5 to 7.5_{Ca} in the top soil, and no less than pH 4.8_{Ca} in the subsoil (Dolling *et al.* 2001; Gazey and Davies 2009). Soil acidification, a key soil condition indicator (NLWRA 2007) is measured by a decline in pH over time. This can occur in the surface and subsurface layers of soil. There are several major causes for the acidification of agricultural soils: removal of agricultural products (most plant and animal products from farms are slightly alkaline); excessive accumulation of organic matter, which contains organic acids, in some circumstances (even though soil carbon also plays a key role in buffering against pH change); excessive use of nitrogenous fertilisers, especially those that lead to release of ammonia into the soil; leaching of fixed, fertiliser and urine-N as nitrate from surface layers to lower layers before plants can utilise it (Scott *et al.* 2000; NLWRA 2001; Gazey and Davies 2009). Understanding the causes will be critical for addressing questions on the efficacy of remedial action in different agricultural land-use scenarios.

The effects of acidification are not easily recognised and hence it is commonly described as an insidious problem in that plant symptoms are less visual and easily misdiagnosed, and production declines are gradual (Scott *et al.* 2000). Impacts can be on-site and related to plant, animal and soil biological performance or off-site, though the link to stream and groundwater acidification is speculative (Cregan and Scott 1998). On-site impacts are usually associated with increases in aluminium (Al) and manganese (Mn) levels with plant toxicity symptoms emerging and a reduction in nutrients such as calcium (Ca), Magnesium (Mg), and Potassium (K) with plant deficiency symptoms emerging (Slattery *et al.* 1989). The reduction in plant biomass production has a major knock-on effect; it reduces the quantity and quality of plant residue entering soils and hence SOC levels and all the associated critical functions (see Section 4, Table 4.1).

Acidification occurs in surface and in subsurface soils. According to the National Water and Land Resources Audit of 2001 (NLWRA 2001), half of the non-rangeland agricultural land in Australia is acidic (surface $\text{pH}_{Ca} \leq 5.5$) and below the optimal level to prevent subsurface acidification. This area, estimated to be of the order of about 49-50 Mha, is 5 times greater than the area affected by salinity. About half of

this, or approximately 17 Mha, has $\text{pH}_{\text{Ca}} \leq 4.8$ and requires immediate remedial action. In WA, almost 8 Mha of the 13 Mha under dryland agriculture are at risk of acidification (Holmes *et al.* 2011). In southern Australia, subsoil acidity occurs on about 24 Mha (Li *et al.* 2010).

Ten years on, the State of the Environment report (Australian State of the Environment Committee 2011) highlights that the severity and extent of acidification has increased in many regions, due, it says, to inadequate treatment, intensification of land management, or both. Although, for three of the four main agricultural industries, the number of businesses applying lime or dolomite to their holdings increased between 1995-96 and 2009-10, the totals by 2009-10 were only between 17 and 21% and most of that increase had occurred by 2001-02 (DAFF 2012a). For cropping, this increase was from 8 to 17% between 1995-96 and 2001-02, rising to 19% by 2009-10 (DAFF 2012a; Barson *et al.* 2012b). Dairy and horticulture started at higher percentages but achieved much smaller increases (DAFF 2012a).

Of even greater concern is the largely unknown extent of subsoil acidification and the intergenerational issues that will arise if this develops to levels where mineral dissolution occurs and soils are beyond remediation. It is clear that subsoil testing to raise awareness of the issue is a critical first step with early evidence of a change in attitude and intention in farmer groups (e.g. Nyabing group) in WA (Wilson *et al.* 2009; Gazey *et al.* 2012).

5.2 Impacts of agriculture and measures that could arrest soil acidification

Broadacre cropping, horticulture, dairy, and grazing all contribute to soil acidification. The Australian State of the Environment Committee (Australian State of the Environment Committee 2011) listed the following summary observations:

- Soil acidification is widespread in the extensive farming lands (cropping, sheep and cattle grazing) of southern Australia;
- Rates of lime application are well short of those needed to arrest the problem;
- Acidification is common in intensive systems of land use (tropical horticulture, sugar cane, dairying);
- Acidification is limiting biomass production in some regions, but the degree of restriction is difficult to estimate;
- Carbon losses are most likely occurring across regions in poor condition, and soil acidification is a major constraint on storing carbon in soils in the future.

Acidification risk areas based on topsoil data from major agricultural land-use categories have been identified (based on a 5 km grid) as a priority for remedial management (Wilson *et al.* 2009). The specific agricultural activities that increase soil acidity are the use of high-analysis nitrogen fertilisers, the large rates of product removal, and the farming of soils that have a low capacity to buffer the decrease in pH (e.g. infertile, light-textured soils) and the soil already has a low pH (Helyar *et al.* 1990; Helyar 1991; Wilson *et al.* 2009).

The five primary actions to address soil acidification are to:

- soil test for pH
- add lime at rates that are effective for arresting acidification
- add lime at high rates, sufficient to reverse acidification in soils that have already acidified
- use acid-tolerant plant species where available (as a short-medium term measure).
- land retirement (this could be considered where it is uneconomic to apply lime and where the benefits of arresting acidification are judged to be sufficiently important – this has not occurred anywhere in Australia to date to our knowledge).

Testing surface and subsurface pH by farmers, on-farm, is the precursor to implementing remedial action. The number of landholders who undertake pH testing has declined slightly (from 07-08 to 09-10) across all industries (grains, horticulture, dairy and grazing) with Queensland being the exception with slight increases in all but the grazing industries (Barson *et al.* 2011, 2102a, b, c). Lime addition and use of acid tolerant species are complementary actions with the fifth action, land-use change, being a more extreme option and not usually considered. The use of acid tolerant species, although a relatively straightforward and cost-effective option, does not address the underlying problem, proving a temporary strategy for 'living with the problem' and probably making it worse. The most widely used remedial action is to add lime to increase surface soil pH and gradually subsurface pH. Information on the neutralizing values of liming material (Goldspink and Howes 2001) and the recommended rates to apply in pasture and cropping systems (Slattery *et al.* 1989; Gazey and Davies 2009) are readily available and supported by online lime calculators for choice of lime, amount to add, and economic benefit (e.g. <http://www.aglime.com.au/liming>; <http://www.soilquality.org.au>).

The adoption of these five primary remedial actions is ultimately influenced by return on investment which is set by regional factors of soil type and rainfall (Helyar 1991; Gazey and Davies 2009; Holmes *et al.* 2011). The impacts of soil acidification and practices that are available to address this widespread problem will now be considered in the context of the four main industry groups.

At a national scale, protocols for monitoring soil pH are established (Grealish *et al.* 2011) but an organised national monitoring system has yet to be implemented.

Broadacre cropping

A consequence of the intensification of broadacre cropping over the past 10-15 years (see Section 4.2) is greater N-fertiliser use and greater product removal leading to increased rates of soil acidification. Liming is regarded as an economically viable option for broadacre cropping, and a lime application strategy must account for a range of factors including type of crop and level of production, type of lime and amount applied, soil texture and rainfall (Slattery *et al.* 1989; Helyar 1991; Helyar *et al.* 1992; Gazey and Davies 2009).

The key management messages for broadacre croppers are that:

- Lime rates should be matched to the soil type and soil pH. The lime requirement (as dolomite or limestone) to raise pH by about one unit varies by soil type, with rates increasing from about 1.5 to 2.5 t/ha of good quality lime on sandy soils to up to 6 t/ha on clay soils (Slattery *et al.* 1989; Aitken *et al.* 1990; Gazey and Davies 2009).
- Varying the rates of lime applied to soils has proved more cost effective than uniform application. This accounts for paddock variability in soil type (see above) and to variable rate N fertiliser applications (Bruce *et al.* 2006).
- Soil samples to assess pH should be taken to depth (down to 30 cm) and composited to account for spatial variability (Slattery *et al.* 1989; Holmes *et al.* 2011) and to assess the occurrence of subsoil acidification (Gazey *et al.* 2012)
- Soil pH should be monitored every three to four years to assess the impact of management and amelioration treatments (Holmes *et al.* 2011).

Lime rates should also consider the crops grown to account for varying tolerances and for loss of alkalinity through product removal (Slattery *et al.* 1989) and to N fertiliser rates to account for increased acidity through nitrate-N drainage (Bruce *et al.* 2006).

Horticulture

The use of high analysis N fertilisers and the high rate of product removal are features of most horticultural enterprises. Horticulture Australia limited (HAL) reports that 11 of the 21 horticultural industries supported by HAL have undertaken soil research (e.g. strawberries, citrus, bananas, blueberries, deciduous orchards, macadamias, and nursery, potatoes, processing tomatoes, turf and vegetables) to counter the problems associated with high fertiliser inputs and product removal. Soil acidification has been identified as one of the six main issues of concern (Horticulture Australia Ltd 2008).

The key management options for mediating soil acidification in horticulture are similar to those for broadacre cropping with liming a key strategy. Nationally about 20% of horticultural businesses apply lime/ dolomite and 25% use pH and nutrient testing (Barson *et al.* 2012c). Horticulturalists tend to use burnt lime (CaO) which reacts more quickly with water (Goldspink and Howes 2001). For intensive industries such as vegetable growing, the high N fertiliser use coupled with irrigation represents a significant risk for acidification through nitrate leaching below the root zone. In extensive perennial-based dryland systems, (e.g. orchards and vineyards), particularly those located in the high rainfall zone, the use of acid tolerant species such as chestnuts and the liming of soils for grape production is recommended (McCarthy *et al.* 1992; Scott *et al.* 2010). The recommended pH_{Ca} for grapevines is 5.5 to 7.5. Outside this range they are likely to suffer toxicity (Al) or deficiency (Fe, Cu, Zn and Mn) (White 2009). Data recording the extent to which lime is applied under vine in Australia is difficult to find.

For many horticultural industries, the cost of liming is relatively small in relation to yield profit so it is more likely that the condition of these soils won't decline from acidification compared to the broadacre cropping industry. As with broadacre industries, liming can be an effective and profitable management strategy for mitigating surface soil acidification provided appropriate rates are applied that account for regional and local (management) factors of soil and plant type and N-fertiliser regimes.

Dairy

Eight of the major dairying areas in Australia occur in the higher rainfall zones (600 mm) of southern Australia (Southern Queensland and Northern NSW) and southern Western Australia. Around 63% of intensively managed grazing, including dairy pastures, area is at low risk of soil acidification (particularly in SA and NSW) and

21% is at high risk (particularly in WA and Vic) (Barson *et al.* 2012a; Dairy Australia 2012).

Due to diminishing returns from milk production dairy farmers nationally have intensified and diversified their production to remain profitable. This has been done by increasing stocking rates, growing irrigated annual fodder crops, moving to mixed livestock systems of beef and dairy, and increasing nutrient inputs (Gourley *et al.* 2007; Bolland and Russell 2010). Many dairy farms also report significant nutrient surpluses, either as a result of high N application rates or by importing feed on farm (Gourley *et al.* 2007). The net effect of these activities is significant acidification, particularly in light textured soils where soil buffering capacity is low. The situation is particularly serious in south-western Australia where most soils used for dairy production have acidified from pH_{Ca} values 5.5–6.5 to pH_{Ca} 3.7–4.5 (McArthur 2004). Aluminium toxicity, induced by soil acidification, is a major problem for dairy production (Bolland and Russell 2010) and is ameliorated by applying sufficient lime to raise the pH of the top 0.10 cm of soil to ≥ 5.5 (Whitten *et al.* 2000). The rate of change was slow, with pH_{Ca} of 5.5 achieved in individual paddocks 9–11 years after the liming program started, with 29% of paddocks not achieving this level despite additions of between 12–21 t/ha lime (Bolland and Russell 2010).

Grazing

Acidification-remediation actions for grazing lands are confined to permanent pasture and mixed farming zones, and subsequent discussion will focus on these systems.

Under grazed permanent pastures, nitrate leaching is considered to be the largest contributor to acidification (Ridley and Coventry 1995). In south eastern Australia (e.g. NSW southern Tablelands and north-eastern Victoria), Scott *et al.* (2000) highlighted three characteristics of acidification; i) the rate of pH decline is slow (50 years or more) and even slower on strongly acidic soils ii) acidity problems are more quickly apparent on light textured soil and iii) soil can be acidic to depths of 60 cm.

The options for managing acidification under grazing systems are listed in Table 5.1 together with the associated constraints (Scott *et al.* 2000). These options are related to increasing perennial pasture content for better uptake of nitrate and for better year round biomass production (Section 4). Specifically there are four listed: 1) to sow perennial grass species rather than annual to access nitrate and prevent leaching; 2) to incorporate agroforestry systems, again to increase rooting depth and nitrate uptake; and 3) to reduce stocking rates on pastures with a high component of native grasses, to maintain vigour of native grasses. This last option will only constitute a minor component of grazing systems (less than 10%) and will therefore

not apply in many cases. Ultimately liming at higher rates is the major solution to reduce soil pH below 10 cm and benefit-cost scenarios for different soil types and rainfall distributions must be articulated.

Table 5.1 Options for management of soil acidity and feasibility in permanent and mixed grazing systems (adapted from Scott *et al.* 2000)

Option	Feasibility	Considerations
1. Modifying the grazing system <ul style="list-style-type: none"> change pasture species and/or grazing management use less fertiliser 	Limited (in permanent pasture systems due to cost and management skills, and also limited to area). This option will also only reduce acidification	Perennial species (e.g. native grasses) <ul style="list-style-type: none"> some scope but very high establishment costs Modification of animal camping behaviour <ul style="list-style-type: none"> high investment in labour, management skills and fencing Increase stocking rate <ul style="list-style-type: none"> likely if farmers more able to afford lime Reduce stocking rate <ul style="list-style-type: none"> likely where there is a reasonable proportion of summer-active native grasses profitability likely lower except maybe for fine wool production Fertiliser use <ul style="list-style-type: none"> avoid elemental S and NH_4^+- fertilisers, otherwise must apply lime to balance (3-7 kg per kg S and N respectively)
2. Breeding and selecting plants for tolerance	Feasible in permanent and mixed grazing systems but is a temporary solution only	Selection of Aluminium tolerant species - most ryegrasses, native grasses, oats and triticale are highly tolerant but can mask and intensify developing problem and does not negate need for lime Breeding must consider other traits such as palatability, persistence and the response of the rhizobial symbiont to acidity. Selection of aluminium tolerant plant varieties and rhizobial strains can be useful as a short –medium term solution (Ridley and Windsor 1992) but can exacerbate acidification in the long-term.
3. Correcting acidity by lime application	Highly feasible but amounts required and time taken dependent on soil type and grazing system (permanent or mixed)	Lime (carbonate) movement is slow <ul style="list-style-type: none"> takes time to move into soil profile, depends on porosity, can be facilitated by tillage and/ or soil fauna higher clay and organic matter soils resist change higher lime rates increase pH to greater depth surface applied lime increases profile pH to greater depth than incorporated lime (Ridley 1995) Response of subterranean clover-based pastures to liming is promising <ul style="list-style-type: none"> sub clover response but variable in magnitude and time; the required 30% increase in stocking rates for economic response has been reported (e.g. Book Book NSW) some nutrients less available limiting rhizobial survival sub clover response less reliable where lime surface applied but likely a matter of time (Ridley and Windsor 1992) Response of perennial-based pastures to liming is promising <ul style="list-style-type: none"> Phalaris, cocksfoot (DM increases) (Ridley and Windsor 1992) Plant yield response is often related to depth of lime incorporation and to rate of application <ul style="list-style-type: none"> the rate of lime required varies with soil type (Ridley 1995)
Management option	Feasibility	Considerations
4. Changing land-use	Technically feasible, politically very difficult!	Forestry/ land retirement means acidification slowed/ less relevant <ul style="list-style-type: none"> forestry is too costly on slopes >20%, location of infrastructure for harvesting trees Land retirement will require public funding Horticulture and cropping means lime amendment is economically achievable (refer above section)

5.3 Evidence of the efficacy of practices to increase soil pH

This section will address the issue of efficacy against the 4 practices listed above.

Test soil for pH

The motivation to test soil requires knowledge of the problem (why it is necessary), instruction on a statistically meaningful sampling design (how to collect the sample), awareness and instruction on best course of action to increase soil pH, and knowledge of economic benefits couched in realistic timeframes. Commercial soil testing facilities are readily available and instruction on testing design is established or under refinement to take greater account of spatial variability and temporal factors that account for the slow rate of change in soil pH (Holmes *et al.* 2011). Yet soil testing for pH (monitored since 2007/08) has declined in 2009-10 (Barson *et al.* 2011; 2012a; b; c). Reasons for this decline are unclear and are likely to be complex and multifaceted (Pannell and Vanclay 2011). Significant motivation will be generated by the promotion of regional data demonstrating the significant benefits to be derived from managing soil pH and the development of a 20-year, \$75 million national soil pH monitoring program (noting that this national program is separate from programs aimed at encouraging local testing) (Grealish *et al.* 2011).

Add lime at rates that are effective for arresting acidification

There is compelling evidence to support the view that the management of soil acidification by liming surface soils can yield significant benefits for broadacre cropping industries. In a long-term trial, known as 'managing acid soils through efficient rotations' (MASTER), wheat crops produced on average, 1.6 t/ha more grain on the limed (2-3.6 t/ha) treatments. Sensitive (barley and wheat) and acid tolerant cereal varieties (e.g. Dollarbird) also yield more (1.6-2 t/ha more) in limed soils (Li *et al.* 2001; Carr *et al.* 2006). Lime-induced yield increases of a similar magnitude have been reported widely in southern Australian broadacre cropping systems in plot trials (Coventry *et al.* 1987; Coventry *et al.* 1989; Slattery *et al.* 1989), even in the presence of soil borne diseases (Coventry *et al.* 1987). According to Li *et al.* (2010), this success, combined with strong grain prices resulted in anecdotal reports of exponential increases in lime applications in the area in the 1990s.

A more recent case study conducted in the Gabby Quoi Quoi Catchment of the Avon River basin in Southern WA, highlighted the increases in soil pH values measured at approximately 300 sites over a 7-year period (1999-2006) after liming (Carr *et al.* 2006). This study reported that 75% of the topsoil and 85% of the mid-soil sampled in 1999 had pH_{Ca} values lower than 5.0, with 15% of these soils having pH values less than 4.0. Re-sampling in 2006 has showed an overall increase in soil pH_{Ca} with

60% topsoil and 69% mid-soil being less than 5.0_{Ca} and no samples found to be below pH 4.0. Yield responses were also measured in wheat (\$28/ha), barley (\$53/ha) and lupin (\$5/ha), although in the latter crop, lime costs were not covered by the increased yield.

In the diverse industries that are collectively grouped into horticulture, the addition of lime is viewed as one of the management strategies for improving the overall health of soils. There are no accessible studies available on the effects of lime rate on biomass production in this industry. The high inputs applied and the short growth phases of vegetable production systems means that the lime-induced response is difficult to assess. Lime addition is therefore seen more as a general soil health maintenance activity (AusVeg 2010).

Despite positive yield responses, national trends in lime/ dolomite use (Barson *et al.* 2011; 2012a; b; c) to manage acidification suggest that there hasn't been much change since 2000/01 or there has been a slight decline depending on industry and state. Many suggest that this could be related to the 10 years of drought during this period. For cereals (majority of broadacre cropping) nationally there was an increase in the percentage of farmers using lime/ dolomite from 1995/96 to 2000/01 but not much change since (except in WA and Tasmania) (Barson *et al.* 2012b). A project in the WA wheatbelt (where sandy soils are at high risk) is showing that 50% of soils tested have subsoil acidification problems, around 40% of broadacre croppers in WA are liming, but lime use is less than half the amount required to manage soil acidification (Gazey *et al.* 2012; Chris Gazey, DAFWA, pers. comm.) For the dairy industry the results are similar, except that liming has decreased in Tasmania and WA since 2000/01 (Barson *et al.* 2012a). In horticulture there was little change in the percentage of farmer's liming between 1995/96 and 2007/08 (Barson *et al.* 2012c). In the grazing industries the percentage of beef cattle/ sheep businesses (outside the rangelands) liming declined between 2007/08 and 2009/10 (Barson *et al.* 2011).

Add lime at high rates, sufficient to reverse acidification in soils that have already acidified

The target values required to arrest acidification are generally high and followed by lower maintenance levels (Li *et al.* 2010). National lime use estimates from the Australian Bureau of Statistics' Agricultural Resource Management Survey show that a total of 4,136,312 tonnes of lime and 302,333 tonnes of dolomite were used in the broadacre cropping, dairy, horticulture and more intensively managed beef cattle/ sheep grazing industries in 2007-08 (Michele Barson, DAFF, pers. comm.) This is

considerably less than the projected requirement for nine million tonnes nationally (Webb *et al.* in preparation).

It is highly likely that these estimated lime requirements reflect the response of the more recalcitrant soils in south western Australia in broadacre and dairy industries where field studies indicate that it may take in excess of 11 years (and likely much more) and between 12–21 t/ha lime to raise the pH_{Ca} to 5.5 (Bolland and Russell 2010).

Use acid-tolerant plant species where available

There is good information available about the natural acid tolerance (and associated Al and Mn tolerance) of a range of pasture and crop plants (Slattery *et al.* 1989; Duncan 1999). The DAFWA Farmnotes soil acidity series (DAFWA 2012) also contains this information. No information was available on the combined use of this acid tolerant species and liming but it could be assumed that both practices are used in many regions that are at high risk of acidifying.

5.4 Concluding remarks

There is compelling evidence to show that liming surface soils increases yields of a wide variety of grasses and legumes. This is based on intensive R&D effort in the 80s-90s on long-term trials in the high rainfall and temperate zones of southern Australia, and more recently in the 1990s-2000s in southern WA field trials. Examples of information packages available are the Department of Agriculture, and Food Western Australia soil acidity series (DAFWA 2012) covering issues such as lime storage, liming rates and quality and expected and actual yield responses. For broadacre cropping and high return industries such as horticulture and dairy, liming can be an effective and profitable management strategy for mitigating surface soil acidification provided appropriate rates are applied that account for regional and local (management) factors of soil and plant type and N-fertiliser regimes.

The efficacy of practices to reduce subsoil acidification is less well established and only demonstrated on a small subset of soil types, but according to Anna Roberts (pers. comm.) the principles are simple – “it is about pH gradient, soil type and rainfall and therefore could be relatively easily calculated”. Notwithstanding the extended time frame for change and the high rates required to shift pH in some soils (of heavier texture) this is a remaining challenge for achieving improvements in soil pH condition. Once subsoil pH testing is adopted more broadly, the mitigation of subsoil acidity with more appropriate lime application rates and frequencies can be implemented in the high-risk agricultural regions.

Box 5.1: Managing Soil pH through a systems approach

System goal

To increase soil pH or slow its decline by managing nitrogen in plant systems.

Considerations

1. Reduce NO_3 availability by using legumes, NH_4 and organic forms of N fertiliser, and maximising N uptake by crops and pastures.
2. Reduce NO_3 leaching by maintaining drier soils and reduced fallow lengths (perennials and higher crop frequency).
3. Balance anion removal in products by liming, presumably this is forever.

Acidification is a constraint to production and C storage, there is reluctance by growers to use more lime and lime application for many farmers is driven by rules of thumb.

These responses are consistent with the soil C responses, provided lime application can be incorporated.

Recommended practices

Apply lime effectively, use organic and NH_4 fertilisers, use more legumes, perennials and increased crop frequency, test soils regularly where $\text{pH} < 6$.

Performance indicators

Trends in soil pH (relevant to support decisions at local to national and international scales), productivity (relevant locally to nationally), leaching of nitrates to subsoil and waterways (relevant locally and regionally).

Conflicts

Suitable machinery for applying lime, especially at depth, higher management inputs required to apply lime at sufficient quantities in some areas and the costs of these inputs encourage some farmers to increase cropping and grazing pressure to maintain cash flow.

6. Wind erosion

6.1 Nature of the issues

Soil erosion is the removal of soil particles from the ground's surface. It is usually brought about by wind and/ or water. The extent to which soils are susceptible to wind erosion depends on a range of factors, including climatic variability, ground cover, topography, the nature and condition of the soil, and the energy of the wind.

Soil particles behave differently depending on the strength of the wind and how well the soil surface is protected by ground cover. As wind erosion intensifies, aggregates can break or abrade, releasing dust into the air (Leys *et al.* 2010). Land management can either moderate or accelerate wind erosion rates, largely depending on how it affects the proportion of bare soil, the dryness and looseness of the ground's surface, and structures that reduce the force of wind (i.e., windbreaks). Grazing by stock, native animals (e.g., kangaroos) and feral animals (rabbits, camels, horses, goats) have major impacts on ground cover and soil physical properties. Such impacts have been exacerbated by the establishment of watering points that allow these animals to be active throughout previously dry landscapes in many parts of Australia (James *et al.* 1999; Landsberg *et al.* 2002). The changes in land cover brought about to establish much of Australia's agriculture have led to an acceleration of wind (and water) erosion (Beadle 1948; Yapp *et al.* 1992; Edwards and Pimentel 1993; Ludwig and Tongway 1995; Wasson *et al.* 1996; Campbell 2008; Hairsine *et al.* 2008; Leys *et al.* 2009).

The on-site impacts of wind erosion include soil loss, reduction in soil nutrients and organic matter (including soil organisms), release of soil carbon to atmosphere, undesirable changes in soil structure, reduced water infiltration and moisture-holding capacity, and exposure of unproductive saline and acid subsoils (Morin and Van Winkel 1996; Belnap and Gillette 1998; Pimentel and Kounang 1998; Lal 2001; Leys *et al.* 2009; McAlpine and Wotton 2009). Off-site impacts include negative impacts on the global climate through positive radiative forcing of dust, physical impacts of dust storms on buildings and equipment, and health impacts of dust for people (Leys *et al.* 2009). The limited data available suggest that the off-site costs of wind erosion can be many times greater than the on-site costs. Williams and Young (1999) estimated direct market values for on-site costs of wind erosion in South Australia to be \$1-6 million per year, compared with an estimated \$11-56 million cost per year for off-site costs (largely associated with human health). The costs borne by Sydney when hit by the 'Red Dawn' dust storm in 2009, including costs associated with

cleaning premises and cars, disruptions to transport and construction, and absenteeism were estimated to be \$330.8 million, while losses of soil fertiliser and carbon to landholders were estimated at \$9 million (Tozer 2012). On the other hand, transport of eroded soil can provide important inputs to nutrient budgets of systems that can trap dust, such as forests and woodlands (McTainsh and Strong 2007).

Several major initiatives have been put in place to improve Australia's ability to monitor wind erosion and to identify priority areas for remedial action (Leys *et al.* 2010; McTainsh *et al.* 2012; Smith and Leys 2009). This will be especially important in the future as climate change is likely to increase the likelihood of soil erosion, due to increased incidence of droughts and reductions in crop production and ground-cover (Leys *et al.* 2009; Soils Research Development and Extension Working Group 2011). Historically, wind erosion has been particularly active in times of drought. In the 1940s and again in 2002 and 2009 there were heightened concerns due to dust storms hitting major Australian towns and cities (McTainsh *et al.* 1990; McTainsh *et al.* 2011). Wind erosion appears to have been reduced substantially since the 1940s, primarily due to better management of vegetation cover on agricultural lands (Australian State of the Environment Committee 2011), but it is expected that the incidence of huge dust storms, like those in 2002, will increase in the future (Leys *et al.* 2009).

6.2 Land management practices in relation to wind erosion

Approaches to reducing wind erosion address three major aspects (Carter 2006):

- Ground cover
- Soil looseness
- Wind velocity

Ground cover is important as it reduces wind speed at the soil surface and captures soils particles mobilised by wind. Soil looseness increases when there is too little vegetation cover, soils are dry, the type of soil contains small particles and/ or the surface is smooth. Maintaining soil moisture, avoiding trampling of exposed or susceptible soil by stock and maintaining rough soil surface are all ways to reduce soil looseness (Findlater *et al.* 1990; Carter *et al.* 1993; Moore *et al.* 2001; Carter 2002; 2006; McTainsh *et al.* 2011). While the velocity of wind is determined by the weather, it can be moderated locally by creating windbreaks.

Cropping and mixed farming

Recent surveys of past soil erosion, using measurement of $^{137}\text{Caesium}$ in soils, have concluded that levels of combined water and wind erosion from cultivated land and rangelands are relatively similar, and as much as eight times greater than from uncultivated areas and forests (Loughran *et al.* 2004; Bui *et al.* 2010). Regions with the largest impacts of wind erosion tend to be focused in arid and semi-arid rangelands of south-western Queensland, western NSW, north-central and north-eastern South Australia and western Western Australia, posing particular challenges for grazing enterprises (Leys *et al.* 2010). The semi-arid agricultural lands of eastern West Australia also have areas of high and very high wind erosion, compared with the generally low erosion levels in the non-agricultural lands of western South Australia, the northern Northern Territory and eastern Western Australia (Leys *et al.* 2010).

The process of cultivation of soil is a key factor affecting potential for both wind and water erosion in broadacre cropping (Freebairn 1992a; b; Freebairn and Loch 1993; Moran 1998; Barson and Lesslie 2004). The effects of cultivation have been likened to a fire passing through ploughed soil, disrupting the activities of soil organisms, oxidising organic matter, reducing soil fertility and often leading to soil structural problems (Australian State of the Environment Committee 2011). Some of these effects can be offset by addition of fertilisers and organic matter, but structural problems are much harder to address. The combination of soil type, moisture, tillage practice, and associated activities like clearing of deep rooted perennials, burning of crop residues, and running of grazing animals on the land can lead to the sorts of structural changes that encourage bare soil (Bartley *et al.* 2006).

The types of land management recommended to reduce wind erosion in cropping and mixed farming zones (McTainsh *et al.* 2011) include:

- Maintenance of adequate plant residue cover for soil erosion protection through the adoption of stubble retention systems;
- The adoption of minimum/ zero tillage systems that protect against erosion and maintain or improve soil structure;
- Avoidance of cultivation in high erosion risk periods;
- Reduction in burning stubbles;
- Use of chemical fallowing rather than tillage;
- Integrated feral fauna and flora control programs, including biological controls;
- Fencing to land class through a developed farm plan;

- Retention of boundary tall perennial vegetation;
- Avoiding grazing erosion-prone areas by fencing these areas;
- Intensive strip grazing/ cropping;
- Land reclamation of degraded areas for both production and conservation uses;
- Involvement of agricultural commodity industries in promotion of better land management practices.

Grazing/ pastoral enterprises

Livestock grazing has been associated with a decline in native perennial cover and an increase in exotic annual cover, reduced litter cover, reduced soil cryptogam cover, loss of surface soil microtopography, increased erosion, changes in the concentrations of soil nutrients, degradation of surface soil structure, and changes in near ground and soil microclimate (Eldridge 1998; Evans 1998; Yates *et al.* 2000; Jansen and Robertson 2001; Landsberg *et al.* 2002; Sparrow *et al.* 2003; Dorrough *et al.* 2004; Hunt *et al.* 2007; Department of the Environment 2009). Recommendations for countering the effects of grazing on soil erosion involve reducing grazing pressure, keeping animals away from riparian areas, and managing movements of cattle using watering points (Andrew 1988; James *et al.* 1999; Dorrough *et al.* 2004; Hunt *et al.* 2007; McTainsh *et al.* 2011). Rotational grazing and cell grazing have been shown to be profitable approaches to managing the impact of grazing on pastures and, therefore, ground cover (McCosker 2000; Southorn and Cattle 2004a; Crosthwaite *et al.* 2008). McTainsh *et al.* (2011) note that pastoral industries have improved in a variety of ways since the 1940s, including better control of total grazing pressure (native, feral and domestic stock).


6.3 Evidence of the effectiveness of management practices for reducing wind erosion

Evidence for the effectiveness of measures to reduce wind erosion come from two types of studies: experimental studies showing relationships between soil movement, wind speed and the state of the soil surface; and evidence of reduced incidence of dust storms as land management practices have improved from the 1940s to the present.

Numerous studies have been performed in Australia, and in comparable ecosystems in other parts of the world, to show that increasing ground cover reduces losses of soil due to both wind and water erosion (Eldridge 1993; Eldridge and Greene 1994;

Erskine and Saynor 1996; Scanlan *et al.* 1996; Carroll *et al.* 2000; Loch 2000; Yates *et al.* 2000; Eldridge and Leys 2003; Durán Zuazo *et al.* 2004; Heywood 2004; Greenway 2005; Bartley *et al.* 2006; Durán Zuazo *et al.* 2006; Raya *et al.* 2006; Silburn *et al.* 2011). Increasingly, evidence is being documented from on-ground initiatives by individual land managers (Jenkins and Alt 2007; Jenkins and Alt 2009).

In semi-arid environments, it has been concluded that ground cover of around 50% is required to keep wind erosion to a minimum (Findlater *et al.* 1990; Leys 1992; Rosewell 1993; Scanlan *et al.* 1996; Leys 1998; Loch 2000; Leys *et al.* 2009; Silburn *et al.* 2011) (Figure 6.1).



The general relationships between ground cover and soil erosion have been known for over 20 years. The main focus of research and development during the past two decades has been on how to achieve ground cover cost-effectively. This is discussed in the following section on water erosion.

The second line of evidence for the effectiveness of better land management (ultimately resulting in improved ground cover) for reducing wind erosion comes from comparisons of Dust Storm Indices (DSI) between the 1940s and the present (McTainsh *et al.* 2011). DSI provides a measure of the frequency and intensity of wind erosion activity. McTainsh *et al.* (2011) showed that mean on-site wind erosion in the 1940s was almost 6 times higher than in the 2000s, and the mean maximum DSI for the 1940s was 4 times that of the 2000s. There are also significant regional differences: wind erosion in the 1940s was much more active in the Mulga, Riverina and Central Australia than in the SA and WA rangelands, and the decrease in wind

erosion between then and the 2000s was much more pronounced in the east and centre of the continent (McTainsh *et al.* 2011). Uptake of measures to improve ground cover was discussed in Section 4 and is also considered in Section 7. Although there have been high rates of adoption among farmers (D'Emden and Llewellyn 2006; Llewellyn and D'Emden 2009; Llewellyn *et al.* 2012), it has not been complete, and so risks of both wind and water erosion remain high in some areas (McTainsh *et al.* 2011).

Box 6.1: Managing wind erosion through a systems approach

System goal

To reduce soil loss from wind erosion.

Considerations

1. Wind speed is reduced by high cover (from soil C actions) and tree windbreaks (probably down fence-lines for operational efficiency). Maintaining ground cover of at least 50% will reduce the risk of soil loss through wind erosion.
2. Particle availability is reduced by limiting concentrated stock movements and tractor operations on very dry surface soils which can generate clay sized particles.

Recommended practices

As for soil C, acidification and water erosion practices.

Performance indicators

Dust monitoring (DEHNSW 2012).

Conflicts

In many cases major changes are needed from traditional practices to ones that build and maintain high levels of ground cover in all seasons and in wet and dry years.

7. Water erosion

7.1 Nature of the issues

Water erosion of soils occurs when soil particles are detached and carried away by water flowing across a landscape. In some cases soil loss is uniform (sheet erosion). In other cases small channels are formed (rill erosion). When the velocity and volume of water are high enough, and the soil surface is vulnerable, deep channels can be cut (gully erosion). Tunnel erosion occurs when the subsoil is removed while the surface soil remains relatively intact, producing tunnels under the soil, which eventually cause the surface to collapse (Coles and Moore 2001).

Like wind erosion (Section 6), the on-site impacts of water erosion include soil loss, reduction in soil nutrients and organic matter (including soil organisms), release of soil carbon to the atmosphere, undesirable changes in soil structure, reduced water infiltration and moisture-holding capacity, and exposure of unproductive saline and acid subsoils (Morin and Van Winkel 1996; Belnap and Gillette 1998; Pimentel and Kounang 1998; Lal 2001; Leys *et al.* 2009; McAlpine and Wotton 2009). Off-site impacts include sedimentation of waterways and impacts on quality of surface water and groundwater (turbidity, nutrient and other chemical loads).

Erosion from hillslopes by water is complex and multifaceted (Figure 7.1). It is determined by the combined effects of:

- the strength of water flow (influenced by the amount and rate of rainfall, the length and steepness of slopes, the degree to which the energy of raindrops is dissipated by ground cover, and whether the water encounters obstacles to its flow)
- the predisposition of soil particles to be dislodged (affected by soil type, ground cover, structural properties of the soil that affect the infiltration rate of water, and the soil's moisture), and
- the presence of obstacles to the flow of sediment from a site (e.g., its roughness and the presence of obstacles such as fallen timber, plant stems or contour banks created to limit erosion).




Figure 7.1: Factors influencing soil erosion by water. Figure was derived from various publications cited in the text

By far the strongest factor mitigating water erosion is ground cover: typically, 20-30% cover reduces erosion by 80-90% across a range of soils and land uses (Freebairn *et al.* 1986; Freebairn and Wockner 1986; Freebairn 1992b; Littleboy *et al.* 1992; Freebairn *et al.* 1993; Freebairn 2004; Gerik and Freebairn 2004; Silburn *et al.* 2007; Freebairn *et al.* 2009). Ground cover can be grasses, herbs, trees, dead plants with root systems still intact, dead plant material (especially branches) lying on the surface, or even stones. The mechanisms by which ground covers prevent erosion are a combination of physical binding (by roots), slowing of over-land flows (by plants, fallen timber, litter, and stones as physical barriers) and dissipation of the energy of raindrops (by foliage) (Freebairn and Wockner 1986; Brandt 1988; Hall and Calder 1993; Daily *et al.* 1997; Loch 2000; Phillips *et al.* 2000; Freebairn *et al.* 2009; McAlpine and Wotton 2009).

It is estimated that current rates of soil erosion by water across much of Australia exceed soil formation rates by a factor of at least several hundred and, in some areas, several thousand (Australian State of the Environment Committee 2011). As a result, the expected half-life of soils (the time for half the soil to be eroded) in some upland areas used for agriculture ranges from less than a century to several hundred years. While the time for total loss of soil is estimated to range from 100-500 or more years in different parts of Australia, it is expected that crops and other plants will respond to small changes in depth of topsoil, so that many areas are at risk of critical decline in productivity in much less than 100 years (Bui *et al.* 2010). Areas at highest

risk include Coastal Queensland, the Wet Tropics, Mitchell Plains grasslands, New England Tablelands, and Victoria River basin in the NT. The 2011 State of the Environment Report concluded that in 9 of Australia's 22 physiographic provinces, the majority of the landscapes have been eroded (by combined wind and water erosion) to the extent that plant growth and agricultural yields have been adversely affected (Australian State of the Environment Committee 2011). In the other 13, it was concluded that management and monitoring are needed or the system of land use will be threatened in the long term.

Drought predisposes land systems to erosion by both wind and water because of reduced soil cover. Major soil erosion accompanied the intense rainfall events and floods that broke the drought of the late 2000s in southern Queensland (Australian State of the Environment Committee 2011).

7.2 Land management practices in relation to water erosion

Land uses that affect water erosion do so primarily via their effects on ground cover, evaporation of soil moisture, soil structure, compaction by heavy equipment or running of stock, and creation of contours that control water flow (Australian State of the Environment Committee 2011).

Broadacre cropping

Many of the effects of cultivation on susceptibility to wind erosion (Section 6) also apply to water erosion. Water erosion associated with cropping was recognised as a serious issue in the 1930s (Carey *et al.* 2004). Different studies report sediment yields from cultivated basins of between 2 and 21 times those from undisturbed native forests (Neil and Galloway 1989; Neil and Fogarty 1991; Erskine *et al.* 2002), although it should be noted that good land management can keep these figures within the low end of this range (Erskine *et al.* 2002). Soil conservation structures (contour banks and grassed waterways) were designed to reduce the slope length and thus net water erosion. These have been implemented extensively in Australia, but have not been sufficient to bring soil erosion within acceptable limits (Freebairn *et al.* 1993; Freebairn *et al.* 2009).

Management of water erosion on cropping lands has increasingly focused on methods of planting and managing crops and controlling weeds that involve little or no tillage, retention of stubble after harvesting, inclusion of a pasture phase between crops and minimisation of the effects of machinery by controlled traffic methodologies (Freebairn *et al.* 1993; Freebairn 2004; Li *et al.* 2007; Silburn *et al.* 2007; Llewellyn and D'Emden 2009; Llewellyn *et al.* 2012). Creating raised beds for

crops in waterlogged areas can create an erosion hazard unless slopes and ground cover are managed carefully (Hamilton *et al.* 2005; Wightman *et al.* 2005)

Over the last 20 years new tillage practices have been developed that maximize water infiltration and reduce runoff; new row spacing and plant arrangement schemes have been developed to reduce soil temperatures and soil evaporation losses. Crop modelling and weather prediction capabilities have been developed to advise farmers on the opportune time of sowing that ensures adequate supply of stored soil water in combination with sufficiently high growing season rainfall probability required to satisfy the crop growth requirements and the farmers' yield goal (Gerik and Freebairn 2004; Australian State of the Environment Committee 2011; McTainsh *et al.* 2011). While including a pasture phase between crops is considered advantageous in managing ground cover, the potential effects of stock on the soil surface during this phase can potentially pose similar problems to those faced on dairy farms, especially if soils are wet (see below).

The uptake of minimum tillage approaches has required two major innovations: equipment capable of planting in stubble; and effective methods for weed control without disturbing the soil (Freebairn 1992; Freebairn and Loch 1993). The advent of better ways to manage heavy vehicles (controlled traffic) has also contributed to reducing runoff-driven erosion (Li *et al.* 2007).

Horticulture

As a form of cropping, horticulture faces many of the same risks as broadacre cropping in terms of encouraging soil erosion. The hardening of soils in many orchards (coalescence) restricts the growth and function of tree roots and infiltration of water to roots (Cockcroft 2012). Two key management innovations in orchards have been control of machinery traffic to minimise soil compaction, and establishment of ground cover plants that both minimise erosion and contribute to the soil ecosystem (Wells and Chan 1996; Dewhurst and Lindsay 1999; Firth *et al.* 1999; Zwieten *et al.* 2001; Reid 2002; McPhee 2009; Loch 2010; Slavich and Cox 2010; HAL 2012a). Increased ground cover is correlated with higher diversity of soil organisms, which has been found to have beneficial effects on water infiltration (and therefore reduced run-off erosion) promotes natural pest control (Colloff *et al.* 2003; Colloff *et al.* 2010).

Dairy

Many dairy farms combine the running of dairy cattle with beef cattle, cropping and/or irrigated pasture production (Ashwood *et al.* 1993). To maintain high production of milk, pastures are fertilized. Key challenges for such enterprises include controlling

sediment (along with nitrogen and phosphorus) losses into waterways, which can be exacerbated by compaction and disturbance of soil by the feet of grazing animals (Nash and Murdoch 1997; Fleming 1998; Fleming and Cox 2001; Fleming *et al.* 2001; Aarons *et al.* 2004; Nash *et al.* 2005; Barlow *et al.* 2007; Chan 2007).

Irrigation itself has the capacity to increase soil erosion by accelerating mineral weathering, transporting and leaching soluble and colloidal material, changing soil structure, and raising the local water table, thereby increasing the risk of salinity (Heywood 2004; Jenkins and Alt 2007; Jenkins and Alt 2009). Irrigation also has the capacity to reverse soil preparation measures such as the tillage that precedes planting.

Grazing

Livestock grazing is the most widespread Australian land use (Section 4). Impacts of livestock grazing on ground cover were discussed in Section 6. These impacts affect vulnerability of landscapes to both water and wind erosion. In addition, as discussed above, grazing during a pasture phase between cropping could increase vulnerability of soils to water erosion by disrupting soil structure and reducing ground cover.

7.3 Evidence of the effectiveness of management practices for reducing water erosion

As mentioned in Section 6, there is an extensive literature showing that increasing ground cover reduces losses of soil due to both wind and water erosion (Eldridge 1993; Eldridge and Greene 1994; Erskine and Saynor 1996; Scanlan *et al.* 1996; Carroll *et al.* 2000; Loch 2000; Yates *et al.* 2000; Eldridge and Leys 2003; Durán Zuazo *et al.* 2004; Heywood 2004; Greenway 2005; Bartley *et al.* 2006; Durán Zuazo *et al.* 2006; Raya *et al.* 2006; Jenkins and Alt 2007; Jenkins and Alt 2009; Silburn *et al.* 2011). Box 7.1 gives an example of how ground cover management, climatic variability and economic pressures can interact to force a region into an 'erosion trap'.


Like wind erosion (Section 6) there is a small number of studies that have focussed on the minimum extent of ground cover needed to avoid soil erosion. While different combinations of cover-types have different effectiveness, largely depending on the proportion and pattern of bare ground (Greene *et al.* 1994; Ludwig *et al.* 2005), some broad guidelines about effective cover have been developed. In general, a higher proportion of cover (70% - Figure 7.2) is recommended to manage water erosion than for wind erosion (50% - Figure 6.1) (Findlater *et al.* 1990; Rosewell 1993; Scanlan *et al.* 1996; Loch 2000; Silburn *et al.* 2011). For environments where rainfall

is moderate to high, and/ or slopes are steep, 80-100% ground cover is recommended (Leys 1992; Lang and McDonald 2005). The standard of 70% is being applied widely by catchment management authorities in northern NSW (Central West Catchment Management Authority 2008; Namoi Catchment Management Authority 2010).

Box 7.1: The Gascoyne Catchment – A Case Study of Water Erosion

Three record flooding events in the Gascoyne Catchment, Western Australia, in the summer of 2010–11, resulted in massive plumes of soil spreading into the ocean at the mouth of the Gascoyne River (Waddell *et al.* 2012). The amount of soil lost during one of the flooding events was an estimated 2,250,000 tonnes. Restoration of damaged land in the Carnarvon area after the three floods required 140,000 tonnes of topsoil. It was concluded that the poor state of the landscapes in the catchment resulted in very much higher losses of soil than would have occurred in a catchment with good ground cover, although the extent of the additional losses could not be determined. The flooding also resulted in damage to infrastructure in the Carnarvon horticulture area.

The Gascoyne Catchment is in a typical ‘erosion trap’. Some of the higher country is protected from erosion by a covering of stones, but other parts have been heavily grazed and are highly degraded. This results in the rapid transfer of sediments and large amounts of water into the lower parts of the catchment. Downslope of the upland areas the landscape is dominated by extensive sheet wash plains. These areas are sources of browse for stock and have been over-utilized, leading to soil instability, when water flows from the upland areas, disrupted water flows and nutrient cycles, and erosion where stock have disrupted the soil surface. As the catchment goes through dry periods, grazing pressure in this part of the catchment increases, making erosion risks worse. In the catchment’s lower reaches, saline alluvial plains are stabilised to some extent by buffel grass, but this is susceptible to fire, the risk of which increases in dry periods. As recovery of these sorts of systems is slow, the challenge of returning this catchment to a state that is resilient to the effects of water in the landscapes, and to climate variations in general, is major.



The main focus of research and development during the past two decades has been on how to achieve appropriate proportions of ground cover cost-effectively. In grazing systems, removal of stock has been shown to allow recovery of ground cover, if conditions are favourable for regrowth of pastures, but recovery of full soil functionality, especially organic matter content, can take years to decades (Braunack and Walker 1985; Basher and Lynn 1996; Lal 1999; Silver *et al.* 2000) and the short-term and longer-term reduction in financial returns can be a disincentive for graziers (Lilley and Moore 2009). Maintaining a diversity of species, especially native plants and soil organisms, at landscape scales, is argued to be an important component of ground cover strategies in grazing systems, as this provides ready sources of species to re-establish ground cover communities after disturbances such as fires and drought (McIntyre 2002; Colloff *et al.* 2010). Restoring and maintaining plant species diversity and community structure is likely to provide greater resilience of ground cover to climatic and other shocks. This will probably require strategies that capture resources, such as water, seeds, nutrients and carbon, increase their retention on-site, and improve microclimate, in addition to removing stock (Yates *et al.* 2000).

Across Australian states, 30-80% of horticultural businesses reported using alternative or cover crops between main crops or using mulching and/ or matting to provide ground cover between crops in 2009-10 (Barson *et al.* 2012c). The proportion of grazing (beef cattle/ sheep) businesses across Australia monitoring ground cover levels has increased from 70% in 2007-08 to 79% in 2009-10, but the

percentage of businesses setting ground cover targets decreased from 40 to 31% in the same period (Barson *et al.* 2011). Similar trends were seen for dairy businesses (Barson *et al.* 2012a).

Detailed research on reduced-tillage approaches has been conducted across Australia (Hamblin *et al.* 1982; Hamblin 1984; Freebairn *et al.* 1986; Hamblin *et al.* 1987; White 1990a; Buckerfield 1992; Freebairn 1992; Kingwell *et al.* 1993; Schmidt and Belford 1993; Schmidt *et al.* 1994; Felton *et al.* 1995; Thomas *et al.* 2007). Conservation tillage has been shown to dramatically reduce soil erosion and provide benefits for production in most areas (Freebairn *et al.* 1986; Freebairn 1992; Radford *et al.* 1993; Thomas *et al.* 2007). No-tillage and reduced tillage (stubble mulch) practices with stubble retention have generally resulted in greater fallow efficiency (gain in soil water during the fallow per unit of rainfall), soil water storage and grain yield, compared with conventional tillage practices, which incorporated stubble into the soil, although lower grain protein content has also been reported for some locations (Freebairn 1992; Radford *et al.* 1993).

These results are supported by around 20 commercial-scale, development and extension experiments across a range of crops and environments in the grain growing areas of Queensland since the 1970s, in which mean grain yield was 9% greater under no-tillage than with stubble incorporation (Thomas *et al.* 2007). There is some evidence that yield responses are likely to be greater where soil water supply limits yield (Freebairn *et al.* 1986; Thomas *et al.* 2007). While it is likely that these general trends will apply in other places with different soil types and production systems, the researchers caution against uncritical generalization without further experimentation (Freebairn *et al.* 2009).

Case studies in Queensland indicate that these benefits can be turned into significantly improved profits from no-tillage compared with traditional tillage, especially when economies of scale can be achieved by applying the same labour and machinery over large areas, and when controlled traffic management is used (Wylie 1997; Gaffney and Wilson 2003).

Some limitations of conservation tillage have been identified. The reduced surface roughness produced by no-till management can lead to enhanced run-off and sediment movement in areas where maintaining high biomass of plants is difficult, or where low cover results from crop failure or grazing (Freebairn *et al.* 2009). In these cases, some tillage might be required to create surface roughness. Since one role of tillage is weed and disease control, crop rotation and other approaches to weed control, such as inversion ploughing every 8-10 years to bury weed-seeds, are

especially important in no-till systems (Douglas and Peltzer 2004; Thomas *et al.* 2007).

As discussed in Sections 4 and 5, the adoption of some form of minimum tillage has increased over the past two decades.

In southern Australia, key factors that have influenced adoption of minimum tillage approaches include machinery costs, perceived lack of convincing evidence of results, and concerns about herbicide resistance and weed control (D'Emden and Llewellyn 2006; Llewellyn and D'Emden 2009; 2010; Llewellyn *et al.* 2012). The main reasons given by adopters for limiting their use of no-tillage approaches include herbicide resistance, weed control issues, soil physical constraints, pests and soil disease. Adoption of no-tillage approaches appears to be leveling out at about 90% of farmers in many regions of Australia (Llewellyn *et al.* 2012).

Box 7.2: Managing water erosion through a systems approach

System goal

To reduce water erosion by reducing suspended sediment and transported sediment.

Considerations

1. Maintain ground cover at better than 50% to reduce raindrop impact and production of suspended sediments. Maintaining good ground cover will also increase biomass available for soil carbon.
2. Increase infiltration (reduce runoff) with adequate ground cover, manage soil moisture to avoid excessive decomposition and waterlogging (as for carbon management), and reduce compaction by using Controlled Traffic (CT) approaches.
3. Where appropriate, manage runoff with designed layouts (controlled traffic farming, diversion and contour banks) to prevent flow concentration (spread runoff evenly across the land). Runoff velocity is then unlikely to reach erosive levels in our landscapes. CT wheel tracks are designed to carry runoff to safe disposal areas (typically diversion channels).

Recommended practices

Soil C and acidification practices, controlled traffic and designed layouts, ground cover management.

Performance indicators

Water erosion control (especially percentage groundcover, turbidity of off-flows, water quality) (relevant at local to regional scales), access and timeliness (relevant at farm scale).

Conflicts

In many cases major changes are needed from traditional practices to ones that build and maintain high levels of ground cover in all seasons and in wet and dry years.

8. Ecosystem services and resilience of soils

8.1 The concept of ecosystem services

The concept of ecosystem services evolved to bridge the perceived gap between economics and ecology. To achieve this it has been necessary to consider at some length how to define and classify ecosystem services so that they not only make sense to a range of stakeholders, but also can be used unambiguously in economic valuation and environmental accounting. Because this process has involved multiple disciplines, there have been different views on how to define terms like ‘processes’, ‘functions’, ‘services’, and ‘value’ (Costanza *et al.* 1997; Daily 1997; de Groot *et al.* 2002; MA 2005; Wallace 2007; Costanza 2008; Fisher *et al.* 2009; TEEB 2009; Dominati *et al.* 2010; Maynard *et al.* 2010; UK National Ecosystem Assessment 2011b; Nahlik *et al.* 2012; Robinson *et al.* 2012). Typologies of ecosystem services have remained fluid with the recognition that services must be identified in relation to those receiving the services, and that this relationship differs with different groups of people, different places and different purposes for considering ecosystem services (de Groot *et al.* 2002; Costanza 2008; Fisher *et al.* 2009).


As our focus in this report is on the links between land management, soil condition and benefits to humans, we have adapted four recent approaches for conceptualising these relationships into the framework shown in Figure 8.1.

Figure 8.1 incorporates several recent conventions designed to reduce inconsistency of terminology and ensure that the direct and indirect contributions of ecosystems are not confused in economic evaluations and environmental accounting:

- Ecosystem services are defined and described (Table 8.1) in terms of what possibilities soil ecosystems make available to humans, without the need for intervention by humans¹; the benefits to humans are identified separately, and require actions or the articulation of needs by humans (Boyd and Banzhaf 2007; Fisher *et al.* 2009; Haines-Young and Potschin 2009).
- We have avoided distinguishing between ecosystem processes and functions, referring only to processes. Ecosystem processes are defined as transformations of inputs into outputs and ecosystem services are defined as the flows that arise from these processes and are of benefit to humans (Dominati *et al.* 2010).

¹ There are several published definitions that meet these criteria, for example “the aspects of ecosystems utilized (actively or passively) to produce human well-being” (Fisher *et al.* 2009)

- We have distinguished between final ecosystem services (those that can be turned directly into benefits by humans) and intermediate ecosystem services (those that support other services but are not used directly for benefit by humans) (de Groot *et al.* 2002; Boyd and Banzhaf 2007; Fisher *et al.* 2009; TEEB 2009; Bennett *et al.* 2010; Dominati *et al.* 2010; Johnston and Russell 2011; UK National Ecosystem Assessment 2011b).
- For consistency with other typologies, we have adopted the broad organising headings of ‘provisioning’, ‘regulating, and ‘cultural’ services (Daily 1999; MA 2005; De Groot *et al.* 2010; Dominati *et al.* 2010).



Although it is potentially confusing to distinguish between final and intermediate ecosystem services, we agree with advocates of this approach that: (i) being strict about final services is essential to avoid double counting of benefits in economic assessments, such as we perform in this report; and (ii) there is a need to recognise a level of aggregation of processes above that of nutrient, water and carbon cycling and the like, by which soils support the final services produced by broader ecosystems.

8.2 Relating soil ecosystem processes to services and benefits

The roles of soils in supporting natural and agricultural ecosystems have been recognised for some time and their importance for providing ecosystem services has been discussed in various recent syntheses (Daily *et al.* 1997; Wall and Virginia 2000; Balmford *et al.* 2002; De Groot *et al.* 2003; Swinton *et al.* 2006b; Dale and Polasky 2007; Kroeger and Casey 2007; Swinton *et al.* 2007b; Turner and Daily 2007; Weber 2007; Bennett *et al.* 2010; Robinson *et al.* 2012). Figure 8.2 and Table 8.1 draw on a number of these syntheses.




Figure 8.2: Interrelationships between living and non-living components of soils, major processes, ecosystem services, benefits to humans and who the beneficiaries are

The diagram synthesises frameworks by: Palm *et al.* (2007); Kibblewhite *et al.* (2008a); Bennett *et al.* (2010); Dominati *et al.* (2010); UK National Ecosystem Assessment (2011a)

Figure 8.2 and Table 8.1 illustrate the complex interrelationships between the living and non-living components of soil, the processes and ecosystem services these interactions generate and the benefits derived by a range of beneficiaries, and seek to simplify this complexity by identifying a relatively small number of ‘final’ ecosystem services and benefits. This figure also emphasises the underpinning importance of soil’s natural capital (including both living and non-living components), which is the key to long-term sustainable management of soils, and maintenance of soil resilience (Lal 1997; Dominati *et al.* 2010; Sylvain and Wall 2011; Robinson *et al.* 2012).

Table 8.1: Description of the broad groups of ecosystem services provided by soils*

Ecosystem services	Description of services and benefits
Provisioning services Provision of fertile soil, natural products and clean water Support for native vegetation Maintenance of genetic diversity Support for structures	<p>Provision ecosystem services are those that either directly provide products that people value or can be used to produce things of value.</p> <p>Products from soils include clean water, bush foods (e.g., witchety grubs, mushrooms), timber, and chemicals and genetic material that might be developed as pharmaceuticals or used in genetic and other technologies in the future.</p> <p>Fertile soil can be used by humans to grow crops. Soil fertility is maintained by a range of processes, including nutrient cycling (distribution of carbon, nitrogen and phosphorus throughout soils by a range of soil organisms), gaseous exchange with the atmosphere (extraction and release of nitrogen and carbon), and the engineering activities of earthworms, insects, fungi and other species (which maintains soil structure, porosity and water-holding and infiltration capacities).</p> <p>By supporting the growth of native forests, woodland and grasslands, soils contribute to the ecosystem services that native vegetation provides, including the provision of fodder for stock.</p> <p>It is often overlooked that the formation of soil by natural processes provides to foundation for anchoring structures such as houses, other buildings and other infrastructure.</p>
Regulating services Water flow regulation Maintenance of landscape (soil) stability Regulation of atmospheric gases Regulation of weather and climate Remediation of wastes Regulation of species and populations Pollination	<p>Regulating ecosystem services are so named because they control biophysical processes in ways that can be beneficial to humans.</p> <p>The structural properties of soils, determined living and non-living components below ground and the vegetation component of the soil-plant ecosystem above ground, affect how water flows across the surface of the ground or infiltrates underground watertables. This affects erosion and damage to human life and property as well as the access of plants, including crops and native vegetation, to water. In Australia, regulation of watertables by soil-plant ecosystems is a key determinant of whether salinity (rising of salt to the surface) becomes a problem.</p> <p>The above processes stabilise landscapes and prevent negative health impacts and damage to property that can accompany dust storms (including major impacts of dust on weather patterns (Mahowald <i>et al.</i> 2010; Rotstayn <i>et al.</i> 2012)). Along with vegetation, soils affect the amount of radiation (heat, light) reflected from the earth to the atmosphere, which affects weather and climate. Evaporation of water into, via soil and vegetation also influences weather and climatic patterns.</p> <p>Extraction of carbon and nitrogen from the air by soils, and release of these elements into the air, are major mechanisms for regulating the composition of the atmosphere, effecting climate and suitability of air for humans.</p> <p>Soils breakdown organic and non-organic compounds, some of which can become toxic to humans, other animals, or plants. Additional investment in waste disposal is needed when this ecosystem service is exceeded by the rate of production of</p>

Ecosystem services	Description of services and benefits
	<p>wastes by humans.</p> <p>The various species living in soil interact with one another and with species living above ground, by eating one another and competing for food and space. In so doing, they regulate one another's numbers and prevent any species increasing to numbers that might be detrimental to ecosystem functions and/ or human activities. Some of these species also play a role in pollinating plants and moving seeds around in landscapes.</p>
<p>Cultural services</p> <p>Contributions to species, ecosystem and landscape diversity</p>	<p>It has been recognised for some time that people draw a wide range of inspiration and both physical and mental health benefits from ecosystems. People identify with certain landscapes ('sense of place'), gain knowledge by studying ecosystems, and often find spiritual connections with the land. In all of the ways discussed above, and more, soil contributes to the diversity and condition of landscapes. Although these cultural benefits are not always easy to define, they are nevertheless vital for humans to thrive mentally and physically.</p>

*Detailed discussions about the nature of ecosystem services in agricultural and other lands in Australia and globally can be found in the following references: Binning *et al.* (2001); de Groot *et al.* (2002); Haygarth and Ritz (2009); Bennett *et al.* (2010); UK National Ecosystem Assessment (2011a; b).

We have chosen to develop our own framework (Figure 8.2) as we have found existing ones to be inconsistent with regard to some of the principles listed in Section 8.1. The following examples illustrate some of these inconsistencies and explain why we have emphasised them in the context of this report:

- Some other frameworks include 'supporting services' as a separate category. In Figure 8.2, these are considered to be part of the 'major processes'.
- When considering 'provisioning services', several other frameworks for ecosystem services from soils and agricultural land include provision of marketable goods, including food (crops and/ or livestock), wood, fibre and others, as ecosystem services (Bennett *et al.* 2010; Dominati *et al.* 2010; UK National Ecosystem Assessment 2011b). Following the principle of separating the services that ecosystems provide from the benefits that are derived with human input (Boyd and Banzhaf 2007; Kroeger and Casey 2007) (see also Table 8.1), we consider that soil ecosystems provide fertile soil but not crops or livestock (Figure 8.2). We do, however, consider provision of edible products from native soil ecosystems (e.g., edible insects and fungi) to be an ecosystem service. This distinction is important because, if we are to assess the value of better management of soil ecosystems we need to be able to account separately for the human inputs and ecosystem responses.
- It is common in ecosystem service typologies to describe 'cultural services' in terms such as 'spirituality', 'knowledge', 'sense of place' and 'aesthetics'. In our framework we interpret these as benefits that are derived by the ways in

which humans interpret landscapes, including soil landscapes, in terms of human needs and values. This is an important distinction because we need to be able to consider how management of soil ecosystems might affect landscapes separately from how these effects might be interpreted by humans.

- It is also common to include ‘control of pests and diseases’ as a ‘regulatory service’. We prefer to describe the service as ‘regulation of species and populations’ because whether or not species are pests depends on human perceptions. This is important because improving the control of potential pests, like aphids in orchards, has been achieved through encouraging soil biodiversity rather than targeting pests per se (Colloff *et al.* 2003; Colloff *et al.* 2010).
- We have included pollination as a soil ecosystem service, because some pollinators (e.g., beetles) have a life-stage that occurs in soil and/ or live in soil as adults. We note, however, that pollination is a final service in some situations (e.g., it contributes directly to production of many crops, separately from the contributions of soil fertility) and an intermediate service in others (e.g., it contributes to the support of native vegetation by soil ecosystems).
- Some other studies have identified ecosystem ‘disservices’, such as salinisation, acidification, erosion and carbon decline (Swinton *et al.* 2007b; Bennett *et al.* 2010). We regard these as symptoms of declines in ecosystem services and we consider them as degradation processes in Figure 8.1, after Dominati *et al.* (2010).

The importance of distinguishing between intermediate and final services was explained in Section 8.1. It can be illustrated in relation to pollination. If this distinction is not made, there is a risk of counting the contribution of pollination more than once in environmental accounting or economic evaluations: one in its own right and again as part of the value of native vegetation. On the other hand, it is important that the contributions of soil biota to fertilising crops are considered in addition to the soil processes that maintain soil fertility, even though the values of both are included in the value of crops produced. This is because the ways in which the benefits are managed by farmers might be different (e.g., farmers might manage soil fertility by addition of fertilisers and might manage pollination by hiring the services of bee-keepers and both of these will be separate items in a farm’s accounts).

Our framework identifies 13 major ecosystem services and 12 groups of benefits from soils. Focusing on benefits and beneficiaries is one way to translate complicated scientific concepts and language for other stakeholders (Ringold *et al.*

2009; Ringold *et al.* 2011). Despite the complexity of the interactions involved, it is possible to make qualitative or semi-quantitative assessments of the relative impacts of different management regimes on different ecosystem services (Foley *et al.* 2005; Bennett *et al.* 2010; Gordon *et al.* 2010) (Figure 8.3). If enough information is available then these benefits can be estimated in monetary terms (Section 9). In Section 8.3, we consider the potential effects of better soil management on ecosystems services in more detail, and in Section 9 the economic implications are considered.

We have depicted only broad groups of beneficiaries in our framework (Figure 8.2 and Table 8.2); when dealing with specific situations it is useful to consider beneficiaries in greater detail than we have (Ringold *et al.* 2009; Ringold *et al.* 2011).




Figure 8.3: Two generalised assessments of differences in ecosystem services from 'natural' ecosystems and agricultural land (Foley *et al.* 2005; Gordon *et al.* 2010)

The further out from the centre the bold line crosses the axis for each ecosystem service the greater the relative production of that service

Table 8.2: Example of the beneficiaries of soil ecosystem services

Beneficiaries	Examples of how they benefit
Farmers and agricultural industries	<p>Production of crops is supported by provision of fertile soil, pollination from animals living in soils and native vegetation, and the role of soil/ plant ecosystems in channelling water into places where it can be used for crops. Raising stock is supported by the role of soils in supporting native (and introduced) pastures and by provision of clean water, filtered and detoxified by soil/ plant ecosystems. Costs of disposing of animal wastes are much lower than they would be if soil ecosystems did not do part of the job.</p> <p>Costs of running machinery are reduced when water has been filtered of sediment by soil ecosystems. Soil provides physical support for farm buildings and structures like dams and levy banks.</p> <p>Stock and crops are protected from heat and floods by native vegetation supported by soil ecosystems, which usually leads to higher yields. The structural components of soils ecosystems, including plant roots, protect against wind and water erosion, reducing costs of replacing nutrients and soil itself and reduced costs of damage.</p> <p>Soil/ plant ecosystems host a range of species that provide pest control by attacking pests of crops. The natural dynamics among species in ecosystems regulated most populations of species and stops them becoming pests or weeds. These processes also control many disease organisms.</p>
Other industries	<p>Industries associated with agriculture, including processors and retailers of food, benefit from the ecosystem subsidisation of food prices – often more than farmers do as profits in these parts of the food supply chain tend to be higher than for farmers.</p> <p>Many other industries rely on clean water and protection from wind and water erosion that could damage infrastructure. Industries that discharge wastes into the environment receive benefits from natural waste breakdown by soil ecosystems. Some industries rely on products from soil ecosystems or ecosystems supported by soil processes (e.g., wildflower harvesting, timber industries, commercial harvesting of fungi or 'bush tucker', peat for fuel).</p>
Individuals, households and communities in rural areas and Individuals, households and communities in rural areas	<p>In both rural and urban areas, individual, households and communities benefit, directly or indirectly, from all soil ecosystem services, but the nature and size of those benefits differs. All Australian households benefit from the production of food and natural products that becomes available in Australian shops. The costs of these products are subsidised by the free soil-fertilisation, water collection, pest control and other services provided to farmers and native vegetation systems by soils.</p> <p>People in both remote and urban areas benefit from water filtration by soil/ plant ecosystems (studies around the world have shown that the cost of providing clean water increases dramatically when catchment areas become degraded). The high health, transport and other impacts and costs incurred by both rural and urban areas during recent dust storms (Leys <i>et al.</i> 2011; Tozer 2012) illustrate the benefits of soil/ plant ecosystems controlling soil stability. Soil stabilisation services, which limit erosion by water and help protect against impacts of flooding, also benefit all people, but especially those living near rivers or in urban areas where water flows could affect life and property.</p> <p>All people benefit from the contributions of soil ecosystems to local regulation of climate and to control of the gaseous composition of the air and air quality (through such processes as absorption of heat, reflection of sunlight, contributions to water cycles that influence rainfall, exchange of gases with the atmosphere, and removal of pollutants and particles from the air).</p> <p>Similarly, all people benefit from the absorption of wastes and pest control by soil ecosystems. People in rural areas may make more direct use of such services and benefits, but people in urban areas still reap the benefits through lower costs of waste disposal than would be the case if soils were not in functional condition. Research in heavily urbanised parts of the world has shown that waste absorption</p>

Beneficiaries	Examples of how they benefit
	<p>capacity of soils is being outstripped by waste production, causing major population-management costs and health risks to be incurred (Folke <i>et al.</i> 1997).</p> <p>Individuals, households and communities are able to receive intellectual stimulation, education, recreational opportunities and various other cultural and spiritual values from ecosystems of which soils are a part. Often people's 'sense of place' is associated with the type and condition of soils present, for example. Conservation of biodiversity is important to many people and this is supported by soil ecosystems. The ways in which cultural ecosystem services are turned into benefits different considerably between people who live close to these services and those who live remotely. For some people, just knowing that ecosystems and biodiversity are functioning well is value in itself (i.e., 'existence value').</p>

8.3 How better management for soil carbon, pH and erosion might affect ecosystem services

Figure 8.3 shows that agriculture generally shifts the balance of ecosystem services in favour of provisioning services while often degrading the processes that lead to regulatory and cultural services. Similar conclusions have been drawn for the world by the Millennium Ecosystem Assessment (MA 2005), for the UK by that nation's National Ecosystem Assessment (UK National Ecosystem Assessment 2011b) and for Australia by various case studies (Binning *et al.* 2001; Abel *et al.* 2003; Karanja *et al.* 2007; Bennett *et al.* 2010; Maynard *et al.* 2010).

As indicated in Figure 8.3B, the aim of modern agricultural management is to restore this balance as much as possible. This is not simply a response to concerns about conservation of biodiversity. As shown in Tables 8.1 and 8.2, there are many benefits that accrue from soil (and other) ecosystems in agricultural landscapes that are socially and/ or economically important to people across society. In this section, we consider how the sorts of best-practice management of soils discussed in previous Sections might be expected to affect ecosystem services and benefits from agricultural landscapes.

The research reviewed in earlier parts of this report indicates that many of the current and emerging approaches to managing soils in Australia appear to be effective, or have the potential to be effective, at addressing the major concerns of declining soil carbon content, increasing pH in some areas, and wind and water erosion (Table 8.3).

It is not easy to capture interactive effects in a table like Table 8.1. While increasing soil organic matter has many benefits for soil structure and processes, for example, excessive accumulation (e.g., in grazing, dairy and some cropping systems) can reduce soil pH (Schumann 1999). Similarly, while inclusion of a pasture phase in

crop rotations provides ground cover and potentially reduces wind and water erosion, if too many stock are run on that pasture then there is the potential for adverse effects on the soil surface that could increase susceptibility to erosion.

Table 8.3: Conclusions from this report about the effectiveness of management practices in Australian agricultural lands for addressing declining carbon content of soil, acidification and wind and water erosion^a

<i>Practice</i>	<i>Type of agriculture</i>	<i>Increases Carbon content</i>	<i>Reduces risk of wind erosion</i>	<i>Reduces risk of water erosion</i>	<i>Reduces risk of soil acidification (low pH)</i>
Soil pH testing	Broadacre cropping	Indirectly	Indirectly	Indirectly	Yes
	Horticulture	Indirectly	Indirectly	Indirectly	Yes
	Dairying	Indirectly	Indirectly	Indirectly	Yes
	Grazing (beef cattle/ sheep meat)	Indirectly	Indirectly	Indirectly	Yes
Soil nutrient testing	Broadacre cropping	Indirectly	Indirectly	Indirectly	Yes
	Horticulture	Indirectly	Indirectly	Indirectly	Yes
	Dairying	Indirectly	Indirectly	Indirectly	Yes
	Grazing (beef cattle/ sheep meat)	Indirectly	Indirectly	Indirectly	Yes
Lime or dolomite applied to reduce soil acidity	Broadacre cropping	Indirectly	Indirectly	Indirectly	Yes
	Horticulture	Indirectly	Indirectly	Indirectly	Yes
	Dairying	Indirectly	Indirectly	Indirectly	Yes
	Grazing (beef cattle/ sheep meat) ^a	Indirectly	Indirectly	Indirectly	Yes
No cultivation/ tillage apart from sowing	Broadacre cropping	Indirectly	Yes	Yes	
Crop residue left intact	Broadacre cropping	Yes	Yes	Yes	
Reduce fallow	Broadacre cropping	Yes	Yes	Yes	
Monitoring of ground cover	Grazing (beef cattle/ sheep meat)	Yes	Yes	Yes	
Use of ground cover management targets*	Grazing (beef cattle/ sheep meat)	Yes	Yes	Yes	
Pasture phase in crop rotations	Broadacre cropping	Yes	Yes	Yes	
Increasing perennial pastures	Grazing (beef cattle/ sheep meat)	Yes	Yes	Yes	

^aThis table draws not only on the material reviewed in this report but also on Barson *et al.* (2011, 2012a, b, c)

The literature also indicates that levels of soil carbon and acid in soils, as well as the extent of wind and water erosion, affect most of the processes expected to generate ecosystem services and therefore the actions to address them are expected to enhance ecosystem services and the benefits flowing from them. The nature and extent of those enhancements, however, will vary with different land systems, land

uses and management regimes (Table 8.4), and improvements cannot be assumed to be linear (see Section 8.4).

Table 8.4: Ways in which actions to address soil condition are likely to affect soil processes and ecosystem services*

Ecosystem services	Practices		
	No cultivation/ tillage apart from sowing/ Crop residue left intact/ Reduce fallow	Managing ground cover above 50%/ Pasture phase in crop rotations/ Increasing perennial pastures	Lime or dolomite applied to reduce soil acidity
Provision of fertile soil	Reduced disturbance is likely to allow soil ecosystems to develop, accumulating soil carbon and nitrogen and engineering soil structure for better water-holding and infiltration capacity	As well as benefits from stabilisation of the soil surface and improved structure and water infiltration, interactions between above ground and below ground ecosystems has the potential to improve carbon and nitrogen cycling.	Reducing acidity will enhance habitat and the activity of many soil organisms. The improvements are likely to be minimal until some pH threshold is reached and soil communities are likely to go through several structural transformations as pH increases.
Support native vegetation	The ability of soils to support native vegetation is likely to be enhanced by reduced use of fertilizers on agricultural land, because fertilizers are likely to change the composition and functioning of native ecosystems. However, if increased use of pest-control chemicals is required then this could have negative impacts on organisms in soils under native vegetation.	Reduced runoff of agricultural chemicals onto soils under native vegetation is likely to be the biggest benefit	Addressing soil acidity on agricultural land might have benefits for soils under adjacent native vegetation by reducing leakage of acid into water tables. However, most cost-effective approaches are likely to only manage topsoil acidity.
Provision of natural products	As above	As above	As above
Provision of clean water	Increased stability of soil, structural involvement of vegetation, and enhanced activity of soil organisms is likely to increase water filtration and detoxification capacity of soils.		To the extent that reduced acidification improves activity of soil organisms and soil structure it will contribute to water filtration and purification.
Maintenance of genetic diversity	Enhancement of the diversity of conditions for soil organisms is likely to improve persistence of genetic diversity both within agricultural soils and in adjacent soils.		As above – reduced acidification is likely to lead to at least small improvements in habitat and genetic diversity below ground.
Water flow regulation	Reduce overland flow of water, reduced evaporation and improved infiltration are all likely to affect hydrological cycles (e.g., increasing recharge of		To the extent that managing acidity improves soil structure and

Ecosystem services	Practices		
	No cultivation/ tillage apart from sowing/ Crop residue left intact/ Reduce fallow	Managing ground cover above 50%/ Pasture phase in crop rotations/ Increasing perennial pastures	Lime or dolomite applied to reduce soil acidity
	water tables, reducing damage from floods)		infiltration rates and/ or allows better establishment of ground cover, it is likely to affect water flows (impacts likely to be small under realistic acid management approaches at present)
Maintenance of landscape (soil) stability	Improved ground cover and minimisation of soil disturbance contribute to soil stability and reduce risks of dust storms, landslides and water erosion		As above
Regulation of atmospheric gases	Improvement of carbon capture by soils will affect atmospheric CO ₂ (indications are that this effect is likely to be small under most realistic scenarios). Depending on the crops or pastures grown, nitrogen exchange with the atmosphere could be affected (this effects is likely to much more significant for soils than the atmosphere)		Small impacts on carbon and nitrogen cycles (as above)
Regulation of weather and climate	Vegetation cover has effects on absorption and radiation of radiant energy from the sun, affecting the temperature of the ground (and hence the environment for below ground organisms). It also affects moisture and air movement close to the ground. There are likely to be effects on local weather (evaporation, cloud formation etc.) but these are likely to be small at the scale of most agricultural management. The exception is when ground cover is inadequate (i.e., the ecosystem service of stabilising soil landscapes is not adequate) and wind erosion results in dust storms that can influence weather considerably (Mahowald <i>et al</i> 2010; Rotstayn <i>et al.</i> 2012).		As above – small impacts to the extent that addressing acidity affects ground cover.
Remediation of wastes	As for provision of clean water		
Regulation of species and populations in soils	To the extent that these approaches encourage species diversity, there will be effects on interactions among species. Community structure is likely to change. There is evidence that improving ground cover can enhance control of above ground pests (e.g. aphids) by below ground species (e.g. in orchards).		To the extent that addressing acidity encourages soil biodiversity (see above) there could be improvements to pest control benefits arising from below-ground population regulation.
Contributions to species, ecosystem and landscape diversity	Improved condition of soils is likely to change the appearance of landscapes and, therefore, the benefits they provide to different groups of people. Perceptions will vary between beneficiary groups. Some will benefit from recreational, spiritual, educational and other cultural aspects of improved condition of native vegetation systems (by experiencing these improvements or just knowing they are occurring). Others will benefit from aesthetic and other cultural aspects of landscapes relating to agricultural productivity. There are likely to be broad cultural benefits from seeing and/ or knowing that degraded landscapes are recovering.		

*This table draws on the rest of this report and, particularly, a number of key synthesis and review

paper (Pimentel *et al.* 1995; Seybold *et al.* 1999; Binning *et al.* 2001; Colloff *et al.* 2003; MA 2005; Lavelle *et al.* 2006; Swinton *et al.* 2006a; Barrios 2007; Swinton *et al.* 2007a; Zhang *et al.* 2007; Haygarth and Ritz 2009; TEEB 2009; Bennett *et al.* 2010; Clothier *et al.* 2011; UK National Ecosystem Assessment 2011a; Griffiths and Philippot 2012; Robinson *et al.* 2012)

8.4 Resilience of soils and associated ecosystems

Resilience is a word and a concept that has become increasingly widely used, across many disciplines, over the past decade (Holling 1996; Folke *et al.* 2002; Folke *et al.* 2004; Walker *et al.* 2004; Walker and Salt 2006; Brand and Jax 2007; Cork 2010a). There is still debate about precise definitions and ways to measure this attribute in relation to ecological, social, organisational and other systems, and it is necessary to review some key aspects of this debate in order to consider resilience of soils.

Often, people equate resilience with ‘health’, ‘condition’, or ‘vigour’ – the ability to ‘bounce back’ after shocks. While soil condition is an important aspect of resilience in many cases, there is much more to soil resilience than condition. This section discusses important concepts that have arisen in the soil literature that relate condition (the subject of the rest of this report) to the broader issue of resilience. These concepts include: debate about whether soils have a ‘single stable state that they return to or whether we have to consider a degree of change in state as part of resilience; the idea that resilience might be different at different scales; the different rates of soil degradation versus recovery; the idea that some degraded states can be highly resilient (i.e., resilience is not always a desirable quality); and the important difference between resilience and resistance to change, which affect the short versus long-term responses of soils.

The 2011 State of the Environment Report (Australian State of the Environment Committee 2011) included, for the first time in state of the environment reporting in Australia, a discussion about soil resilience. This discussion focussed on the key aspects of soil condition that allow it to continue to function through perturbations like climatic variation and change and physical disruption by land management practices. It included that good-quality and resilient land has these related features:

- Leakage of nutrients is low.
- Biological production is high relative to the potential limits set by climate.
- Levels of biodiversity are relatively high.
- Rainfall is efficiently captured and held within the root zone.
- Rates of soil erosion and deposition are low, with only small quantities transferred out of the system (e.g. to the marine environment).

- Contaminants are not introduced into the landscape, and existing contaminants are not concentrated to levels that cause harm.
- Systems for producing food and fibre for human consumption do not rely on large net inputs of energy.

The State of the Environment report also pointed to the fact that older, more weathered soils, such as those in most of Australia, are less able to return to their original state after perturbations than younger soils. It discussed the role of clays in allowing some Australian soils (e.g., Vertosols) to recover from compaction. This issue is discussed in relation to resistance versus resilience of soils below. It also discussed the importance of considering thresholds of change, especially with respect to organic matter decline, soil acidity and erosion. The significance of thresholds in relation to soil resilience is also discussed further below.

Since Holling's (1996) landmark paper, a distinction has been made between 'engineering resilience' (return of a system to a previous state after perturbation) and 'ecological resilience' ("the capacity of a system to absorb disturbance and reorganize while undergoing change so as to still retain essentially the same function, structure, identity, and feedbacks") (Walker *et al.* 2004). Folke *et al.* (2002) concluded that resilient ecosystems: "can cope, adapt, or reorganize without sacrificing the provision of ecosystem services".

A key difference between these two approaches is that the former assumes that the system has a single stable state (or if there are alternative states they should be avoided), while the latter assumes that ecosystems can exist in multiple stable states and that resilience is the property of the system that keeps it within the bounds of a particular state (Botton *et al.* 2006).

When considering multiple stable states, the concept of 'hysteresis' becomes important (Lal 1997; Seybold *et al.* 1999; Potts *et al.* 2006). Hysteresis is the difference between degradation and recovery phases, in terms of the rates of recovery and the processes involved. For example, resilient soils often will take much longer to recover their functions than it took to lose them (Lal 1997). This concept is particularly relevant when considering the ability of soils to cope with declining organic matter or increasing acidity. Natural processes or human intervention can help soils rebuild carbon stores and enhance the many processes reliant on carbon and/ or living components of soil when soils have sufficient reserves of minerals and retain sufficient diversity of living components (Seybold *et al.* 1999; Botton *et al.* 2006; Jiang and Patel 2008; Griffiths and Philippot 2012; Kuske *et al.* 2012), but the record of past perturbations is important and recovery can

take decades (Kuske *et al.* 2012). Similarly, recovery from acidification can be very long term, especially if sub-soils are affected (Section 5).

Another way to interpret hysteresis is that degraded states often have high resilience and/ or resistance to remediation. The broader literature on resilience has recognised that ecological and/ or social resilience is not always desirable to humans. Apart from highly acidified, carbon-depleted or eroded soils, polluted soils can have high resilience (Botton *et al.* 2006).

The concept of 'panarchy' is also particularly relevant to considering resilience of soils. This is the idea that the resilience of any 'system' is affected by other systems operating at higher and lower scales (Gunderson and Holling 2002). For example, the resilience of the soil ecosystem at a paddock scale will be influenced both by ecosystems operating within the soil and by processes occurring at landscape, regional and even larger scales, including interaction between soils, plants, animals and the atmosphere and interactions between ecological and human social systems. Most soil recovery mechanisms and ecosystem services are biologically mediated, including cycling of nutrients, detoxification of pollutants, and suppression of pathogenic organisms (Seybold *et al.* 1999). Neither recovery of soil organic matter nor rebuilding of resistance to wind and water erosion can be accomplished without considering inputs from plants as organic matter and through their mutualistic associations with soil organisms. Also, as explained below, resilience of soils cannot be considered without reference to human social and economic processes.

Research on ecological and social resilience has emphasised the importance of the question: "resilience *of* what *to* what?" (Carpenter *et al.* 2001). Defining 'essential functions, feedbacks and identify' (*of* what) is essential if we are to judge whether these are being retained. Systems might have resilience to some 'specified' (known, previously experienced) pressures but not others (*to* what). The characteristics that give a system specified resilience can be different from those that give 'general' resilience (Walker and Salt 2006).

Most often, soil resilience has been defined as the capacity of a soil to recover its functional and structural integrity after a disturbance (see reviews by Lal (1997); Seybold *et al.* (1999); Botton *et al.* (2006)). This resembles the engineering concept of resilience, although there has been recent discussion about the concept of multiple stable states applied to soils. For example, research on soil microbial populations indicates that community composition and structure change dramatically with wetting and drying and other perturbations resulting in alternative stable states that exhibit hysteresis (Seybold *et al.* 1999; Botton *et al.* 2006; Potts *et al.* 2006;

Jiang and Patel 2008; Griffiths and Philippot 2012). At the scale of managing agricultural enterprises, however, the essential functions required of soils are defined by the uses that land managers wish to make of the soils. Lal (1997) pointed out that these uses are influenced by: “the socio-economic and political forces that govern land use, land rights, institutional support, and income”.

A related concept of ‘resistance’ refers to tendency of a system’s attributes (e.g., structures and functions) to not fluctuate when perturbed (Lal 1997; Seybold *et al.* 1999; Botton *et al.* 2006). For example, some soils resist compaction and retain their porosity while others suffer compaction but are able to regain porosity after a period of time (Seybold *et al.* 1999). These differences between resistant and resilient soils are important as they affect responses of crops and pastures in the short and long term. Similarly, ground cover above a critical threshold confers resistance to wind and water erosion (Sections 6 and 7), whereas resilience to wind and water soil erosion is a function of the depth and type of soil and the rate of soil formation, which, in many parts of Australia, is many times slower than rates of erosion (Section 7.1).

Often the distinction between resilience and resistance is blurred. For example, different wetlands in the Murray Darling Basin have very different abilities to neutralise acids formed when sediments are exposed by dry periods (Glover *et al.* 2011). This ‘acid neutralising’ capacity confers both resistance and resilience (within the limits of the system’s buffering capacity) on these wetlands.

Research on ecological resilience generally has revealed the importance of considering thresholds of change (rapid, often irreversible changes that take a system into a different state) (Walker and Salt 2006). An important aspect of the resilience of a soil ecosystem is its ability to stay away from such thresholds and, in general, its resilience will be lower the less disturbance is required to push the system through a threshold of change. A range of thresholds have been suggested for soil ecosystems (Lal 1997):

- An organic carbon threshold (varying with soil type but usually 1-2% in surface layers) below which physical and chemical fertility effectively collapse and after which recovery of critical carbon fractions can take decades (Baldock and Skjemstad 1999; Australian State of the Environment Committee 2011);
- A soil pH threshold (around 4.2) below which aluminium toxicities emerge and the soil becomes very difficult to remediate (Australian State of the Environment Committee 2011) (Section 4);

- Ground cover thresholds (50-70%) below which soils are vulnerable to erosion by wind and water (Section 6);
- A postulated lower vegetation-cover threshold (20% in Chinese grasslands), below which ecosystems cannot recover by themselves from sustained degeneration of the vegetation community, erosion of the surface soil and declining soil fertility (Gao *et al.* 2011);
- Non-linear changes in many soil properties (e.g., water flux, porosity, mineral dissolution rates, redox potential and acid-base reactions as carbon is added (Chadwick and Chorover 2001);
- Thresholds of inadequate sediment flows (resulting in the loss of beaches, storm protection, nutrient inputs, etc.) or excessive flows (resulting in lake, reservoir and wetland infilling, coral reef smothering, etc.) (Apitz 2012);
- Physical damage to biocrusts (e.g., by grazing), in concert with changing temperature and precipitation patterns, has potential to alter performance of dryland ecosystems for decades (Kuske *et al.* 2012);
- Catastrophic shifts in soil-vegetation systems due to interactions between herbivores, plants and below-ground ecological systems (van de Koppel *et al.* 1997);
- Over-saturation of soil nutrients leading to accelerated leaching to water courses (Heckrath *et al.* 1995);
- Local extinction of certain strains of bacteria when soils become contaminated by toxic pollutants (Chaudri *et al.* 2008);
- Thresholds of suitability of soils when used for sub-optimal purposes (e.g., using soils as raw materials or using soils suitable for growing food as a platform for building upon) (Haygarth and Ritz 2009).

Multiple factors influence soil resistance and resilience (Lal 1997; Seybold *et al.* 1999; Botton *et al.* 2006; Zhang *et al.* 2010; Griffiths and Philippot 2012). They are partly related to soil properties such as organic matter, aggregation, the quantity and quality of carbon inputs, clay content and soil pH. Terrain characteristics, landscape position, parent material, climate, water balance, vegetation and soil biodiversity are also important. Research on the contributions of the living components of soils to soil resilience has focused primarily on microbial populations (Seybold *et al.* 1999; Botton *et al.* 2006; Zhang *et al.* 2010; Griffiths and Philippot 2012). This research reveals no simple general rules but suggests that the diversity of functional traits of species is important, as is the structure of communities (lower resilience when

communities have a highly uneven balance between species or are dominated by a few species). There is an expectation that high levels of functional redundancy, i.e., a high number species performing the same function, might act as a buffer against the effect of biodiversity loss on functioning. Resilience and resistance become much more complex issues under extreme perturbations such as contamination of soils with toxic compounds, which select very rapidly for species that can deal with the challenges.

Each of the best-practice management approaches to dealing with soil carbon, pH, and the threat of erosion (i.e., those summarised in Table 8.3) potentially contributes to the requirements for increasing resilience after perturbations. Processes important for returning soil function after perturbation include new soil formation, aggregation, soil organic matter accumulation, nutrient cycling and transformation, leaching of excess salts, and increases in biodiversity, including species' succession (Lal 1997). When applying best-practice management for specific challenges to soil condition, however, it will be important to consider how the range of management practices being implemented interact with one another and to consider the specified as well as the general resilience of the resulting soil ecosystems. For example, managing ground cover to appropriate targets can improve soil carbon status, and reduce wind and water erosion, while managing soil acidity through liming can also overcome a major constraint to building carbon and having adequate ground cover (Table 8.3).

Approaches to assessing soil resilience involve assessing actual functionality through time, or indicators of functionality, in relation to reference states, and considering thresholds of undesirable change (such as those discussed above) and how to avoid them (Lal 1997; Seybold *et al.* 1999; Botton *et al.* 2006).

8.5 Economic values of soil ecosystem services and resilience

Many of the benefits that can come from ecosystem services can be expressed in monetary terms, because they include goods that are sold in markets or involve other financial transactions that reveal people's willingness to pay for the benefits (Costanza *et al.* 1998; Bennett 1999; Bockstael *et al.* 2000; Gillespie *et al.* 2008; TEEB 2009; UK Government 2011). Resilience has been included as a benefit from ecosystems in some recent typologies (TEEB 2008).

The economic values of soil ecosystem services have been estimated in a variety of ways in different studies in different parts of the world. The approach taken depends on the questions being asked. Some studies have estimated the replacement cost of soil ecosystem services. When we consider how processes like large-scale nutrient

and water cycling, extraction of nutrients and carbon from the atmosphere, acid-based balance, waste breakdown, regulation of hydrology and pest control could be replaced by engineered alternatives, including provision of fertilizers and other chemical components, the costs are massive (Daily *et al.* 1997; Sandhu *et al.* 2008).

Replacement costs of soil ecosystem services are not, however, relevant to the questions being asked in this report (Section 9). The value that farmers, and others who use ecosystem services in production of goods and services, (i.e., 'producers') might get from better soil management is more appropriately estimated as the difference between what they would be willing to accept as payment for the goods and the price they receive in the markets ('producer surplus'). The contribution of ecosystem services to producer surplus is a function of how much their use reduces production costs. The proportion of total ecosystem service production that is used depends on the time period, from very small over a short time period to total use if an ecosystem is totally degraded in the long term, and the degree to which natural capital is consumed by the production activity. The value that consumers (including the broader public) get from ecosystem services is most appropriately estimated as the difference between what they would be willing to pay for the benefits and what it actually costs them (consumer surplus). Consumer surplus is complex to assess. It can be partly estimated by assessing consumers' willingness to pay for access to ecosystem services (TEEB 2008; MacDonald *et al.* 2011; CSIRO 2012) but this often will not take account of the savings that people make through such benefits as better mental and physical health.

Section 9 considers the economic benefits of better soil management in Australia, by considering the net benefits across a range of case studies. Management practices are not the only factors affecting the adequacy of soil ecosystem services to meet human needs. Climatic factors obviously play a major role, and it is important that soils are managed appropriately for the climate they are exposed to. This is a key component of best-practice management. In Australia, drought should no longer be used as an excuse for degradation of soils as management of soil resilience should include management for wet and dry periods. Apart from factors affecting the supply of ecosystem services, demand for them is an important consideration. Demand for ecosystem services is affected by where and how people live, infrastructure for turning services into benefits, and economic pressures coming from outside a region or Australia. We are unable to take these extrinsic factors into account in this project, but they should be considered as part of broader population planning in Australia in the future (Cork 2010b).

9. Private and public benefits of soils and soil management

9.1 Introduction

This section takes the discussion of ecosystem services in the previous Section a step further and reviews estimates of the value of ecosystem goods and services provided by Australian soils managed for agriculture. It addresses the following questions.

What is the nature of benefits from improving agricultural soil condition?

Who benefits from improving agricultural soil condition?

How significant might these benefits be?

How might Australia realise these benefits?

In this review, we have considered the net benefits that are likely to flow from improved soil condition and better quality soil ecosystem services from agricultural lands. We have not tried to address questions of how to optimise benefits from soil ecosystem services, nor how to balance public and private investment in soil condition.

9.2 What is the nature of benefits from improving agricultural soil condition?


Understanding the benefits from improving agricultural soil condition requires a framework for distinguishing between benefits to human wellbeing; final ecosystem services; the natural capital (or soil condition) which underpins those services; soil depreciation and accumulation processes; and the external drivers which influence soil condition. The framework we use in this report is discussed in Section 8.

In this report, we focus on the marginal change in benefits that ultimately come from a change in land management practices. In the short term, these benefits are generally improvements to agricultural productivity or the reduced cost of impacts off-site from agricultural lands. In some cases, the benefit may come from keeping open future options to produce different types of crops in response to changing market demand or climate. In theory, soil conditions which support a wider range of future uses will be reflected in a higher capital value of the land (Gretton and Salma 1996).

9.3 Who benefits from improving agricultural soil condition?

Benefits can improve the wellbeing of private landholders, or the public, or both. They may occur at a local (on-site), regional, national or global spatial scale. They may be realised over short (1-5 year), medium (5-30 year) or long (30-100 year) timeframes.

Figure 9.1 shows some examples of who benefits from final soil ecosystem services, where the benefits are realised, and over what timeframe. It also shows whether the value of the benefit is in the form of a flow of services (similar to financial interest), an option to maintain future benefits (similar to insurance) or a stock of soil condition (similar to financial capital).



Soils can provide many different ecosystem services (Figure 9.1). Yet not all services can be provided at once. Land management decisions involve trade-offs between different types of benefits (Robertson 1987). For example, in the short-term, at least, there can be trade-offs between stock production and maintenance of ground cover. The perspectives of different beneficiaries lead to a range of views about which benefits are more important.

For landholders, agricultural systems primarily produce food and fibre. The challenge is to optimise long-term production, and build soil resilience to external drivers such as climate and degrading processes such as erosion or acidification. This means producing stable agricultural returns without compromising the future ability of soil to support crops or livestock, or increasing soil vulnerability to erosion and acidification.

However, the sixty percent of the land mass managed for agriculture is part of the broader Australian landscape. At this scale a number of ecosystem functions are important, for which soils may provide supporting services. For example, soils support the production of native grasses, which are both habitat and food for Australia's diverse range of native fauna.

Australia's agricultural industries are also part of the broader Australian economy, and the global system of food trade. In this context, reliability of agricultural production is important for contributing to Australia's economic stability. As the global population rises, the reliability of Australian agricultural production may also be important for food security. This could become increasingly significant as external drivers such as commodity prices and weather may be more volatile in the future (OECD/FAO 2011).

In this report, we have focused on benefits to land managers at the farm scale- and to the Australian public at a local, regional or national scale.

9.4 How significant might these benefits be?

We have reviewed existing economic studies to assess what is known about the magnitude of benefits from improved land management practices. We selected studies that have a clear link between costs or benefits and soil ecosystem services. However, the economic values estimated are not all attributed to changes in soil condition. Other factors and agricultural inputs also contribute.

Table 9.1: Gross value of agricultural production (ABS 2011a)

Industry sector	Gross value of agricultural production – average 2008 – 2010 (\$ billion/yr)
Broadacre cropping	9.6
Beef/sheep grazing	9.8
Horticulture (excluding grapes)	8.4
Dairy	4.0
Other agriculture	9.8
Total	41.2

Australia's total average agricultural production is \$41.2 billion per year. The four industry sectors covered in this report, account for around 75% of this production (Table 9.1).

Given the importance of these industries, a number of previous economic studies have examined the links between soil condition, agricultural production and public benefits. Table 9.2 outlines the most relevant previous studies.

The benefits of improving agricultural soil condition need to be calculated against a baseline soil condition. In many cases, the baseline soil condition is on a moving trajectory. The nature of this trajectory, and its likely impact on agricultural production or off-site environmental impacts, is complex. Some aspects of soil condition may be improving, while others may be declining. Both land management practices and external drivers may be responsible for these changes.

The very act of farming alters soil condition (Robertson 1987). Managing land for agriculture shifts soil ecosystem processes toward increased production of crops or pasture, and away from other intermediate or final services (Pretty 2008). In some cases, these other services can be augmented, replenished or replaced by external inputs. For example, adding phosphorus fertilizer can augment the ability of soils to provide nutrients. Trace elements lacking in Australia's weathered soils may be replenished by agricultural practices. Organic matter lost due to erosion or intensive cropping may be replaced by manure or green waste.

However, soil degradation problems occur if land management practices produce short term gain, at the cost of declining soil condition (Robertson 1987). The benefits of improved land management practices therefore depend on improved productivity, calculated against the expected cost of a continuing decline in soil condition. However, improved productivity may be seen in short to medium term, whereas the costs of inaction may only be apparent in longer term.

The complexity of agricultural and natural systems, as well as gaps in knowledge and data, make it difficult to accurately predict the economic impacts of changing land management practices (Gillespie *et al.* 2008; Rolfe *et al.* 2008). Many studies to date have focused on the private benefits of near-term production values; with some estimates of the avoided public cost of damage from erosion or rising water tables; as well as public willingness to pay for environmental benefits. In most cases, these estimated values are specific to a certain region, and may not apply across the diverse range of soil types, conditions, land-use and land management practices found in Australia.

Table 9.2: Existing estimates of the value of costs or benefits related to land management practice (footnotes explained at end of table)

Ecosystem service ²	Benefits	Time-frame ³	Net value	Example ⁴	Comment
Provisioning services					
Soil condition suitable for growing crops or feed (through maintenance or improvement against declining baseline)	Increased nutrients	Short	Positive if private benefits > private costs	\$14-\$16 per hectare additional production potential from reducing acidity in NSW (Walpole <i>et al.</i> 1996). \$10.8-\$16.5 billion NPV of additional production through lime/ gypsum treatment of the 4% of land at risk of acidity and sodicity where soil treatment is profitable (at a 10% discount rate) (Hajkowicz and Young 2002). It has been estimated that Western Australian farmers face an opportunity cost of lost agricultural production from soil acidity of around \$498 million/ year (Herbert 2009).	These figures over-estimate soil ESS as they include benefits from other inputs paid for by farmers. They are not marginal values, as they assume all soil degradation is avoided.
	Greater economic stability	Long	Positive	Avoided cost of 9 cents per household for every 10 persons remaining in regional communities (Hajkowicz and Young 2002).	Choice modelling shows Australians perceive rural depopulation as a cost.
Native vegetation support (through maintenance of soil condition)	Maintenance of existing native vegetation	Medium - Long	Positive if public and private benefits > private costs	Willingness to pay \$2.90 per household per year over 15 years for a 1% improvement in healthy vegetation in Qld (Windle and Rolfe 2007).	Not all public value can be attributed to soils, as other economic inputs may be required. Private costs will depend on land-management practices, offset to some extent by private benefits e.g. shade for livestock (Fischer <i>et al.</i> 2009).
	Regeneration of native vegetation	Medium-Long	Positive if public and private benefits > private costs	Willingness to pay 7 cents per household per year for every additional 10,000 ha of farmland repaired or bushland protected (Hajkowicz and Young 2002).	As above
	Reduced cost of water treatment and equipment maintenance	Short	Positive	Avoided cost of \$0.8-\$2.0 billion NPV for a 1%-10% decline in water quality based on downstream infrastructure costs of turbidity due to erosion (Hajkowicz and Young 2002).	Not all this value can be attributed to soils as other interventions, like tree planting and erosion control, may be required. Additional public and private benefits would flow from avoiding raised nutrient levels and eutrophication.
Regulating services					
	Protection from	Medium	Positive	\$62 m per year avoidable costs of rising water	Not all this value is due to water table

Ecosystem service ²	Benefits	Time-frame ³	Net value	Example ⁴	Comment
	change in water table levels			impacts on public and private infrastructure based on projections from 2000 to 2020 (Hajkowicz and Young 2002).	levels. Some is due to the avoidable cost of damage from salinity.
Landscape (soil) stabilisation	Protection from erosion	Medium	Positive	<p>Avoided cost of \$2 worth of fertiliser lost with every tonne of soil erosion prevented (Raupach, McTainsh, and Leys 1994)</p> <p>Avoidable private costs of \$5.72-\$8.09 per hectare in net agricultural income in 1989/90, for Lachlan valley and Orange SLA (Mallawaarachchi 1993; Mallawaarachchi, T., Young, M., Walker, P. and Smyth 1994).</p> <p>Estimated public value of \$0.5billion PV for erosion control outcomes from National Heritage Trust investments (Gillespie <i>et al.</i> 2008).</p> <p>Estimated \$23 million per year total off-site wind erosion costs for South Australia. Most of this is health related costs (Williams and Young 1999).</p> <p>Estimated >\$400 million cost of 2009 'Red Dawn' dust storm in Sydney. Most of this is cleaning and lost work time (Tozer 2012). Regional dust storm events are more frequent, their economic impacts are likely to be lower because regional populations are smaller and regions have fewer infrastructure assets. Nevertheless, estimates of the offsite impacts of dust erosion on the Mildura region show that costs to the regional economy are approximately \$3 million annually (Tozer 2012).</p>	<p>Not all this value can be attributed to soil structure stabilisation. Other interventions, like tree planting and erosion control works, may be required.</p> <p>Health related costs assume asthma rates are linked to wind erosion and dust.</p>
Gas regulation	Reduction in carbon dioxide emissions	Long	Positive if public and private benefits > private costs	Recent research suggests improved management could provide relative gains of 0.2-0.3 tonnes of C per ha/year for cropland, and 0.1-0.3 tonnes of C per ha/year for pasture (Sanderman, Farquharson, and Baldock 2010).	Estimating the market value is difficult as enhanced carbon stocks may be lost due to drought or changes in land management practices.
Cultural services					
Existence of soils in good condition	Environmental health	Medium	Positive	Willingness to pay \$3.70 per household per year for 15 years for a 1% improvement in soils in good condition in Qld (Windle and Rolfe 2007).	Choice modelling shows Australians are generally willing to pay for the environmental benefits associated with

Ecosystem service ²	Benefits	Time-frame ³	Net value	Example ⁴	Comment
					improved soil condition.

¹Estimates of the potential cost of increased water turbidity were not included as they can't be clearly attributed to changes in soil condition. We have been unable to find published information about whether turbidity is due to soil erosion or sediments already in streams. Estimates of the costs of turbidity, from all sources, are available in Hajkiewicz and Young (2002).

²This table does not include supporting services, as these generally increase the benefits realised from other ecosystem services rather than directly benefiting humans.

³Timeframes are defined as short (1-5 years), medium (5-30 years) and long (30-100 years).

⁴Unless noted otherwise, all estimates of value are in the dollars of the year of the original study.

Nevertheless, some insights can be drawn from consistency of findings across the range of valuations shown in Table 9.2:

- The lost value of crop yields due to soil acidity may be high. Additional production potential of \$14-\$16 per hectare (1996 dollars) is at least 4% of average NSW broadacre cropping revenue of around \$400 per hectare (ABS 2011a; Walpole *et al.* 1996). Compared to an average annual \$9.6 billion gross value of production of broadacre crops, an NPV of \$16 billion for treating 4% of the land at risk of acidification or sodicity represents a significant opportunity (Hajkowicz and Young 2002). The estimated \$498 million/ year of lost production due to acidity in Western Australia was the highest cost of any hazard, followed by salinity (\$344 m/yr), surface compaction (\$333 million/ year), and water repellence (\$251 million/ year), and far exceeding the estimated losses due to wind erosion (\$71 million/ year), waterlogging/ inundation (\$29 million/ year), soil structure decline (\$15 million/ year), and water erosion (\$10 million/ year) (noting that these hazards are not independent) (Herbert 2009).
- Aggregate public costs of erosion are high, particularly during intense dust storms. Private costs of erosion may be slightly lower than those of acidity, estimated at \$6-\$8 per hectare in NSW (Table 9.2) (but note the comparison for Western Australia, above). However, erosion may have more significant long-term impacts as soil-loss is irreversible.
- Willingness to pay estimates indicate Australians recognise the value of public investment to improvement soil condition, regional jobs, and maintenance of farmland vegetation (assuming Queensland residents surveyed by Windle and Rolfe (2007) are representative of the broader Australian population (Table 9.2)).

Recent assessments of the extent and risk of land degradation have also suggested which industries are likely to benefit most from improving soil condition.

- Managing soil acidity is likely to benefit broadacre cropping, of which 36% is at high risk, and intensively managed grazing land, of which 21% is at high risk (Barson *et al.* 2011, 2012b). Tropical horticulture and dairying are also at risk but available data are too coarse to allow accurate assessment to be made for these industries (Michele Barson, DAFF, pers. comm.)
- Reducing soil loss through wind erosion is most likely to benefit beef and sheep grazing in the rangelands, as well as broadacre cropping in WA (Smith and Leys 2009).

- Reducing soil loss through water (sheet and rill) erosion is most likely to benefit broadacre cropping, as well as sugarcane and other horticulture in Qld (Hairsine *et al.* 2009).
- Increasing soil carbon is most likely to benefit horticulture, broadacre cropping and grazing in NSW, Qld and WA (Baldock *et al.* 2009).

9.5 How might Australia realise these benefits? Examples through case studies

Australia's soils, their condition, land-use and management practices, are highly variable. This makes it difficult to present simple conclusions that apply to all soils in Australia. Instead, we have used case studies of specific industries and, in some cases, locations to draw general findings.

We selected four case studies to demonstrate the issues relevant to considering the private and public benefits of improving soil condition. For each case study we have highlighted how land management practices can:

- Improve agricultural production by reducing or removing soil constraints, stabilising profits, or increasing efficiency of resource use
- Reduce or avoid environmental impacts off agricultural lands
- Address land degradation that occurs over different timeframes
- Face barriers to implementation in addition to costs of implementation
- Be widely applied in Australia.

Case study 1: Reducing soil erosion in broadacre cropping – northern NSW

Broadacre cropping is an important agricultural sector for Australia. Australia wide, production of cereals such as wheat and barley, pulses such as lupins and chick peas, and oilseeds such as canola and sunflower, has contributed \$9.6 billion per year² in gross value of agricultural production from 22 million hectares cultivated across all states (ABS 2011b)³. In NSW, 5.8 million hectares produce almost \$2 billion per year⁴ in gross value of agricultural production (ABS 2011b).

²² Based on averages for 2008 to 2010

³ Note that the gross annual value of production for broadacre cropping given in Section 4.2 (around \$13 billion) included cotton, hay and sugar cane in addition to the crops included in this section

⁴⁴ Based on averages for 2008 to 2010

However, poor structure, low water permeability and soil acidity limit yields in various cropping regions across Australia (Beeston *et al.* 2005). Improving soil condition can increase crop yields, by improving the quality of soil ecosystem services. Maintenance of good ground cover levels results in more stable soils, which reduces wind and water erosion, and therefore the loss of soil nutrients and carbon, which can support crop production. Soils in good condition are also more able to provide nutrients and moisture when crops need it. Where nutrient availability is not a constraint, the level of soil moisture at the time of sowing directly influences the final crop yield (Day *et al.* 2008).

This case study considers the benefits already gained from improving soil structure and water permeability in northern NSW (Table 9.3). Given the benefits of new land management practices are uncertain (due to high variability in soil types, crop types, weather patterns and barriers to adoption), an historical example can offer greater insight than forecasts.

The evolution of farming systems has increased yields in part by improving soil condition, often overcoming negative impacts on soil condition caused by earlier farming practices. Conventional farming systems used before the 1970s tilled the soil, which destroyed the soil structure and increased vulnerability to wind and water erosion (Scott and Farquharson 2004).

Since the 1970s, conservation farming has sought to maintain soil structure and fertility by leaving crop residues on or near the surface. Weed growth is reduced by using herbicides rather than tilling the soil (Barr and Cary 1992). Conservation farming can increase agricultural production, reduce soil loss through wind and water erosion, lower greenhouse gas emissions and improve water use efficiency. Conservation tillage is a key part of conservation farming⁵. Across Australia, 95% of cropped land is now managed with some level of conservation tillage (Barson *et al.* 2012b).

As adoption of newer farming systems increases, significant private benefits from improved soil ecosystem services are often seen. The value of these benefits can be estimated, although it is difficult to separate the contribution of soil ecosystem services from other human and environmental impacts. Public benefits are harder to quantify, but can still be significant.

In northern NSW, existing estimates of net private benefits from conservation farming are significant at an industry level. Between 1970 and 2000, the net present

⁵ Conservation tillage involves no-till or minimal till practices, combined with direct drill seeding techniques.

value of increased agricultural production was estimated at over \$200 million (Scott and Farquharson 2004). At the farm-scale, returns on capital invested increased by around 3.5% within five years of adopting no-tillage practices, compared to conventional farming as a baseline.^{6,7}

A return below commercial investment benchmarks may explain why widespread adoption of conservation farming took several decades. However, conventional farming has shown rapidly declining crop yields and quality in around 20 years (Scott and Farquharson 2004). Using a longer timeframe may therefore show much higher returns on capital invested, compared to business as usual. While these higher private benefits may be clear in hindsight, farmers may be unlikely to take the risk of adopting new practices without some public investment in research, development and extension to prove they work.

Significant public benefits also came from adoption of conservation farming. Soil erosion was reduced by an estimated 18 million tonnes per year (Scott and Farquharson 2004). Public benefits from the increase in gross value of agricultural production would have flowed through increased economic activity at local, regional and state levels.

The key findings from this case study are:

- Private benefits of improved crop yields were apparent within 5 years.
- Public benefits included reduced off-site environmental costs of dust, and possibly greater economic contributions from the broadacre cropping industry.
- Over a 5 year time-frame, private returns on capital invested were below commercial rates. However, over a 20-year timeframe they may be much higher.
- The low 5-year returns on capital invested and risk aversion to adopting new practices may have been barriers to private investment in improving soil condition.
- Conservation farming practices are now widely used within Australia, with current rates of adoption 95%.

⁶ This is based on whole farm budget estimates in two locations.

⁷ The total private benefits of increased agricultural production across northern NSW were estimated at \$224m (no till) and \$586m (no till plus reduced tillage). The benefit share due to NSW Government investment was estimated at 35 % of these total figures, giving \$78.4m (no till) and \$205.4m (no till plus reduced tillage).

Table 9.3: Full range of benefits and beneficiaries – Reducing soil erosion in broadacre cropping

Beneficiaries	Ecosystem services	Benefits	Costs	Time-frame	Expected net value
Private land-holders	Landscape (soil) stabilisation Soil condition for crops	Avoided cost of lost nutrients and carbon due to erosion Increased soil nutrients over time Increased soil carbon and moisture Avoided cost of lime (reduced need for fertilizers and, therefore, reduced acidity risk)	Short term reduction in production due to nutrients and carbon retained in soil Costs of fertilizers and herbicides	Medium	Positive net benefits from increased farm productivity, profitability and sustainability mean these practices are being rapidly adopted in various regions of Australia (Sanderman <i>et al.</i> 2010)
Public	Landscape (soil) stabilisation Soil condition for crops	Reduced risk of erosion and downstream pollution Increased economic activity More stable farm profitability	Incentives for changed land management, where net private benefits are marginal	Medium	Positive regional and national net value due to avoided costs of erosion and pollution, greater agricultural economic activity, and possibly avoided costs of exceptional circumstances assistance.

Case study 2: Managing acid soils in broadacre cropping - Western Australia

Wheat production in Western Australia was worth \$1.8 billion in 2010, measured as gross value of production (ABS 2011a). This is equivalent to 38% of Australia's total crop. The largest wheat producing area is the Avon River Basin, which covers about 45% of the Western Australian wheatbelt (Gazey and Andrew 2010). Other significant areas are the northern and southern wheatbelts. Soil acidity is a significant constraint to increasing wheat yields.

Soil acidity reduces the ability of soil to provide nutrients and moisture for crop production. As Figure 9.2 shows, crop yields decline rapidly as the pH of surface soil drops below a target of 5.5⁸. For subsurface soils, a pH below 4.8 is a significant constraint to root growth. Acidic topsoils reduce the efficiency of nutrient use, leading to higher costs of fertilizers. Acidic subsurface soils can have toxic levels of aluminium which reduce crop root growth, leading to lower nutrient uptake, less efficient water use and lower crop yields.

Vulnerability to soil acidity is widespread in Western Australia's wheatbelt. In the Avon River Basin, almost 80% of topsoil samples are below a target pH of 5.5, while

⁸ pH is a measure of potential hydrogen. A higher pH indicates more basic conditions, while a lower pH indicates more acidic conditions.

50% of subsurface soil samples were below the target of 4.8 (Gazey and Andrew 2009). Similar results were found in the northern and southern wheatbelts, where more than 80% of topsoil samples were below a pH of 5.5. Coarse textured sands and gravels account for 90% of all soils affected. (Davies, Gazey, Bowden, *et al.* 2006).




Figure 9.2: Example of output from the acidity relative yield model for four plant tolerance classes within a given Al/Mn solubility class (Dolling *et al.* 2001)

Soil acidity can be reversed by adding lime to soils. However, if insufficient lime is added to agricultural soils at risk they gradually become more acidic. Failure to slow or reverse topsoil acidification generally leads to subsurface acidification. This is much more expensive to fix, and may need special equipment to inject lime deep into the subsurface soil (Davies, Gazey, and Tozer 2006).

Acid soils reduce wheat production in Western Australia by an estimated \$300-\$400 million per year (Gazey and Andrew 2010).⁹ The average loss in wheat yield is 8-12% (Davies, Gazey, and Tozer 2006). Grain yield responses to surface liming are often 10-15% and may increase with time. Subsurface liming can increase yields by 30-40% (Davies, Gazey, and Tozer 2006).

The public costs of acid soils may also be significant. There is speculation that acidification from agriculture might result in acid running off into local streams, with

⁹ It has been estimated that the overall cost of lost agricultural production from soil acidity in WA is around \$498 million/ year (Herbert 2009).

costs imposed on downstream water users, but evidence is not yet available (Cregan and Scott 1998, Hamblin 1996). Reduction in the quality and quantity of high value wheat grain also has negative impacts on local, regional, state and national economies. While there is little evidence linking off-farm impacts directly to agricultural practices (see Section 5.1), inaction on soil acidity does have public costs. Other public costs of soil acidification are longer term and associated with the risk of wind and water erosion on highly acidic soils which support little ground cover and the possibility of having to take land out of production because subsoil acidification is too costly to remediate.

Applying sufficient surface lime to treat acidity through the soil profile can be a cost-effective way to improve soil condition and the quality of ecosystem services it provides. Recent results from long-term field trials show that significant yield increases can be achieved in both the short and medium term, if sufficient lime is applied. Yield increases can be long-lasting, and may increase over time.

- A trial of 2 t/ha surface lime applied to sandy gravel at Bindi Bindi in the northern wheatbelt showed yield increases of over 10% within the first 2 years. Similar yield increases were still being achieved 8 years after lime was first applied. Net of amortised liming costs, grain income increased by \$87/ha (25%) in year 8 (Davies, Gazey and Tozer 2006).
- A trial of 5 t/ha surface lime, followed by a further 1 t/ha 10 years later showed wheat yields were 20% higher 17 years after the initial application. Surface lime was applied to a yellow sandy earth Tenosol at Kellerberrin in the Avon River Basin (Gazey and Andrew 2010).

In addition to improved yields, farmers can benefit from lower fertilizer use and a greater choice of crops to plant in rotations (Table 9.4). This flexibility can allow farmers to take advantage of volatility in international commodity prices, and better manage soil fertility by rotating crops. On-farm environmental benefits can include reduced weed growth, soil degradation and risk of wind erosion of soils (Davies, Gazey and Tozer, 2006; Gazey and Andrew 2010).

However, the amount of lime currently applied is not enough to adequately treat existing and on-going acidification in Western Australia (Hajkiewicz and Young 2002). Farmers often cite economic factors (upfront costs, returns and cash-flow constraints) as barriers to applying lime (Fisher *et al.* 2010). Yet focus groups suggest many farmers are convinced of the benefits of liming and need better information on how much lime to apply, and how to make it cost-effective. Others are

less convinced and have information needs for how liming works, what the benefits are, how much to apply, and the economics of liming (Fisher *et al.* 2010).

The key findings from this case study are:

- Private benefits of increased crop yields can be seen within 2 years. Gross margins (net of costs of lime) and yield gains are enduring and may increase over 10 to 15 years. A greater range of viable crop choices can allow farmers to better manage soil and respond to climate variability while taking advantage of fluctuating commodity prices.
- Public benefits may include reduced off-site environmental costs of water pollution, although this cannot be confirmed. Higher and more stable long-term economic contributions from the wheat industry may be another benefit, as are longer –term avoidance of soil erosion and loss of productive land.
- The private costs of inaction can rise significantly over time. Unless surface acidity is treated with enough lime, subsurface soil acidity can become an enduring constraint to cropping. Treatment of subsurface soil acidity is more expensive and technically difficult than applying surface lime.
- Barriers to private investment in improving soil condition may be lack of information on how, where and when to apply lime cost-effectively.
- Managing soil acidity by applying surface lime is relevant to around 80% of the West Australian wheat belt.

Table 9.4: Full range of benefits and beneficiaries – Managing acid soils in broadacre cropping

Beneficiaries	Ecosystem services	Benefits	Costs	Time-frame	Expected net value
Private land-holders	Soil condition for crops	Increased crop yields of >10% Reduced weed growth Wider range of choices for crop rotation Increased fertilizer use efficiency Increased water use efficiency	Purchase and application of lime to soil surface	Short – Medium	Positive and enduring if soil is tested regularly and sufficient lime applied. The longer-term net value should be compared to the cost of deep ripping of soil and injection of lime to reverse the sub-soil acidification that would occur if no action were taken.
Public	Soil condition for crops	Higher crop yields and greater choice of crop rotations increases and	Nil	Medium – Long	Positive

Beneficiaries	Ecosystem services	Benefits	Costs	Time-frame	Expected net value
		stabilizes regional and national economic contribution of agriculture Reduced offsite impacts of wind and water erosion, long term loss of land from production (intergenerational issue)			

Case study 3: Increasing soil carbon in irrigated horticulture – southern Australia

Horticulture is Australia's third largest agricultural industry, with an average \$8.4 billion annual gross value of production over 2008 to 2010 (ABS 2011a). Horticulture includes a diverse range of industries– with fruit and vegetables the largest product sectors (NLWRA 2008). The horticulture industry covers all states and a wide range of climate zones and types of soil. Irrigation is an important contributor to horticultural production, accounting for over 70% of the gross value of production in 2009-10 (ABS 2011b).

Under irrigation, Australian soils with poor structure can harden and significantly constrain horticultural production by restricting the growth of tree roots and their ability to take up water. In general, this is due to the age of Australian irrigated soils. Loams and fine sandy loams, in particular, may lack minerals that maintain soil porosity and structure under irrigation (Cockcroft 2012). Soils with high organic matter, are thought to be less likely to have this problem (Cockcroft and Olsson 2000).

This is a particular problem for commonly irrigated soils in Victoria's Goulburn Valley, the Murrumbidgee Irrigation Area in NSW and the Barossa Valley in South Australia. Red-brown earths account for a large amount of irrigated tree fruit, vines and vegetable production in southern Australia (Cockcroft 2012). These soils are vulnerable to hardening.

The opportunity cost of reduced crop yields due to poor soil condition may be high. An unpublished study suggests that for a fruit crop such as pears, Australian yields of 35 tonnes per hectare are well below the best international yield of 180 tonnes per hectare (Cockcroft 2012). Australian horticultural crops grown on poor soil types can average as low as 10 tonnes per hectare, while those grown on the best soils can achieve yields of 50 tonnes per hectare (Cockcroft 2012).

Increasing soil organic carbon in the root zone can significantly enhance agricultural productivity for a wide range of crops (Lal 2010). As shown in Table 9.5, increasing

soil carbon improves the quality of several final ecosystem services from soil: The increase in soil aggregation and available water capacity are among the important benefits of higher soil organic carbon (Lal 2010).

For horticulture, the conventional recommendation is to add organic carbon directly to the soil to supplement minimal till and controlled traffic techniques (Pattinson, *et al.* 2010; HAL 2010). Organic carbon may be in the form of manure, green waste or biochar.

An alternative approach of planting rye grass in fruit orchards has shown economic benefits in field trials in the Goulburn Valley. This method involves growing ryegrass in winter and mulching it onto the roots of trees in summer. The roots of ryegrass are thought to increase soil organic carbon by increasing biological activity within a sheath that protects organic matter from being consumed by worms and other soil biota (Cockcroft 2012). While this mechanism has not yet been fully studied, field results are promising. Preliminary trials suggest soil carbon and structure is reported to increase within a few months, although rye-grass may need to be planted two years ahead of fruit trees to get the best results.

Private benefits for farmers are primarily from higher fruit yields (Cockcroft 2012). In field trials since the 1980's, the best commercial yields have been double those achieved in 1965. However, it is important to note that factors other than increases in soil organic carbon and soil structure may be responsible for some of this increase¹⁰. Other benefits include trees with stronger and deeper root systems that should be more robust to a wide range of environmental pressures (Murray 2007). Farmers may also benefit from lower operating costs due to more efficient use of irrigation water, fertilizers and pesticides.

Private costs are relatively low, but do involve time and labour (Cockcroft 2012). For the best results, poor soils need to be planted with rye-grass for 2 years before planting trees. Orchards then need to be cultivated every 6 months to build up a bed of soil around the trees. To maximise increases in crop yield, changes to pruning practices and management of leaf to fruit ratio might also be required.

The public benefits from improved soil condition and higher quality ecosystem services are difficult to quantify. Reduced erosion of soil by water and/ or wind would be important if land cover were being increased towards 50%, but above this level of cover the likely off-site impacts on the public are likely to be small (Sections 6 and 7).

¹⁰ This does not account for any increases in crop yield due to other changes in orchard management systems since the mid-1960's. For example, the shift from flood irrigation to spray irrigation, building soil beds around trees, and loosening subsoil before planting fruit trees.

Other public benefits include the reduced pollution of streams and surface water due to greater water-use efficiency by fruit trees and increased removal of carbon from the atmosphere. If higher fruit yields lead to increased total gross value of production for the industry, there will also be public benefits that flow through greater local, regional and national economic activity. With the possible exception of investment in information for fruit growers to encourage building up soil carbon, public costs are nil or very low.

The key findings from this case study are:

- Private benefits of higher fruit yields are related to improved soil structure, nutrient and water conditions.
- Public benefits may include lower off-site environmental costs of water pollution, flowing from improved soil condition and more efficient use of water.
- There may be a lag of several years between action to improve soil condition and higher fruit yields. However, evidence of soil condition is visible within a few months.
- Barriers to private investment may be the availability of labour for orchard cultivation and new pruning practices.

Table 9.5: Full range of benefits and beneficiaries – Increasing soil carbon in irrigated horticulture

Beneficiaries	Final ecosystem services	Benefits	Costs	Timeframe	Expected net value
Private landholders	Landscape (soil) stabilisation Soil with nutrient and water conditions suitable for growing crops	Higher crop yields Increased efficiency of water-use and possibly fertilizer-use	Additional time and labour to plant and mulch ryegrass	Short – Medium	Positive. Field trials indicate fruit crops could be double those achieved with conventional orchard soil management systems not designed to build soil carbon.
Public	Landscape (soil) stabilisation Provision of clean water	Reduced erosion Reduced pollution of streams and surface water	Nil	Medium – Long	Positive

Case study 4: Reducing wind erosion in grazing areas - Rangelands

Australia's rangelands cover 81% of the continent (Bastin and ACRIS Management Committee 2008). They include a diverse range of relatively intact ecosystems, such

as tropical savannas, woodlands, shrublands and grasslands. Extensive grazing on native pastures takes place across the rangelands (Australian Government 2012). Much of Australia's \$7.4 billion/ year¹¹ beef grazing industry is located in the rangelands (ABS 2011a). Sheep grazing is also an important industry in some areas.

The causal links between over-grazing, loss of ground cover and soil degradation are well established. In one recent study, spatial comparisons of sites in semi-arid woodlands with different histories of grazing pressure demonstrated reductions in shrub cover and increases in bare soil at the most disturbed sites (Eldridge *et al.* 2011). Reduced soil stability and nutrient levels were obvious at the most disturbed sites, while sites with low levels of disturbance showed no physical or chemical degradation of their soils. Episodes of severe degradation occur when stocking rates remain high during droughts and ground cover declines due to over-grazing¹² (Stafford Smith *et al.* 2007). This decline is essentially permanent. While partial recovery of ground cover can occur during periods of higher rainfall, this requires even lower stocking rates than usual and may be unprofitable (Stafford Smith *et al.* 2007).

However, graziers lack visible signs to indicate when slowly declining soil condition may tip into irreversible degradation. Most of the time, soil condition declines slowly and may still support regrowth of perennial pastures (Ash *et al.* 2002). However, during episodes of drought the vulnerability of soil in poor condition becomes apparent. Impacts of severe soil degradation during drought condition include dust storms, erosion scalds and gullies (Stafford Smith *et al.* 2007). The gap in time between taking action to maintain soil condition and visible evidence of the avoided costs of erosion may be years or decades.

Severe soil degradation can impose significant private and public costs. Degraded soils lose the soil organic carbon, nutrients and structure needed to support perennial grasses on which stock graze (Ash *et al.* 2002). This has private costs, as long-term sustainable stocking rates may be reduced to as little as 40% of the average before degradation (Stafford Smith *et al.* 2007). The costs of rehabilitating land rise significantly for more extreme degradation, as grazing may not be possible for years while soil and pastures recover (Land & Water Australia 2005). Where soil condition is too poor to support the regrowth of perennial native grasses, even with

¹¹ This is the 2008– 2010 average gross value of agricultural product from livestock slaughtering of cattle and calves.

¹² A common pattern of decline in ground cover and soil condition following over-stocking during droughts has occurred in seven major episodes of land degradation since 1898 (Stafford Smith *et al.* 2007).

good rainfall, the land may need to be retired unless farmers can afford fertilizers to grow introduced pastures (Ash *et al.* 2002).

Several studies have estimated the off-site costs of dust storms in the order of millions of dollars. The 'Red Dawn' dust storm that hit Sydney in September 2009 is estimated to have cost over \$400 million in cleaning costs and lost work hours (Tozer 2012). This dust was lifted from the far west and northwest of NSW, and the Lake Eyre Basin, due to drought and extreme wind conditions (Leys *et al.* 2011; Tozer 2012). Earlier estimates of \$23 million per year for the cost of less severe dust storms in Adelaide included potential impacts on human respiratory health (Williams and Young 1999). Other public costs for which values have not been estimated include increased nutrient levels in waterways (Leys *et al.* 2011).

Keeping ground cover intact can reduce soil degradation and maintain forage for cattle (Stafford Smith *et al.* 2007). Reducing stocking rates to match pasture cover and condition is the main management practice to achieve this in the rangelands. A large decrease in the frequency of dust storms reaching east coast cities since the 1940's may be due to graziers monitoring ground cover levels in paddocks and setting targets for ground cover management¹³ (Australian State of the Environment Committee 2011; Barson *et al.* 2011). However, recent large dust storms, such as the 'Red Dawn' event in 2009, suggest Australia's management of ground cover is not yet sufficient to avoid wind erosion and soil degradation during extended droughts (Leys 2012).

Reducing stocking rates can provide net benefits to graziers. Although the net economic value may be small in the short-term, the longer-term economic benefits include more stable profits and reduced risk of negative cash returns (O'Reagain *et al.* 2011). Soil ecosystem services contribute to these benefits by providing conditions that allow a diverse range of perennial grasses to thrive (Table 9.6). The magnitude of both short-term benefits and the longer-term reduction in the risk of negative returns due to soil and pasture degradation will depend on the underlying soil type and condition.

More stable profits are likely to be the main private benefit of moderate stocking rates, where soil and pasture condition is not highly degraded (Table 9.6). This assumes the economics of rangelands grazing are similar to other areas of Australia. In northern tropical savanna regions, economic modelling based on the results of long-term field trials suggests that pastures maintained in good condition produce

¹³ In 2009-10, 79 % of grazing businesses monitor ground cover and 31 % set targets. Figures are for grazing businesses both within and outside the rangelands. It is worth noting that the number of businesses setting ground cover targets declined from 40 % in 2007 – 08 to 31 % in 2009-10.

slightly higher and more stable cash returns over 25 years than pasture in a deteriorated condition¹⁴ (Land & Water Australia 2005). However, returns were much higher and more consistent than for highly degraded pastures, which produced negative cash returns more than half the time.

By contrast, increasing stocking rates may provide marginal increases in profit, but reduce the resilience of native perennial pastures by driving declining soil condition. According to one economic modelling study in the rangelands for a typical 40,000 ha property in the Mitchell grass plains in Queensland and the Northern Territory, a 3% increase in cattle led to less than 1% profit increase and long term decline in soil condition and hydrology (Macleod and McIvor 2004).

The public benefits of avoiding episodic and ongoing erosion of bare soil are likely to be high (Table 9.6). These include reductions in the annual off-site cost of wind erosion to cities and regional towns, and potentially a lower risk of extreme dust storms. While there are currently no available economic estimates, the value of avoiding water erosion will depend on both management of critical areas of soil and the sensitivity of the catchment receiving sediment (Waddell et al. 2012; see also Box 4 in Section 7). If private profits are more stable, there may also be less need for publicly funded payments to farmers during droughts.

The key findings from this case study are:

- Private benefits of maintaining soil condition and ground cover are primarily more stable grazing profits over time. Using moderate stocking rates to achieve this may have a small positive or negative impact on profits in any given year.
- Public benefits are primarily lower off-site environmental, health and cleaning costs of dust due to wind erosion. On average, these could be worth tens of millions of dollars per year. In some years this may be hundreds of millions of dollars.
- Benefits occur within graziers' decision-making timeframes, but are only visible by comparison to poorly managed areas or during extended droughts.
- Barriers to managing ground cover levels are likely to be the lack of visible indicators of long-term benefits, as well as short-term financial pressure to increase stocking levels.

¹⁴ Cash returns were estimated as a net present value over 25 years using a 6 % discount rate.

- Managing ground cover levels to avoid erosion and maintain soil condition suitable for pastures is relevant to all of the rangelands grazing industry, which covers much of inland Australia.

Table 9.6: Full range of benefits and beneficiaries – Reducing wind erosion in grazing areas

Beneficiaries	Final ecosystem services	Benefits	Costs	Timeframe	Expected net value
Private landholders	Landscape (soil) stabilisation Soil condition for pasture	Avoided cost of rehabilitating or abandoning land Avoided cost of cattle feed during dry periods	Marginal reduction in profits due to lower stocking rate	Short – Medium	Positive due to more stable profits (assuming grazing operation not fully funded by equity, debt levels and interest costs would be lower if profits are stable)
Public	Landscape (soil) stabilisation Soil condition for pasture	Avoided costs of erosion Avoided public costs of drought impacts	Nil	Medium – Long	Positive due to avoided costs of land rehabilitation; health and other costs of erosion; and possibly lower publicly-funded payments to farmers during droughts

9.6 General findings

Australia's soils, their condition, land-use and management practices, are highly variable. This makes it difficult or impossible to present simple conclusions that apply to all soils in Australia.

However, we can draw some general conclusions from the case studies covered in this chapter.

1. The economic benefits from improving soil condition depend on the nature of the soil degradation process. There are three relevant factors:
 - How vulnerable is the soil to crossing a tipping point beyond which agricultural production is constrained?
 - Can the soil be returned to a condition that supports unconstrained agricultural production?
 - How much might this cost and how long will it take?

2. The soil conditions and threatening processes considered in this report vary widely in ways that impact the magnitude, timing and scale of the benefits of improving soil condition, and the costs of inaction:
 - Acidification may be reversible, although the cost of this can increase significantly past certain thresholds of soil condition. The most apparent costs of inaction are at the farm scale, although they may not be visible to land managers.
 - Some forms of soil organic carbon can be replenished within farm planning timeframes. Although the cost of doing so has not been established, private benefits at the farm scale may be visible to land managers where initial soil organic content is low.
 - Erosion causes a permanent loss of soil and associated nutrients that can impose long-term costs on land managers. However, these may not be obvious except during severe droughts or rainfall. Public costs can also be high due to the impact of dust storms.
3. Net private, and public benefits are positive and enduring for the land management practices covered in these case studies. In some cases the private economic benefits are more stable, rather than higher, profits.
4. However, other barriers to private investment may need to be addressed for these land management practices to be widely applied:
 - Conservation farming has already demonstrated improvement in broadacre crop yields in northern NSW. However, slow rates of adoption may have been due to low initial returns on investment.
 - Surface liming to manage acidity can increase and stabilise the profitability of wheat crops in Western Australia. However, land managers may need information about how, when and where to apply lime cost-effectively.
 - Increasing soil organic carbon can improve fruit yields for horticulture on red-brown earths in southern Australia. Barriers to private investment may be the availability of labour for orchard cultivation and new pruning practices.
 - Running moderate stocking rates can provide more stable long-term profits to grazing in the rangelands with impacts on average annual profits likely to be minimal. However, barriers to managing ground

cover levels are likely to be the lack of visible indicators of long-term benefits, as well as short-term financial pressure to increase stocking levels.

5. Over time, the ability to estimate public benefits should improve as data and knowledge about soils, land management practices and ecosystem processes develops. Some high-priority areas for research appear to be:
 - The impacts of soil acidity on surface and groundwater pollution
 - How to encourage strategic approaches to maintaining ground cover above critical thresholds (e.g., 50%) to reduce wind and water erosion from the rangelands during long droughts

10. Summary and conclusions

This project addressed two over-arching questions:

- What evidence exists about how improving land management practices will lead to reduced soil loss (through water and wind erosion) and improved soil condition (especially through reduced impacts of soil acidification and increased organic matter content)?
- How might reducing soil loss and improving soil condition result in improvements in the quantity and quality of ecosystem services and benefits delivered from agricultural lands, including cleaner air, improved water quality, reduced greenhouse gas emissions, and more productive soils?

The answers to these questions are summarised in the following sub-sections.

10.1 Improving the organic matter status of soils

Soil organic matter (SOC) contributes to a range of critical functions of soils, including: holding releasing plant nutrients; involvement in ion exchange; increasing soil water holding capacity; playing a role in building and maintaining soil structure and strength and reducing susceptibility to erosion; influencing water infiltration capacity surface runoff; providing a source energy for soil biota; buffering against fluctuations in soil acidity; and, moderation of soil temperature through its effect on soil colour and reflective capacity (Section 4).

These functions of SOC can be associated with provisioning, regulating and cultural ecosystem services as well as the soil processes that support these services.

The amount of SOC that accumulates is the balance between the amount of carbon added to the soil and the amount lost through degradation. Land-use change (including agriculture) has reduced SOC in many places around the world through both reductions in inputs and increases in losses. In Australia, clearing of native vegetation for primarily agricultural purposes has caused a 40-60% decrease in SOC stocks from pre-clearing levels.

Interpreting research on the effects of soil management practices on SOC is complicated because many studies have not been able to control all variables (e.g., rainfall, soil type, time since last cultivation, and the depth at which measurements are made all affect SOC accumulation). How sustained any increases might be is also subject to conjecture as there are limited long-term studies of these systems

across Australia, and rates of accumulation are highest in surface soils, which are also most vulnerable to disturbance.

There is good evidence that management of cropland to reduce disturbance, thereby reducing carbon losses, and increase carbon inputs (e.g., minimising tillage, retaining stubble, and/ or planting pastures between crops) has decreased rates of SOC loss compared with traditional practices, but has so far not resulted in absolute increases in SOC on average across Australia.

The greatest theoretical potential for building SOC is the addition of organic materials such as manure and green waste and the inclusion of a pasture phase in a cropping sequence, and/or transformation from cropping to permanent pasture and retirement and restoration of degraded land. Due to their relatively recent emergence there is very little scientific evidence that associates these sorts of carbon-enhancing practices with increased SOC in Australian broadacre cropping. There are likely to be some tradeoffs involved with such approaches, such as increased nutrient requirements for soil biota as their energy source is enhanced.

For horticulture, dairy and grazing industries, evidence of the efficacy of management strategies to increase SOC is difficult to find in the primary literature.

Horticulture in the past has often involved high losses of carbon to the atmosphere compared with other land uses. Like broadacre cropping, best-practice management of horticultural systems involves minimizing disturbance and compaction of soils (by machinery), maintaining ground cover, and improving inputs of carbon. Limited evidence suggests that these approaches are effective in managing soil carbon as they are for cropping.

Grazing by livestock (e.g. beef and sheep) can impact directly on SOC and nitrogen cycling by modifying plant biomass inputs into soil (shoot and root material) and by reducing ground cover and thereby exposure of SOC-rich surface layers to wind and water erosion, and can also impact indirectly by modifying soil structure. Management options to avoid and overcome these impacts have focussed on increasing carbon inputs (e.g., increasing productivity using irrigation and fertilization and addressing acidification) and reducing disturbance to soils and the potential for erosion (e.g., time controlled or rotational grazing and shifting to perennial pasture species). Research on the impacts of these options on SOC is limited, but a small number of studies in south-eastern Queensland and northern NSW have indicated short-term increases in herbage mass, SOC, nitrogen, and ground-litter, and reduced runoff and soil loss under time-controlled grazing compared to continuous grazing.

Dairy systems generally have high levels of SOC, due to high inputs of manure and fertilizers, but loss of soil carbon can occur and best-practice management seeks to minimize damage to soil from stock and loss of soil by erosion.

Sequestering carbon as way to reduce atmospheric carbon-dioxide is a somewhat separate issue to enhancing SOC to improve soil function. It appears that the potential for reduced or no-tillage (direct-drilling) and stubble-retention to sequester additional carbon and mitigate green house gas emission is limited in low-rainfall areas, in contrast to areas with higher rainfall and greater biomass production.

10.2 Improving the pH (acid-bases balance) of soils

There are several major causes for the acidification of agricultural soils, including: removal of agricultural products; excessive accumulation of organic matter; excessive use of nitrogenous fertilisers; and leaching of fixed, fertiliser and urine-N as nitrate from surface layers to lower layers before plants can utilise it. Impacts of soil acidification on-site and related to plant, animal and soil biological performance or off-site, though the link to stream and groundwater acidification is speculative (Section 5). On-site impacts include aluminium (Al) and manganese (Mn) toxicity affecting plants and plant nutritional problems caused by reduced availability of nutrients such as calcium (Ca), Magnesium (Mg), and Potassium (K). The resulting reduction in plant biomass production reduces the quantity and quality of plant residue entering soils and hence SOC levels.

Acidification occurs in both surface and subsurface soils, the latter being of increasing concern in parts of Australia (e.g., WA). Soil acidification is widespread in the extensive farming lands (cropping, sheep and cattle grazing) of southern Australia, and appears to be getting worse rather than better, and it is common in intensive systems of land use (tropical horticulture, sugar cane, dairying).

The use of high analysis nitrogen fertilisers and a high rate of product removal are features of most horticultural enterprises and about half of the horticultural industries have undertaken research to counter these potential problems. Due to diminishing returns from milk production dairy farmers nationally have intensified and diversified their production to remain profitable. This led to has increased stocking rates, use of irrigated annual fodder crops, use of mixed livestock systems of beef and dairy, and nutrient inputs, resulting in significant acidification, particularly in light textured soils where soil buffering capacity is low (e.g., in south-western Australia). Under grazed permanent pastures, nitrate leaching, as a result of over-fertilization and overstocking, is considered to be the largest risk in relation to acidification

Across most agricultural systems, the primary actions to address soil acidification are to: test soil pH regularly and at a range of depths; add lime at rates that are effective for arresting acidification; add lime at high rates, sufficient to reverse acidification in soils that have already acidified; use acid-tolerant plant species where available (as a short-medium term measure); and, retire land in the extreme. Management of potential acidity in many grazing systems consists of: sowing perennial grass species and/or agroforestry systems, to increase rooting depth and nitrate uptake; and reducing stocking rates on pastures with a high component of native grasses, to maintain vigour of the grasses.

Around 50% of dairy, broadacre cropping and horticulture businesses test for pH regularly (a slight decline over the past few years) and around 30% of grazing businesses (also a decline). Far fewer go on to apply lime or dolomite. It has been concluded that lime applications across Australia is far short of what is needed to arrest, let alone reverse, the rate of soil acidification. The use of acid tolerant species, although a relatively straightforward and cost-effective option, does not address the underlying problem, proving a temporary strategy for 'living with the problem' and probably making it worse.

There is compelling evidence to show that liming surface soils can increase yields of a wide variety of grasses and legumes (including many broadacre crops such as wheat and barley), so long as strategies are matched to soil type and pH, paddock variability, intended crops grown and fertiliser rates, and soil is tested regularly at a range of depths. These conclusions are based on intensive R&D effort in the 80s-90s on long-term trials in the high rainfall and temperate zones of southern Australia, and more recently in the 1990s-2000s in southern WA field trials. For broadacre cropping and high return industries such as horticulture and dairy, liming can be an effective and profitable management strategy for mitigating surface soil acidification provided appropriate rates are applied that account for regional and local (management) factors of soil and plant type and N-fertiliser regimes. The efficacy of practices to reduce subsoil acidification is less well established and only demonstrated on a small subset of soil types.

10.3 Minimising erosion of soils by wind

The extent to which soils are susceptible to wind erosion depends on a range of factors, including climatic variability, ground cover, topography, the nature and condition of the soil, and the energy of the wind. Land management can either moderate or accelerate wind erosion rates, largely depending on how it affects the proportion of bare soil, the dryness and looseness of the ground's surface, and

structures (stems, leaves, clumps of plants) that reduce the force of the wind. Grazing by stock, native animals (e.g., kangaroos) and feral animals (e.g., rabbits, camels, horses, goats) have major impacts on ground cover and soil physical properties. The changes in land cover brought about to establish much of Australia's agriculture have led to an increase in wind (and water) erosion.

The on-site impacts of wind erosion include soil loss, reduction in soil nutrients and organic matter (including soil organisms), release of soil carbon to the atmosphere, undesirable changes in soil structure, reduced water infiltration and moisture-holding capacity, and exposure of unproductive saline and acid subsoils. Off-site impacts include negative impacts on the global climate through positive radiative forcing of dust, physical impacts of dust storms on buildings and equipment, and health impacts of dust for people. The limited data available suggest that the off-site costs of wind erosion can be many times greater than the on-site costs. Historically, wind erosion has been particularly active in times of drought. In the 1940s and again in 2002 and 2009 there were heightened concerns due to dust storms hitting major Australian towns and cities.

Approaches to reducing wind erosion address three major aspects (Carter 2006): Ground cover; soil looseness; and, wind velocity. Ground cover is important as it reduces wind speed at the soil surface and captures soils particles mobilised by wind. Soil looseness increases when there is too little vegetation cover, soils are dry, the type of soil contains small particles and/ or the surface is smooth. Maintaining soil moisture, avoiding trampling of exposed soil by stock and maintaining rough soil surface are all ways to reduce soil looseness. While the velocity of wind is determined by the weather, it can be moderated locally by creating windbreaks.

Levels of combined water and wind erosion from cultivated land and rangelands are relatively similar, and as much as eight times greater than from uncultivated areas and forests. Management involves: protecting or encouraging ground cover, including avoidance of cultivation; control of pests that destroy ground cover and/or disturb the surface of soil; minimizing the area and intensity of grazing and cropping; and, managing movements of stock in dry areas using strategic placement of watering points.

Numerous studies have been performed in Australia, and in comparable ecosystems in other parts of the world, to show that increasing ground cover reduces losses of soil due to both wind and water erosion. As a general rule, it has been concluded that ground cover of around 50% is required to keep wind erosion to a minimum

across a range of climatic conditions and soil types (this level of cover achieves around an 80-90% reduction in erosion compared with bare soil).

The general relationships between ground cover and soil erosion have been known for over 20 years. The main focus of research and development during the past two decades has been on how to achieve ground cover cost-effectively.

Another line of evidence for the effectiveness of better management of ground cover and soil surface properties for reducing wind erosion comes from data showing that Dust Storm Indices (DSI) in the 1940s were on average four times higher in the 1940s than in the 2000s (management of ground cover has improved substantially since the 1940s). Despite these improvements, it is expected that the incidence of huge dust storms, like those in 2002, will increase in the future as parts of Australia go through long dry periods.

10.4 Minimising erosion of soils by water

Water erosion of soils occurs when soil particles are detached and carried away by water flowing across a landscape (Section 7). Like wind erosion, the on-site impacts of water erosion include soil loss, reduction in soil nutrients and organic matter (including soil organisms), release of soil carbon to the atmosphere, undesirable changes in soil structure, reduced water infiltration and moisture-holding capacity, and exposure of unproductive saline and acid subsoils. Off-site impacts include sedimentation of waterways and impacts on quality of surface water and groundwater (turbidity, nutrient and other chemical loads).

It is estimated that current rates of soil erosion by water across much of Australia exceed soil formation rates by a factor of at least several hundred and, in some areas, several thousand. While the time for total loss of soil is estimated to range from 100-500 or more years in different parts of Australia, it is expected that crops and other plants will respond to small changes in depth of topsoil, so that many areas are at risk of critical decline in productivity in much less than 100 years. Areas at highest risk include Coastal Queensland, the Wet Tropics, Mitchell Plains grasslands, New England Tablelands, and Victoria River basin in the Northern Territory.

Many of the effects of cultivation on susceptibility to wind erosion also apply to water erosion. Water erosion associated with cropping was recognised as a serious issue in the 1930s and has been a concern ever since. Horticulture faces many of the same risks of water erosion as broadacre cropping. Reduction of ground cover by livestock grazing can greatly increase vulnerability of landscapes to water erosion.

Key challenges for dairy enterprises include controlling sediment (along with nitrogen and phosphorus from fertilizers) losses into waterways, which can be exacerbated by compaction and disturbance of soil by the feet of grazing animals.

Land management practices designed to minimise water erosion seek to: increase ground cover above a critical threshold; minimise evaporation of soil moisture; maintain soil structure; limit compaction by heavy equipment or running of stock; and/ or, create of contours that control water flow.

There is an extensive literature showing that increasing ground cover reduces losses of soil due to water erosion. Typically, 20-30% cover reduces erosion by 80-90% across a range of soils and land uses. Ground cover can be grasses, herbs, trees, dead plants with root systems still intact, dead plant material (especially branches) lying on the surface, or even stones. While different combinations of cover-types have different effectiveness, In general, 70% ground cover is recommended to manage water erosion, although 80-100% cover is recommended where rainfall is moderate to high and slope are steep.

Reduced tillage has been shown to dramatically lower soil erosion and provide benefits for crop production and improved profits compared with traditional cultivation in a range of climates and soil types. This is especially true when economies of scale can be achieved by applying the same labour and machinery over large areas, and when controlled traffic management is used. Some limitations of conservation tillage have been identified, such as reduced surface roughness and enhanced run-off and sediment movement in areas where maintaining high biomass of plants is difficult or where low cover results from crop failure or grazing, but such issues can be managed cost-effectively.

In grazing systems, removal of stock has been shown to allow recovery of ground cover, if conditions are favourable for regrowth of pastures, but recovery of full soil functionality, especially organic matter content, can take years to decades.

10.5 improvements in the quantity and quality of ecosystem services and benefits delivered from agricultural lands

The living and non-living components of soil ecosystems interact to mediate a range of processes that would require engineering at an unprecedented scale to replicate (Section 8). These processes transform natural resources into forms that are potentially of benefit to humans and in so doing they are said to provide 'ecosystem services' (Table 10.1).

Table 10.1: Ecosystem services from soils and the benefits potentially derived (summarised from Section 8)

Ecosystem services		Potential benefits
Provisioning services		
Provision of fertile soil		Crops, meat, and other food
Support native pastures, foods, fibre, flowers and other above-ground natural raw materials		Natural products to support industries and lifestyles, bush food
Provision of natural products from soil		Natural products to support industries and lifestyles, food
Provision of clean water		Water of a quality suitable for drinking, recreation, use in industries, machinery etc.
Maintenance of genetic diversity		Intellectual stimulation, cultural value, moral value, potential for new foods and other products
Support for structures		Physical support for building and other infrastructure
Regulating services		
Water flow regulation		Protection from wind and water erosion and floods, prevention of salinity, storage of water
Maintenance of landscape (soil) stability		Protection from wind and water erosion, including risk to lives from land slippages, protection from damage and adverse health and climatic effects from dust storms
Regulation of atmospheric gases		A liveable atmosphere, physical and mental health and well being, liveable climate
Role (with vegetation) in regulation of weather and climate		A liveable climate
Breakdown of wastes and toxins		Disposal of wastes, health and wellbeing benefits
Regulation of species and populations in soils		Reduced risks of pests and diseases, reduced need for chemicals, health and financial benefits
Pollination and seed dispersal		Contributes to production of crops and native vegetation and the benefits that provides
Cultural/ habitat services		
Contributions to species, ecosystem and landscape diversity		Intellectual stimulation, knowledge, cultural and spiritual values (e.g., sense of place)

Management of land for agriculture dramatically changes the balance among ecosystem services, increasing some provisioning services, decreasing some regulating services and changing the nature of many cultural services. One aim of improved agricultural management is to adjust this balance to meet a wider range of private and public needs.

The research reviewed in this report has shown that best-practice approaches to managing soil carbon, acidity and wind and water erosion are generally effective at addressing those issues and improving soil condition generally. Practices like minimal tillage, maintaining ground cover above 50%, adding organic matter to soil

(within limits), and managing the impacts of stock and machinery on soil disturbance and compaction, have beneficial outcomes for all aspects of soil condition. These practices, therefore, potentially enhance most ecosystem services and allow most of the benefits that come from those services to be increased.

The beneficiaries include farmers, agricultural industries, communities, families and individuals in regional areas and in cities. It is possible to estimate the magnitude of these benefits under different conditions in the future, but it is not meaningful to make a single estimate of future value because of the many combinations of management practices, soil types, climatic variations, products, market opportunities, demographic changes, and demands of consumers over the coming decades. Some general conclusions can, however, be made:

- There are achievable opportunities to address declining soil carbon and increasing acidity and reduce wind and water erosion and at the same time improve profitability of agriculture and deliver a range of public benefits (which in some cases will be worth more than the private benefits in terms of health and wellbeing outcomes);
- To do this it will be important to consider the ability of soil ecosystems to cope with ongoing and potential future shocks (i.e., their adaptive capacity and resilience), which cannot be considered in isolation from the adaptive capacity and resilience of the humans who manage agricultural landscapes;
- The resilience of soils in many parts of Australia depends strongly on building and maintaining soil carbon stocks, which affect a wide range of functions, including nutrient cycling and water infiltration and storage, and the ability of landscapes to retain topsoil;
- Another key aspect of the resilience of Australian soils is their ability to avoid passing through thresholds of change, some of which could be, to all intents and purposes, irreversible;
- Such thresholds include critical proportions of ground cover (50-70% depending on factors like rainfall and slope), below which erosion accelerates dramatically, carbon-content thresholds, and thresholds of acidification, especially of subsoil, which currently cannot be addressed economically by most agricultural industries.

10.6 Summary


In summary, this report finds that:

- Efforts by farmers, graziers and governments since the 1970s have achieved many improvements in soil condition.
- Not only farmers, but all people stand to receive major financial and social gains from better management of soils.
- There is a great potential to achieve further benefits from improving soil condition and the quality of ecosystem services.
- Each of the four main indicators of soil condition—soil acidity, soil carbon, wind erosion and water erosion—can be improved by wider adoption of best-practice soil management.
- Appropriate practices do exist, and the benefits of greater adoption are significant to those involved in agriculture and to the wider public.

References

- Aarons S. R., Melland A. R. & Gourley C. J. P. (2004) Nutrient distribution within a dairy farm. In: SuperSoil 2004. 3rd Australian New Zealand Soils Conference, University of Sydney, Australia. p. Published on CDROM. <http://www.regional.org.au/au/pdf/asssi/supersoil2004/1622_aaronss.pdf>
- Abel N., Cork S., Gorddard R., Langridge J., Langston A., Plant R., Proctor W., Ryan P., Shelton D., Walker B. & Yialeloglou M. (2003) *Natural Values: Exploring Options For Enhancing Ecosystem Services In The Goulburn Broken Catchment*. CSIRO, Canberra.
- ABS. (2011a) Gross value of irrigated agricultural production, 2000-01 to 2009-10 (4610.0.55.008). Australian Bureau of Statistics, Canberra, <<http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4610.0.55.008Main+Features12000-01%20to%202009-10?OpenDocument>> [Accessed June 2, 2012].
- ABS. (2011b) Value of agricultural commodities produced, Australia, 2009-10 (7503.0). Australian Bureau of Statistics, Canberra, <<http://www.abs.gov.au/ausstats/abs@.nsf/mf/7503.0>> [Accessed June 2, 2012].
- Agtrans Research. (2007) An Economic Analysis of the GRDC Investment in Precision Agriculture Variable Rate Technology (VRT). Grains Research and Development Corporation, Canberra.
- Aitken R. L., Moody P. W. & McKinley P. G. (1990) Lime requirement of acidic Queensland soils 1. Relationships between soil properties and pH buffer capacity. *Australian Journal of Soil Research* **28**, 695-701.
- Andrew M. H. (1988) Grazing impact in relation to livestock watering points. *Trends in Ecology & Evolution* **3**, 336-9.
- Apitz S. E. (2012) Conceptualizing the role of sediment in sustaining ecosystem services: Sediment-ecosystem regional assessment (SECoRA). *The Science of the total environment* **415**, 9-30.
- Ash A., Corfield J. & Ksikisi T. (2002) The Ecograz Project : Developing Guidelines To Better Manage Grazing Country. Meat & Livestock Australia and the Queensland Government.
- Ashwood A., Kerr D. & Chataway R. (1993) Northern dairy feedbase 2001. 5. Integrated dairy farming systems for northern Australia. *Tropical Grasslands* **27**, 212-28.
- Australian Government. (2012) Rangelands Overview - Australia. Australian Natural Resources Atlas. <<http://www.anra.gov.au/topics/rangelands/overview/index.html>> [Accessed June 2, 2012].
- Australian State of the Environment Committee. (2011) Land. In: *State of the Environment 2011 (SoE 2011)* pp. 260-9. Australian Government, Canberra.
- AusVeg. (2010) EnviroVeg. AusVeg, Camberwell, Victoria, <<http://ausveg.com.au/enviroveg/programs.htm>> [Accessed 12 July, 2012].
- Baldock J., Grundy M., Wilson P., Jacquier D., Griffin T., Chapman G., Hall J., Maschmet D., Crawford D., Hill J. & Kidd D. (2009) Identification of Areas Within Australia with the Potential to Enhance Soil Carbon Content. CSIRO, Canberra.
- Baldock J. & Skjemstad J. (1999) Soil organic carbon/soil organic matter. In: *Soil Analysis: An Interpretation Manual* (eds K. Peverill, L. Sparrow and D. Reuter) pp. 159-70. CSIRO Publishing, Collingwood.
- Balmford A., Bruner A., Cooper P., Costanza R., Farber S., Green R., Jenkins M., Jefferiss P., Jessamay V., Madden J., Munro K., Myers N., Naeem S., Paavola J., Rayment M., S Rosendo, Roughgarden J., Trumper K. & Turner R. K. (2002) Economic reasons for conserving wild nature. *Science* **297**, 950-3.
- Bardgett R. D., Anderson J. M. & Behan-Pelletier V. (2001) The influence of soil biodiversity on hydrological pathways and the transfer of materials between terrestrial and aquatic ecosystems. *Ecosystems* **5**, 421-9.
- Barlow K., Nash D. & Hannah M. (2007) The effect of fertiliser and grazing on nitrogen export in surface runoff from rain-fed and irrigated pastures in south-eastern Australia. *Nutrient Cycling in Agroecosystems* **77**, 69-82.
- Barr N. F. & Cary J. W. (1992) *Greening a Brown Land*. Macmillan Education Australia, South Melbourne.
- Barrios E. (2007) Soil biota, ecosystem services and land productivity. *Ecological Economics* **64**, 269-85.
- Barson M. & Lesslie R. (2004) Land management practices – why they are important and how we know this. In: Land Management Practices Information Priorities, Classification and Mapping – Towards an Agreed National Approach, Canberra Winery, Canberra. Unpublished.
- Barson M., Mewett J. & Paplinska J. (2011) Land management practice trends in Australia's grazing (beef cattle/sheep)

- industries. Caring for our Country Sustainable Practices Fact Sheet 2. Department of Agriculture, Fisheries and Forestry, Canberra.
<http://www.daff.gov.au/__data/assets/pdf_file/0009/2148714/practices_grazing_fact_sheet.pdf>.
- Barson M., Mewett J. & Paplinska J. (2012a) Land management practice trends in Australia's dairy industry. Caring for our Country Sustainable Practices Fact Sheet 1. Department of Agriculture, Fisheries and Forestry, Canberra.
<http://www.daff.gov.au/__data/assets/pdf_file/0008/2148713/practices_dairy_fact_sheet.pdf>.
- Barson M., Mewett J. & Paplinska J. (2012b) Land management practice trends in Australia's broadacre cropping industries. Caring for our Country Sustainable Practices Fact Sheet 3. Department of Agriculture, Fisheries and Forestry, Canberra. <http://www.daff.gov.au/__data/assets/pdf_file/0006/2148711/Practices_-_Broadacre.pdf>.
- Barson M., Mewett J. & Paplinska J. (2012c) Land management practice trends in Australia's horticulture industry. Caring for our Country Sustainable Practices Fact Sheet 4. Department of Agriculture, Fisheries and Forestry, Canberra.
<http://www.daff.gov.au/__data/assets/pdf_file/0010/2148715/practices_horticulture_fact_sheet.pdf>.
- Bartley R., Roth C. H., Ludwig J., McJannet D., Liedloff A., Corfield J., Hawdon A. & Abbott B. (2006) Runoff and erosion from Australia's tropical semi-arid rangelands: Influence of ground cover for differing space and time scales. *Hydrological Processes* **20**, 3317-33.
- Basher L. R. & Lynn I. H. (1996) Soil changes associated with cessation of sheep grazing in the Canterbury high country, New Zealand. *New Zealand Journal of Ecology* **20**, 179-89.
- Bastin G. & ACRIS Management Committee. (2008) Rangelands 2008 - Taking the Pulse. National Land and Water Resources Audit, Canberra.
- Beadle N. C. W. (1948) *The Vegetation and Pastures of Western New South Wales with Special Reference to Soil Erosion*. T.H. Tennant, Government Printer, Sydney.
- Beeston G., Stephens D., Nunweek M., Walcott J. & Ranatunga K. (2005) GRDC strategic planning for investment based on agro-ecological zones. Bureau of Rural Sciences, Canberra.
- Belnap J. & Gillette D. A. (1998) Vulnerability of desert biological soil crusts to wind erosion: the influences of crust development, soil texture, and disturbance. *Journal of arid environments* **39**, 133-42.
- Bennett J. (1999) Estimating the Values of Environmental Impacts of Agriculture. In: Country Matters, Canberra. Bureau of Rural Sciences, Canberra.
<http://live.greeningaustralia.org.au/nativevegetation/pages/pdf/Authors%20B/7_Bennett.pdf>.
- Bennett L. T., Mele P. M., Annett S. & Kasel S. (2010) Examining links between soil management, soil health, and public benefits in agricultural landscapes: An Australian perspective. *Agriculture, Ecosystems & Environment* **139**, 1-12.
- Binning C., Cork S., Parry R. & Shelton D. (2001) Natural assets: An inventory of ecosystem goods and services in the Goulburn Broken catchment. CSIRO, Canberra.
- Bockstael N. E., Freeman A. M., Kopp R. J., Portney P. R. & Smith V. K. (2000) On measuring economic values for nature. *Environmental Science & Technology* **34**, 1384-9.
- Bolland M. D. A. & Russell W. K. (2010) Changes in chemical properties of 48 intensively grazed, rain-fed dairy paddocks on sandy soils over 11 years of liming in south-western Australia. *Soil Research* **48**, 682-92.
- Botton S., van Heusden M., Parsons J. R., Smidt H. & van Straalen N. (2006) Resilience of microbial systems towards disturbances. *Critical reviews in microbiology* **32**, 101-12.
- Bowman K. (2008) Economic and Environmental Analysis of Converting to Controlled Traffic Farming. In: 6th Australian Controlled Traffic Conference pp. 61-8. <www.ctfsolutions.com.au/ebm3-doc/.../Bowman%20Economics.pdf>.
- Boyd J. & Banzhaf S. (2007) What are ecosystem services? The need for standardized environmental accounting units. *Ecological Economics* **63**, 616-26.
- Brand F. S. & Jax K. (2007) Focusing the meaning(s) of resilience: Resilience as a descriptive concept and a boundary object. *Ecology and Society* **12**, [on line] <http://www.ecologyandsociety.org/vol12/iss1/art32/ES-2007-134.pdf>.
- Brandt J. (1988) The transformation of rainfall energy by a tropical rain forest canopy in relation to soil erosion. *Journal of Biogeography* **15**, 41-8.
- Braunack M. V. & Walker J. (1985) Recovery of some surface soil properties of ecological interest after sheep grazing in a semi-arid woodland. *Austral Ecology* **10**, 451-60.

- Bruce S. E., Howden S. M., Graham S., Seis C., Ash J. & Nicholls A. (2006) Pasture-cropping: effect on biomass, total cover, soil water and nitrogen. Pasture Cropping & No Kill Cropping, Wellington, NSW, <http://www.nokillcropping.com/index.php?option=com_content&view=article&id=50:pasture-cropping-effect-on-biomass-total-cover-soil-water-a-nitrogen&catid=40:research-findings&Itemid=63> [Accessed June 15, 2012].
- Buckerfield J. C. (1992) Earthworm populations in dryland cropping soils under conservation-tillage in south australia. *Soil Biology and Biochemistry* **24**, 1667-72.
- Bui E. N., Hancock G. J., Chappell A. & Gregory L. J. (2010) Evaluation of Tolerable Erosion Rates and Time to Critical Topsoil Loss in Australia. CSIRO, Canberra.
- Campbell A. (2008) Managing Australia's Soils: A Policy Discussion Paper. Department of Agriculture, Fisheries and Forestry, Canberra.
- Carey B. W., Leach G. J. & Venz B. (2004) Soil erosion in Queensland cropping lands – an historical perspective and new challenges. In: Conserving Soil and Water for Society: Sharing Solutions. Proceedings of the 13th International Soil Conservation Organisation Conference (eds S. R. Raine, A. J. W. Biggs, N. W. Menzies, D. M. Freebairn and P. E. Tolmie) p. 8. Australian Society of Soil Science Incorporated/International Erosion Control Association (Australasia), Brisbane.
- Caring for our Country. (2012) Managing Soil Acidity on WA wheat belt farms. <www.nrm.com.au>.
- Carpenter S. R., Walker B. H., Anderies J. M. & Abel N. (2001) From metaphor to measurement: Resilience of what to what? *Ecosystems* **4**, 765-81.
- Carr S., York D., Andrew J. & Gazey C. (2006) Economic returns from monitoring and remediation of acid soils with lime application. Australian Government, Canberra, <http://www.soilhealthknowledge.com.au/images/PDFfiles/acidity_case_study_2006.pdf>.
- Carroll C., Merton L. & Burger P. (2000) Impact of vegetative cover and slope on runoff, erosion, and water quality for field plots on a range of soil and spoil materials on central Queensland coal mines. *Australian Journal of Soil Research* **38**, 313-28.
- Carter D. (2002) The amount of stubble needed to reduce wind erosion. Farmnote no. 67/2002. Department of Agriculture Western Australia, Perth. <http://www.agric.wa.gov.au/objtwr/imported_assets/content/lwe/land/erosion/fn067_2002.pdf>.
- Carter D. (2006)  Wind erosion - prevention and management. Department of Agriculture and Food, Western Australia, Perth, WA, <http://www.agric.wa.gov.au/PC_92240.html> [Accessed 28 June, 2012].
- Carter M. & Steed G. (1992) The effects of direct drilling and stubble retention on hydraulic properties at the surface of duplex soils in north-eastern Victoria. *Soil Research* **30**, 505-16.
- Carter M. R., Parton W. J., Rowland I. C., Schultz J. E. & Steed G. R. (1993) Simulation of soil organic carbon and nitrogen changes in cereal and pasture systems of southern Australia. *Australian Journal of Soil Research* **31**, 481-91.
- Central West Catchment Management Authority. (2008) Perennial Pastures on the Central West Plains. A Best Management Practice Guide for the Central West Catchment. NSW Government and the Central West Catchment Management Authority. <cw.cma.nsw.gov.au/_literature_37462/Plains>.
- Chadwick O. A. & Chorover J. (2001) The chemistry of pedogenic thresholds. *Geoderma* **100**, 321-53.
- Chan K. Y. (2007) Soil compaction controls the abundance, biomass and distribution of earthworms in a single dairy farm in south-eastern Australia. *Soil & Tillage Research* **94**, 75-82.
- Chan K. Y., Heenan D. P. & So H. B. (2003) Sequestration of carbon and changes in soil quality under conservation tillage on light-textured soils in Australia: a review. *Australian Journal of Experimental Agriculture* **43**, 325-34.
- Chan Y. (2008) Increasing soil organic carbon of agricultural land. Primefact 732. NSW Department of Primary Industries, Wagga Wagga.
- Chaudri A., McGrath S., Gibbs P., Chambers B., Carlton-Smith C., Bacon J., Campbell C. & Aitken M. (2008) Population size of indigenous *Rhizobium leguminosarum* biovar trifolii in long-term field experiments with sewage sludge cake, metal-amended liquid sludge or metal salts: Effects of zinc, copper and cadmium. *Soil Biology & Biochemistry* **40**, 1670–80.
- Clothier B. E., Hall A. J., Deurer M., Green S. R. & Mackay A. D. (2011) 9. Soil ecosystem services - sustaining returns on investment into natural capital. In: *Sustaining Soil Productivity in Response to Global Climate Change: Science,*

- Policy, and Ethics* (eds T. J. Sauer, J. M. Norman and M. V. K. Sivakumar) pp. 117-39. John Wiley & Sons, Inc.
- Cockcroft B. (2012) Soil Management For Australian Irrigated Horticulture. National Program for Sustainable Irrigation, Canberra, <<http://npsi.gov.au/files/products/national-program-sustainable-irrigation/npsi0212/npsi0212-soil-management-australian-irrigated-hort.pdf>> [Accessed June 2, 2012].
- Cockcroft B. & Olsson K. A. (2000) Degradation of soil structure due to coalescence of aggregates in no-till, no-traffic beds in irrigated crops. *Australian Journal of Soil Research* **38**, 61-70.
- Coles N. & Moore G. (2001) 7.2 Runoff and water erosion. In: *Soilguide: A Handbook for Understanding and Managing Agricultural Soils. Bulletin 4343* (ed G. Moore) pp. 223-42. Agriculture Western Australia, Perth, WA.
- Colloff M. J., Fokstuen G. & Boland T. (2003) Toward the triple bottom line in sustainable horticulture: Biodiversity, ecosystem services and an environmental management system for citrus orchards in the riverland of South Australia. CSIRO, Canberra. <<http://www.ecosystemservicesproject.org/html/publications/index.htm>>.
- Colloff M. J., Pullen K. R. & Cunningham S. A. (2010) Restoration of an Ecosystem Function to Revegetation Communities: The Role of Invertebrate Macropores in Enhancing Soil Water Infiltration. *Restoration Ecology* **18**, 65-72.
- Cork S. (2010a) *Resilience and Transformation - Preparing Australia for Uncertain Futures*. CSIRO Publishing, Collingwood, Victoria, Australia.
- Cork S. (2010b) Ways forward in the population and environment debate. Parliament of Australia, Department of Parliamentary Services, Canberra, <<http://www.aph.gov.au/library/pubs/PEPU/PopulationEnvironDebate.pdf>> [Accessed 6 May 2011, 2011].
- Costanza R. (2008) Ecosystem services: multiple classification systems are needed. *Biological Conservation* **141**, 350-2.
- Costanza R., Ayres R. U., Daly H. E., El Serafy S., Herendeen R. A., Hueting R., Norgaard R. B., Opschoor J. B., Pimentel D., Rees W. E., Templet P. H., Toman M. & Turner R. K. (1998) Special section: Forum on valuation of ecosystem services. *Ecological Economics (Amsterdam)* **25**, 1-72.
- Costanza R., d'Arge R., Groot R., Farber S., Grasso M., Hannon B., Limburg K., Naeem S., O'Neill R. V., Paruelo J., Raskin G., Sutton P. & Vandenbelt M. (1997) The value of the world's ecosystem services and natural capital. *Nature (London)* **387**, 253-60.
- Coventry D. R., Reeves T. G., Brooke H. D., Ellington A. & Slattery W. J. (1987) Increasing wheat yields in north-eastern Victoria by liming and deep ripping. *Australian Journal of Experimental Agriculture* **27**, 679-85.
- Coventry D. R., Walker B. R., Morrison G. R., Hyland M. T., Avery J. C., Maden J. J. L. & Bartram D. C. (1989) Yield responses to lime of wheat and barley on acid soils in north-eastern Victoria. *Australian Journal of Experimental Agriculture* **29**, 209-14.
- Cregan P. D. & Scott B. J. (1998) Soil acidification-an agricultural and environmental problem. . In: *Agriculture and the Environmental Imperative* (eds J. Pratley and A. Robertson). CSIRO Publishing, Collingwood.
- Crosthwaite J., Malcolm B., Moll J. & Dorrough J. (2008) Future investment in landscape change in southern Australia. *Landscape Research* **33**, 225-39.
- CSIRO. (2012) Assessment of the ecological and economic benefits of environmental water in the Murray–Darling Basin. The final report to the Murray–Darling Basin Authority from the CSIRO Multiple Benefits of the Basin Plan Project. CSIRO, Australia.
- D'Emden F. H. & Llewellyn R. S. (2006) No-till adoption decisions in southern Australian cropping and the role of weed management. *Australian Journal of Experimental Agriculture* **46**, 563-9.
- DAFF. (2012a) Australia's agriculture, fisheries and forestry at a glance. Department of Agriculture, Fisheries and Forestry, Canberra, <<http://www.daff.gov.au/about/publications/glance2012>>.
- DAFF. (2012b) Australian horticulture fact sheet. Department of Agriculture Fisheries and Forestry, Canberra, <http://www.daff.gov.au/__data/assets/pdf_file/0020/2109206/Horticulture-fact-sheet-January-2012.pdf> [Accessed 11 July, 2012].
- DAFWA. (2012) Soil acidity publications. Department of Agriculture and Food WA, Perth, <http://www.agric.wa.gov.au/PC_92430.html> [Accessed 1August, 2012].
- Daily G. C. (1997) *Nature's Services - Societal Dependence on Natural Ecosystems*. Island Press, Washington.

- Daily G. C. (1999) Developing a Scientific Basis for Managing Earth's Life Support Systems. *Conservation Ecology* **3**, [on line] <http://www.consecol.org/vol3/iss2/art14>.
- Daily G. C., Matson P. A., Vitousek P. M. & Daily G. E. (1997) Ecosystem services supplied by soil. In: *Nature's Services* (ed G. C. Daily) pp. 113-32. Island Press, Washington.
- Dairy Australia. (2012) The Australian dairy industry. <<http://www.dairyaustralia.com.au/Industry-overview/About-the-industry.aspx>>.
- Dale V. H. & Polasky S. (2007) Measures of the effects of agricultural practices on ecosystem services. *Ecological Economics* **64**, 286-96.
- Davies S., Gazey C., Bowden B., Gool D. v., Gartner D., Liaghati T. & Gilkes B. (2006) Acidification of Western Australia's agricultural soils and their management. In: Groundbreaking Stuff. Proceedings of the 13th Australian Agronomy Conference. Perth, Western Australia (eds N. Turner and T. Acuna) p. [on line]. The Regional Institute Ltd. <http://www.regional.org.au/au/asa/2006/concurrent/soil/4555_daviess.htm>.
- Davies S., Gazey C. & Tozer P. (2006) Long-term productivity and economic benefits of subsurface acidity management from surface and subsurface liming. In: Agribusiness Crop Updates 2006. Perth, Western Australia pp. 15-6. Department of Agriculture and Food and the Grains Research and Development Corporation. <http://www.agric.wa.gov.au/objtwr/imported_assets/content/fcp/farmingsystems2006_1.pdf>.
- Day P., Cribb J., Hannam R., Price P., Krause M., Wylie P., Burgi A. & Riches D. (2008) Managing Complex Systems : Preliminary findings from Grain & Graze. Land & Water Australia, Canberra.
- De Groot R., Van der Perk J., Chiesura A. & van Vliet A. (2003) Importance and threat as determining factors for criticality of natural capital. *Ecological Economics* **44**, 187-204.
- De Groot R. S., Alkemade R., Braat L., Hein L. & Willemen L. (2010) Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecological Complexity* **7**, 260-72.
- de Groot R. S., Wilson M. A. & Boumans R. M. J. (2002) A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecological Economics* **41**, 393-408.
- DEHCD. (1978) A basis for soil conservation policy in Australia: Commonwealth and State Government Collaborative Soil Conservation Study - Report 1. Department of Environment Housing and Community Development, Canberra.
- DEHNSW. (2012) DustWatch. Department of Environment and Heritage, NSW, Sydney, <<http://www.environment.nsw.gov.au/dustwatch/>> [Accessed 1 August, 2012].
- DEWHA. (2009) Assessment of Australia's Terrestrial Biodiversity 2008. W. Australian Government Department of the Environment, Heritage and the Arts, Canberra.
- Dewhurst S. & Lindsay C. (1999) Development of good farm management guidelines for sustainable fruit and vegetable production. Horticultural Research & Development Corporation and the Queensland Fruit and Vegetable Growers, Sydney. <http://books.google.com.au/books?id=xDsnAQAAACAAJ&dq=intitle:Development+of+good+farm+management+guidelines+for+sustainable+fruit+and+vegetable+production&cd=1&source=gbs_api>.
- Dolling P., Ferris D. & Fedorenko D. (2010) Initial farmer perceptions on pasture cropping in the Northern Agricultural Region of Western Australia. Future Farm Industries CRC, Perth.
- Dolling P. J., Moody P. W., Noble A. D., Helyar K. R., Hughes B., Reuter D. J. & Sparrow L. (2001) Soil Acidity and Acidification. National Land and Water Resources Audit, Canberra.
- Dominati E., Patterson M. & Mackay A. (2010) A framework for classifying and quantifying the natural capital and ecosystem services of soils. *Ecological Economics* **69**, 1858-68.
- Doran J. W. & Zeiss M. R. (2000) Soil health and sustainability: managing the biotic component of soil quality. *Applied Soil Ecology* **15**, 3-11.
- Dorrough J., Yen A., Turner V. & Clark S. G. (2004) Livestock grazing management and biodiversity conservation in Australian temperate grassy landscapes. *Australian Journal of Agricultural Research* **55**, 279-95.
- Douglas A. & Peltzer S. C. (2004) Managing herbicide resistant annual ryegrass (*Lolium rigidum* Gaud.) in no-till systems in Western Australia using occasional inversion ploughing. In: Proceedings of the Fourteenth Australian Weeds Conference. Council of Australasian Weed Societies (CAWS).

- <http://www.caws.org.au/awc/2004/awc200413001.pdf>.
- Duncan M. (1999) Pastures and acid soils. NSW Agriculture, Armidale.
http://www.dpi.nsw.gov.au/__data/assets/pdf_file/0005/167261/acid-soil-pastures.pdf.
- Durán Zuazo V. H., Martínez J. R. F., Pleguezuelo C. R. R., Martínez Raya A. & Rodríguez B. C. (2006) Soil-erosion and runoff prevention by plant covers in a mountainous area (SE Spain): implications for sustainable agriculture. *The Environmentalist* **26**, 309–19.
- Durán Zuazo V. H. D., Martínez J. R. F. & Raya A. M. (2004) Impact of vegetative cover on runoff and soil erosion at Hillslope Scale in Lanjaron, Spain. *The Environmentalist* **24**, 39–48.
- Earl J. M. & Jones C. E. (1996) The need for a new approach to grazing management - Is cell grazing the answer? *The Rangeland Journal* **18**, 327-50.
- Edwards K. (1993) Soil erosion and conservation in Australia. In: *World soil erosion and conservation* (ed D. Pimentel) pp. 147-70. Cambridge University Press, Cambridge, UK.
- Eldridge D. J. (1993) Cryptogams, vascular plants, and soil hydrological relations: some preliminary results from the semiarid woodlands of eastern Australia. *Western North American Naturalist* **53**, 48-58.
- Eldridge D. J. (1998) Trampling of microphytic crusts on calcareous soils, and its impact on erosion under rain-impacted flow. *Catena* **33**, 221-39.
- Eldridge D. J. & Greene R. S. B. (1994) Microbiotic soil crusts-a review of their roles in soil and ecological processes in the rangelands of Australia. *Soil Research* **32**, 389-415.
- Eldridge D. J. & Leys J. F. (2003) Exploring some relationships between biological soil crusts, soil aggregation and wind erosion. *Journal of arid environments* **53**, 457-66.
- Eldridge D. J., Val J. & James A. I. (2011) Abiotic effects predominate under prolonged livestock-induced disturbance. *Austral Ecology* **36**, 367–77.
- ERIN. (2009a) Indicative locations where improving soil and land management practices to reduce soil loss from wind erosion will provide the biggest benefits (map). Environmental Resources Information Network Canberra, <http://nrmonline.nrm.gov.au/catalog/mql:2348> [Accessed June 2, 2012].
- ERIN. (2009b) Indicative locations where improving soil and land management practices to reduce soil loss from hillslope (sheet and rill) erosion will provide the biggest benefits (map). Environmental Resources Information Network, Canberra, <http://nrmonline.nrm.gov.au/catalog/mql:2386> [Accessed June 2, 2012].
- ERIN. (2009c) Indicative locations where improving soil and land management practices to increase soil organic matter will provide the biggest benefits (map). Environmental Resources Information Network, Canberra, <http://nrmonline.nrm.gov.au/catalog/mql:2390> [Accessed June 2, 2012].
- Erskine W. D., Mahmoudzadeh A. & Myers C. (2002) Land use effects on sediment yields and soil loss rates in small basins of Triassic sandstone near Sydney, NSW, Australia. *Catena* **49**, 271-87.
- Erskine W. D. & Saynor M. J. (1996) Success of soil conservation works in reducing soil erosion rates and sediment yields in central eastern Australia. In: *Erosion and Sediment Yield: Global and Regional Perspectives*. Proceedings of the Exeter Symposium, July 1996. IAHS Publ. no. 236 pp. 523-30. IAHS.
https://itia.ntua.gr/hsj/redbooks/236/iahs_236_0523.pdf.
- Evans R. (1998) The erosional impacts of grazing animals. *Progress in Physical Geography* **22**, 251-68.
- Felton W. L., Marcellos H. & Martin R. J. (1995) A comparison of three fallow management strategies for the long-term productivity of wheat in northern New South Wales. *Australian Journal of Experimental Agriculture* **35**, 915–21.
- Findlater P. A., Carter D. J. & Scott W. D. (1990) A model to predict the effects of prostrate ground cover on wind erosion. *Australian Journal of Soil Research* **28**, 609-22.
- Firth D. J., McFadyen L. M., Jones R. M. & Cook B. G. (1999) Evaluation of groundcovers for macadamia orchards. Horticultural Research and Development Corporation, Sydney.
http://scholar.google.com/scholar?q=related:GGQIMJzHwnsJ:scholar.google.com/&hl=en&num=30&as_sdt=0,5.
- Fischer J., Stott J., Zenger A., Warren G., Sherren K. & Forrester R. I. (2009) Reversing a tree regeneration crisis in an endangered ecoregion. *Proceedings of the National Academy of Sciences of the United States of America* **106**,



10386-91.

- Fisher B., Turner K., Zylstra M. & Brouwer R. (2008) Ecosystem services and economic theory: integration for policy-relevant research. *Ecological Applications* **18**, 2050-67.
- Fisher B., Turner R. & Morling P. (2009) Defining and classifying ecosystem services for decision making. *Ecological Economics* **68**, 643-53.
- Fisher J., Gazey, C., Andrew, J., Carr, S. (2010) Two levels of information are required to assist extension of liming in Western Australia. In: Food Security from Sustainable Agriculture. Proceedings of 15th Agronomy Conference, Lincoln, New Zealand (eds H. Dove and R. A. Culvenor) p. [on line]. The Regional Institute Ltd. <http://www.regional.org.au/au/asa/2010/farming-systems/participatory-research-extension/7121_fisherjs.htm#TopOfPage.
- Fleming N. (1998) Phosphorus and carbon losses off dairy catchments located on duplex soils. In: Agronomy, Growing a Greener Future? Proceedings of the 9th Australian Agronomy Conference, Charles Sturt University, Wagga Wagga, NSW. (eds D. L. Michalk and J. E. Pratley) p. [on line]. The Regional Institute Ltd. <<http://www.regional.org.au/au/asa/1998/8/021fleming.htm>.
- Fleming N. K., Cox J. W., Chittleborough D. J. & Dyson C. B. (2001) An analysis of chemical loads and forms in overland flow from dairy pasture in South Australia. *Hydrological Processes* **15**, 393-405.
- Fleming N. K., Cox J. W., Fleming N. K. & Cox J. W. (2001) Carbon and phosphorus losses from dairy pasture in South Australia. *Australian Journal of Soil Research* **39**, 969-78.
- Foley J. A., DeFries R., Asner G. P., Barford C., Bonan G., Carpenter S. R., Chapin F. S., Coe M. T., Daily G. C., Gibbs H. K., Helkowski J. H., Holloway T., Howard E. A., Kucharik C. J., Monfreda C., Patz J. A., Prentice I. C., Ramankutty N. & Snyder P. K. (2005) Global consequences of land use. *Science* **309**, 570-4.
- Folke C., Carpenter S., Elmqvist T., Gunderson L., Holling C. & Walker B. (2002) Resilience and sustainable development: Building adaptive capacity in a world of transformations. *Ambio* **31**, 437-40.
- Folke C., Carpenter S., Walker B., Scheffer M., Elmqvist T., Gunderson L. & Holling C. S. (2004) Regime shifts, resilience and biodiversity in ecosystem management. *Annual Review of Ecology Evolution and Systematics* **35**, 557-81.
- Folke C., Jansson L., Larsson J. & Costanza R. (1997) Ecosystem appropriation by cities. *Ambio* **26**, 167-72.
- Freebairn D. (2004) Some observations on the role of soil conservation structures and conservation. *Journal of the Australian Association of Natural Resource Management* **7**, 8-13.
- Freebairn D., Ward L., Clarke A. & Smith G. (1986) Research and development of reduced tillage systems for Vertisols in Queensland, Australia. *Soil & Tillage Research* **8**, 211-29.
- Freebairn D. M. (1992) Managing Resources-the soil resource: erosion, stubble management and catchment. In: Looking Back - Planning Ahead. Proceedings of the 6th Australian Agronomy Conference, The University of New England, Armidale, New South Wales (eds K. Hutchinson and P. Vickery) p. [on line]. The Regional Institute Ltd. <<http://regional.org.au/au/asa/1992/invited/p-05.htm?print=1>.
- Freebairn D. M., Loch R. J. & Cogle A. L. (1993) Tillage methods and soil and water conservation in Australia. *Soil & Tillage Research* **27**, 303-25.
- Freebairn D. M. & Silburn D. M. (2004) Soil conservation in Australia's semi arid tropics: Pathways to success, and new challenges. In: Conserving Soil and Water for Society: Sharing Solutions. ISCO 2004 - 13th International Soil Conservation Organisation Conference, Brisbane (eds S. R. Raine, A. J. W. Biggs, N. W. Menzies, D. M. Freebairn and P. E. Tolmie) p. [on line]. ASSSI/IECA, Brisbane, Qld). <<http://tucson.ars.ag.gov/isco/isco13/PAPERS%20F-L/FREEBAIRN.pdf>.
- Freebairn D. M. & Wockner G. H. (1986) A study of soil erosion on vertisols of the eastern Darling Downs, Queensland. I. Effects of surface conditions on soil movement within contour bay catchments. *Australian Journal of Soil Research* **24**, 135-58.
- Freebairn D. M., Wockner G. H., Hamilton N. A. & Rowland P. (2009) Impact of soil conditions on hydrology and water quality for a brown clay in the north-eastern cereal zone of Australia. *Australian Journal of Soil Research* **47**, 389-402.
- Gaffney J. & Wilson A. (2003) The economics of zero tillage and controlled traffic farming for Western Downs farms. In: Soil Management for Sustainability. Proceedings of International Soil Tillage Research Organisation 16th Triennial Conference pp. 458-64. International Soil Tillage Research Organisation.

- Gao Y., Zhong B., Yue H., Wu B. & Cao S. (2011) A degradation threshold for irreversible loss of soil productivity: A long-term case study in China. *Journal of Applied Ecology* **48**, 1145-54.
- Gazey C. (2008) Case Study 3. Farmer experience. In: *Soil Health Knowledge Bank: an Australian Government Initiative*, Canberra.
- Gazey C. & Andrew J. (2009) Soil pH in northern and southern areas of the WA wheatbelt. Department of Agriculture and Food, Western Australia, Perth.
<http://www.agric.wa.gov.au/objtwr/imported_assets/content/lwe/land/acid/north_and_south_ph_report_151208.pdf>.
- Gazey C. & Andrew J. (2010) Getting the soil pH profile right helps with weed control and sustainability. In: *Soil Solutions for a Changing World*. 9th World Congress of Soil Science pp. 232–5, Brisbane.
<<http://www.idd.go.th/swcst/Report/soil/symposium/pdf/0700.pdf>>.
- Gazey C. & Davies S. (2009) Soil acidity: A guide for WA farmers and consultants. Department of Agriculture and Food, Western Australia, Perth.
- Gazey C., Fisher J. & Andrew J. (2012) Taking soil acidity seriously – results of grower surveys. Department of Agriculture and Food, Western Australia, Perth,
<http://biosecurity.wa.gov.au/objtwr/imported_assets/content/fcp/cu2012_final_2012_crop_updates_proceedings.pdf#page=132> [Accessed July 18, 2012].
- Gerik T. & Freebairn D. (2004) Management of extensive farming systems for drought-prone environments in North America and Australia. In: *New Directions for a Diverse Planet*. Proceedings of the 4th International Crop Science Congress. Published on CDROM, Published on CDROM. <http://cropsscience.org.au/icsc2004/pdf/2084_gerik.pdf>.
- Gillespie R., Dumsday R. & Bennett J. (2008) Estimating the value of environmental services provided by Australian farmers. Australian Farm Institute, Surrey Hills, Australia.
- Glover F., Whitworth K. L., Kappen P., Baldwin D. S., Rees G. N., Webb J. A. & Silvester E. (2011) Acidification and buffering mechanisms in acid sulfate soil wetlands of the Murray-Darling Basin, Australia. *Environmental Science & Technology* **45**, 2591-7.
- Goldspink B. H. & Howes K. M. (2001) *Fertilisers for Wines Grapes*. Department of Agriculture Western Australia, Perth.
- Gordon L. J., Finlayson C. M. & Falkenmark M. (2010) Managing water in agriculture for food production and other ecosystem services. *Agricultural Water Management* **97**, 512-9.
- Gourley C. J. P., Powell J. M., Dougherty W. J. & Weaver D. M. (2007) Nutrient budgeting as an approach to improving nutrient management on Australian dairy farms. *Australian Journal of Experimental Agriculture* **47**, 1064-74.
- Grealish G., Clifford D., Wilson P. & Ringrose-Voase A. (2011) National Soil Condition Monitoring for soil pH and soil carbon. Objectives, Design, Protocols, Governance and Reporting. CSIRO Land and Water Science Report for Caring for our Country. CSIRO, Canberra.
- Greene R. S. B., Kinnell P. I. A. & Wood J. T. (1994) Role of plant cover and stock trampling on runoff and soil-erosion from semi-arid wooded rangelands. *Australian Journal of Soil Research* **32**, 953.
- Greenway M. (2005) The role of constructed wetlands in secondary effluent treatment and water reuse in subtropical and arid Australia. *Ecological Engineering* **25**, 501–9.
- Gretton P. & Salma U. (1996) Land degradation and the Australian agricultural industry. Productivity Commission, Canberra.
- Griffiths B. S. & Philippot L. (2012) Insights into the resistance and resilience of the soil microbial community. *FEMS microbiology reviews* **In press**.
- Gunderson L. & Holling C. S. (2002) *Panarchy: Understanding Transformations in Human and Natural Systems*. Island Press, Washington, D.C. USA.
- Guo L. B. & Gifford R. M. (2002) Soil carbon stocks and land use change: a meta analysis. *Global Change Biology* **8**, 345-60.
- Haines-Young R. H. & Potschin M. B. (2009) Methodologies for defining and assessing ecosystem services. JNCC, Project Code C08-0170-0062. The University of Nottingham, Centre for Environmental Management, Nottingham, UK.
- Hairsine P. B., Barson M., Randall L. & Wilkinson S. N. (2008) Identification of areas in Australia where soil loss from hillslope erosion could be reduced. CSIRO, Canberra. <<http://nrmonline.nrm.gov.au/catalog/mql:2238>>.
- Hajkowicz S. & Young M. (2002) Value of returns to land and water and costs of degradation. A consultancy report to the

- National Land and Water Resources Audit. CSIRO, Canberra.
- HAL. (2003) A survey approach to investigate the soil factors associated with the productivity and sustainability of vegetable production in Australia (abstract of Project VG99057). Horticulture Australia Limited.
<http://www.horticulture.com.au/reports/search_final_reports_result.asp?src=&orgid=0&projid=375&strSearch=&strProjectNo=VG99057&strIndustry=0-All&strSortby=date&strDisplay=title&pageno=1>.
- HAL. (2008) Horticulture For Tomorrow - Horticulture's Environmental Contribution. Horticulture Australia Ltd,
<http://www.horticulturefortomorrow.com.au/about_horticulture/environmental_contribution.asp> [Accessed 12 July, 2012].
- HAL. (2010) Climate Change: Managing Variability and Carbon. Horticulture Australia Ltd.
<<http://ausveg.businesscatalyst.com/rnd/fact%20sheets/Climate%20Change%20and%20Managing%20Carbon.pdf>>.
- HAL. (2011) Benchmarking Soil Health for Improved Crop Health, Quality and Yields in the Temperate Australian Vegetable Industries. Horticulture Australia Limited.
- HAL. (2012a) Developing sustainable vegetable production systems. Horticulture Australia Ltd.
- HAL. (2012b) Quantifying carbon footprints in the Bundaberg region - a vehicle towards horticultural sustainability (abstract of Project HG09032). Horticulture Australia Limited.,
<http://www.horticulture.com.au/reports/search_final_reports_result.asp?src=&orgid=0&projid=4649&strSearch=&strProjectNo=HG09032&strIndustry=0-All&strSortby=date&strDisplay=title&pageno=1> [Accessed July 15, 2012].
- Hall R. L. & Calder I. R. (1993) Drop size modification by forest canopies: measurements using a disdrometer. *Journal of Geophysical Research* **98**, 18465–70.
- Hamblin A. (1996) Agronomy and the environment. In: *Agronomy - Science with its sleeves rolled up*. Proceedings of the 8th Australian Agronomy Conference, The University of Southern Queensland, Toowoomba, Queensland (eds D. L. Michalk and J. E. Pratley) p. [on line]. The Regional Institute Ltd.
<<http://www.regional.org.au/au/asa/1996/invited/033Hamblin.htm>>.
- Hamblin A., Tennant D. & Perry M. W. (1987) Management of soil water for wheat production in Western Australia. *Soil Use and Management* **3**, 63-9.
- Hamblin A. P. (1984) The effect of tillage on soil surface properties and the water balance of a xeralfic alfisol. *Soil & Tillage Research* **4**, 543-59.
- Hamblin A. P., Tennant D. & Cochrane H. (1982) Tillage and the growth of a wheat crop in a loamy sand [Western Australia]. *Australian Journal of Agricultural Research* **33**, 887-97.
- Hamilton G., Bakker D., Houlbrooke D. & Spann C. (2005) A Manual for Raised Bed Farming in Western Australia. The Department of Agriculture, Western Australia, Perth.
<http://www.agric.wa.gov.au/objtwr/imported_assets/content/lwe/land/cult/raisedbeds.pdf>.
- Haygarth P. & Ritz K. (2009) The future of soils and land use in the UK: Soil systems for the provision of land-based ecosystem services. *Land Use Policy* **26S**, S187-S97.
- Heckrath G., Brookes P. C., Poulton P. R. & Goulding K. W. T. (1995) Phosphorus leaching from soils containing different phosphorus concentrations in the Broadbalk Experiment. *Journal of environmental quality* **24**, 904–10.
- Helyar K., Cregan P. & Godyn D. (1990) Soil acidity in New-South-Wales - Current pH values and estimates of acidification rates. *Soil Research* **28**, 523-37.
- Helyar K. R. (1991) The management of acid soils. In: *Plant-Soil Interactions at Low pH* (eds R. J. Wright, V. C. Baligar and R. P. Murrman). Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Helyar K. R., Agriculture N., Institute A. R., Research W. & Council D. (1992) Predicting Lime Response and Managing Acid Soil Infertility: Final Report for the Wool Research and Development Council. N.S.W. Agriculture and Agricultural Research Institute, Sydney. <<http://books.google.com.au/books?id=GkSttgAACAAJ>>.
- Herbert A. (2009) Opportunity Costs of Land Degradation Hazards in The South-West Agriculture Region. Resource Management Technical Report 349. Department of Agriculture and Food, Western Australia, Perth.
- Heywood J. (2004) Improving riparian condition and controlling erosion. Queensland Murray Darling Committee, Toowoomba, Queensland. <<http://nrmonline.nrm.gov.au/catalog/mql:1300>>.
- Hillel D. (1980) *Introduction to Soil Physics*. Academic Press, San Diego.

- Holling C. S. (1996) Engineering resilience versus ecological resilience. In: *Engineering within Ecological Constraints* (ed P. C. Schulze) pp. 31-44. National Academy Press, Washington, D.C.
- Holmes K., Bellamy P. & Dixon J. (2011) Designing a soil pH monitoring network for the Western Australian wheatbelt. Department of Agriculture and Food WA, Perth.
- Hooper S. & Levantis C. (2011) Physical and financial performance benchmarks for grain producing farms , New South Wales – Victoria slopes agroecological zone. ABARES, Canberra.
- Hunt L. P., Petty S., Cowley R., Fisher A., Ash A. J. & MacDonald N. (2007) Factors affecting the management of cattle grazing distribution in northern Australia: preliminary observations on the effect of paddock size and water points. *The Rangeland Journal* **29**, 169.
- James C. D., Landsberg J. & Morton S. R. (1999) Provision of watering points in the Australian arid zone: a review of effects on biota. *Journal of arid environments* **41**, 87-121.
- Jansen A. & Robertson A. I. (2001) Relationship between livestock management and the ecological condition of riparian habitats along an Australian floodplain river. *Journal of Applied Ecology* **38**, 63-75.
- Jenkins A. & Alt S. (2007) Soil erosion solutions. Helping North Coast landholders reduce soil erosion. Soil erosion projects completed in the Northern Rivers, NSW 2005/06. NSW Government, Sydney.
<<http://www.dpi.nsw.gov.au/agriculture/resources/soils/erosion/soil-erosion-solutions>>.
- Jenkins A. & Alt S. (2009) Soil Erosion Solutions. Helping North Coast landholders reduce soil erosion. ~~Soil erosion~~ Soil erosion projects completed in the Northern Rivers, NSW 2007/2008. NSW Government, Sydney.
<<http://www.dpi.nsw.gov.au/agriculture/resources/soils/erosion/info-sheet>>.
- Jiang L. & Patel S. N. (2008) Community assembly in the presence of disturbance: a microcosm experiment. *Ecology* **89**, 1931-40.
- Johnston R. J. & Russell M. (2011) An operational structure for clarity in ecosystem service values. *Ecological Economics* **70**, 2243-9.
- Jones D. L., Rousk J., Edwards-Jones G., DeLuca T. H. & Murphy D. V. (2012) Biochar-mediated changes in soil quality and plant growth in a three year field trial. *Soil Biology and Biochemistry* **45**, 113-24.
- Karanja F., Reid N., Cacho O. & Kumar L. (2007) Evaluating the impact of integrated catchment management interventions on provision of ecosystem services using GIS. In: 25 years Landscape Ecology: Scientific Principles in Practice. Proceedings of the 7th IALE World Congress, 8-12 July 2007, Wageningen, The Netherlands. (eds R. G. H. Bunce, R. H. G. Jongman, L. Hojas and S. Weel). International Association of Landscape Ecology (IALE). <http://une-au.academia.edu/Karanja/Papers/246151/Evaluating_the_impact_of_integrated_catchment_management_interventions_on_provision_of_ecosystem_services_using_GIS>.
- Kemp A. & Connell P. (2001) Impact of land degradation on Australian Agriculture: A land values approach. ABARE report to National Land and Water Resources Audit, Canberra.
- Kemp D. R. & Dowling P. M. (2000) Towards sustainable temperate perennial pastures. *Australian Journal of Experimental Agriculture* **40**, 125-32.
- Kibblewhite M., Ritz K. & Swift M. (2008) Soil health in agricultural Systems. *Philosophical Transactions of the Royal Society of London B* **362**, 685-701.
- Kimetu J. M. & Lehmann J. (2010) Stability and stabilisation of biochar and green manure in soil with different organic carbon contents. *Soil Research* **48**, 577-85.
- Kingwell R. S., Pannell D. J. & Robinson S. D. (1993) Tactical responses to seasonal conditions in whole-farm planning in Western Australia. *Agricultural economics* **8**, 211-26.
- Kirkegaard J., Kirkby C. & Gupta V. (2007) Management Practices for Building Soil Carbon. In: The Healthy Soils Symposium: Can Australian Soils Sustain our Agricultural Systems? pp. 74-8. Land & Water Australia, Canberra.
<<http://lwa.gov.au/files/products/healthy-soils-sustainable-farms/pn30193/pn30193.pdf>>.
- Kroeger T. & Casey F. (2007) An assessment of market-based approaches to providing ecosystem services on agricultural lands. *Ecological Economics* **64**, 321–32.
- Kuske C. R., Yeager C. M., Johnson S., Ticknor L. O. & Belnap J. (2012) Response and resilience of soil biocrust bacterial communities to chronic physical disturbance in arid shrublands. *The ISME journal* **6**, 886-97.

- Lal R. (1997) Degradation and resilience of soils. *Philosophical Transactions of the Royal Society of London B* **352**, 997-1010.
- Lal R. (1999) Soil management and restoration for C sequestration to mitigate the accelerated greenhouse effect. *Progress in Environmental Science* **1**, 307-26.
- Lal R. (2001) Soil degradation by erosion. *Land Degradation & Development* **12**, 519-39.
- Lal R. (2004) Soil carbon sequestration to mitigate climate change. *Geoderma* **123**, 1-22.
- Lal R. (2010) Enhancing eco-efficiency in Agro-ecosystems through soil carbon sequestration. *Crop Science* **50S**, S120-S31.
- Land & Water Australia. (2005) ECOGRAZE — Sustainable Grazing Management for Northern Australia. Land & Water Australia, Canberra. <<http://lwa.gov.au/files/products/land-and-water-australia-corporate/ew071245/ew071245-cs-10.pdf>>.
- Landsberg J., James C. D., Maconochie J., Nicholls A. O., Stol J. & Tynan R. (2002) Scale-related effects of grazing on native plant communities in an arid rangeland region of South Australia. *Journal of Applied Ecology* **39**, 427-44.
- Lang D. & McDonald W. (2005)  Maintaining ground cover to reduce erosion and sustain production. NSW Department of Primary Industries, Gunnedah, NSW. <http://www.dpi.nsw.gov.au/__data/assets/pdf_file/0018/162306/ground-cover-for-pastures.pdf>.
- Lavelle P., Decaëns T., Aubert M., Barot S., Blouin M., Bureau F., Margerie P., Mora P. & Rossi J. P. (2006) Soil invertebrates and ecosystem services. *European Journal of Soil Biology* **42S**, S3-S15.
- Leys J., Butler H., Yang X. & Heidenreich S. (2010)  CEMSYS modelled wind erosion. Department of Environment, Climate Change and Water NSW on behalf of the Department of Environment, Water, Heritage and the Arts, Sydney. <<http://nrmonline.nrm.gov.au/catalog/mql:2239>>.
- Leys J., Heidenreich S. K., Strong C. L., McTainsh G. H. & Quigley S. (2011) PM10 concentrations and mass transport during “Red Dawn” – Sydney 23 September 2009. *Aeolian Research* **3**, 327-42.
- Leys J., Smith J., Macrae C., Rickards J., Yang X., Hairsine P., Dixon J. & McTainsh G. (2008) Improving the capacity to monitor wind and water erosion: A Review. Australian Government Department of Agriculture, Fisheries and Forestry, Canberra.
- Leys J., Smith J., MacRae C., Rickards J., Yang X., Randall L., Hairsine P., Dixon J. & McTainsh G. (2009) Improving The Capacity To Monitor Wind And Water Erosion: A Review. Department of Agriculture, Fisheries and Forestry, Canberra. <<http://nrmonline.nrm.gov.au/catalog/mql:2243>>
- Leys J. F. (1992) Cover levels to control soil and nutrient loss from wind erosion on sand plain country in central NSW. In: Australian Rangelands in a Changing Environment, 7th Biennial Conference pp. 84–91. Australian Rangeland Society.
- Leys J. F. (1998) Wind Erosion Processes and Sediments in Southeastern Australia. PhD thesis, Griffith University, Brisbane.
- Li G. D., Helyar K. R., Conyers M. K., Cullis B. R., Cregan P. D., Fisher R. P., Castleman L. J. C., Poile G. J., Evans C. M. & Braysher B. (2001) Crop responses to lime in long-term pasture-crop rotations in a high rainfall area in south-eastern Australia. *Australian Journal of Agricultural Research* **52**, 329-41.
- Li G. D., Singh R. P., Brennan J. P. & Helyar K. R. (2010) A financial analysis of lime application in a long-term agronomic experiment on the south-western slopes of New South Wales. *Crop and Pasture Science* **61**, 12-23.
- Li Y. X., Tullberg J. N. & Freebairn D. M. (2007) Wheel traffic and tillage effects on runoff and crop yield. *Soil & Tillage Research* **97**, 282-92.
- Lilley J. M. & Moore A. D. (2009) Trade-offs between productivity and ground cover in mixed farming systems in the Murrumbidgee catchment of New South Wales. *Animal Production Science* **49**, 837.
- Littleboy M., Silburn D. M., Freebairn D. M., Woodruff D. R., Hammer G. L. & Leslie J. K. (1992) Impact of soil erosion on production in cropping systems. I. Development and validation of a simulation model. *Australian Journal of Soil Research* **30**, 757-74.
- Llewellyn R. S. & D'Emden F. H. (2009) Understanding the drivers of no-till adoption in Australian agriculture. In: Landscapes Transformed: The Quiet Triumph of Conservation Tillage and Direct Seeding pp. 70-9. University of Saskatchewan Saskatoon, Saskatchewan. <<http://www.kis.usask.ca/CTConference.html>>.
- Llewellyn R. S. & D'Emden F. H. (2010) How much is enough: the steep final steps to extensive no-tillage cropping in Australia.

- In: Food Security from Sustainable Agriculture. Proceedings of 15th Agronomy Conference 2010 (eds H. Dove and R. A. Culvenor). The Regional Institute Ltd. <http://www.regional.org.au/au/asa/2010/crop-production/soil-water/7180_llewellynrs.htm>.
- Llewellyn R. S., D'Emden F. H. & Kuehne G. (2012) Extensive use of no-tillage in grain growing regions of Australia. *Field Crops Research* **132**, 204-12.
- Loch R. (2010) Optimising Turf Use to Minimise Soil Erosion on Construction Sites. Horticulture Australia Ltd, Sydney. <http://www.horticulture.com.au/reports/search_final_reports_result.asp?src=&orgid=0&projid=3702&strSearch=&strProjectNo=TU08033&strIndustry=0-All&strSortby=date&strDisplay=titledesc&pageno=1>.
- Loch R. J. (2000) Effects of vegetation cover on runoff and erosion under simulated rain and overland flow on a rehabilitated site on the Meandu Mine, Tarong, Queensland. *Australian Journal of Soil Research* **38**, 299-312.
- Loughran R. J., Elliott G. L., McFarlane D. J. & Campbell B. L. (2004) A survey of soil erosion in Australia using Caesium-137. *Australian Geographical Studies* **42**, 221-33.
- Ludwig J. A. & Tongway D. J. (1995) Desertification in Australia: An eye to grass roots and landscapes. *Environmental monitoring and Assessment* **37**, 231-7.
- Ludwig J. A., Wilcox B. P., Breshears D. D., Tongway D. J. & Imeson A. C. (2005) Vegetation patches and runoff-erosion as interacting ecohydrological processes in semiarid landscapes. *Ecology* **86**, 288-97.
- MA. (2005) *Millennium Ecosystem Assessment. Ecosystems and Human Well-Being: Current State and Trends*. Island Press, Washington DC USA.
- MA. (2005) *Ecosystems and Human Well-Being: Synthesis*. Island Press, Washington DC, USA.
- MacDonald D. H., Bark R., Garrick D., Banerjee O., Connor J. & Morrison M. (2011) Multiple benefits through the life cycle of the basin plan. In: *Basin Futures: Water Reform in the Murray-Darling Basin* (eds D. Connell and R. Q. Grafton) pp. 263-75. ANU E-Press, Canberra.
- MacEwan R., Gardner W., Ellington A., Hopkins D. & Bakker A. (1992) Tile and mole drainage for control of waterlogging in duplex soils of south-eastern Australia. *Australian Journal of Experimental Agriculture* **32**, 865-78.
- MacKenzie D. (2010) Soil Carbon Sequestration under Pasture in Australian Dairy Regions. Dairy Australia Review. Dairy Australia, Melbourne. <http://www.dairyingfortomorrow.com/uploads/documents/file/Reports/Final%20Soil%20Carbon%20Sequestration%20in%20Aust%20dairy%20regions%20report%20feb%2010.pdf>.
- Macleod N. & Mclvor J. (2004) Assessing financial and environmental impacts of management options: Managing for biodiversity in the rangelands. CSIRO, Canberra.
- Mahowald N. M., Kloster S., Engelstaedter S., Moore J. K., Mukhopadhyay S., McConnell J. R., Albani S., Doney S. C., Bhattacharya A., Curran M. A. J., Flanner M. G., Hoffman F. M., Lawrence D. M., Lindsay K., Mayewski P. A., Neff J., Rothenberg D., Thomas E., Thornton P. E. & Zender C. S. (2010) Observed 20th century desert dust variability: impact on climate and biogeochemistry. *Atmospheric Chemistry and Physics* **10**, 10875-93.
- Mallawaarachchi T. (1993) Economic costs of cropland erosion: An application of GIS-based resource accounting approach (as reported on Envalue database). In: 37th Annual Conference of the Australian Agricultural Economics Society, University of Sydney. <http://www.environment.nsw.gov.au/envalueapp/studydetail.asp?id_study=358>.
- Mallawaarachchi T., Young M., Walker P. & Smyth B. (1994) Economic assessment of land management options: A GIS-based natural resource accounting approach (as reported on Envalue database). In: *Procedures for Economic Assessment of Management Options for Dryland Salinity: LWRRDC Occasional Paper No. 06/94*. LWRRDC, Canberra.
- Mason W. K. & Kay G. (2000) Temperate pasture sustainability key program. *Australian Journal of Experimental Agriculture* **40**, 121-4.
- Maynard S., James D. & Davidson A. (2010) The Development of an Ecosystem Services Framework for South East Queensland. *Environmental Management* **45**, 881-95.
- McAlpine K. G. & Wotton D. M. (2009) Conservation and the delivery of ecosystem services. New Zealand Department of Conservation, Wellington, NZ. <<http://www.doc.govt.nz/upload/documents/science-and-technical/sfc295entire.pdf>>.
- McArthur W. M. (2004) *Reference Soils of South-Western Australia*. Australian Society of Soil Science, Perth

- McCarthy M. G., Dry P. R., Hayes P. R. & Davidson D. M. (1992) Soil management and frost control. In: *Viticulture. Volume 2. Practices*. (ed B. G. Coombe) pp. 149-77. Winetitles, Adelaide.
- McCosker T. (2000) Cell grazing - The first 10 years in Australia. *Tropical Grasslands* **34**, 207-18.
- McIvor J. G. & McIntyre S. (2002) Understanding grassy woodland ecosystems. In: *Managing and Conserving Grassy Woodlands* (eds S. McIntyre, J. G. McIvor and K. M. Heard) pp. 1-23. CSIRO Publishing, Camberwell, Victoria.
- McPhee J. E. (2009) Controlled Traffic Farming Systems for the Tasmanian vegetable industry (Reporting on HAL Project VG07058). Horticulture Australia Ltd, Sydney. <<http://ecite.utas.edu.au/56242>>.
- McTainsh G., Lynch A. & Burgess R. (1990) Wind erosion in eastern Australia. *Australian Journal of Soil Research* **28**, 323-39.
- McTainsh G. & Strong C. (2007) The role of aeolian dust in ecosystems. *Geomorphology* **89**, 39-54.
- McTainsh G., Strong C., Leys J., Baker M., Tews K. & Barton K. (2009) Wind erosion histories, model input data and community DustWatch. Atmospheric Environment Research Centre, Griffith University, Brisbane.
- McTainsh G. H., Leys J. F., O'Loingsigh T. & Strong C. L. (2011) Wind erosion and land management in Australia during 1940-1949 and 2000-2009. Report prepared for the Australian Government Department of Sustainability, Environment, Water, Population and Communities on behalf of the State of the Environment 2011 Committee. DSEWPaC, Canberra.
- Mele P. M. & Carter M. R. (1993) Effect of climatic factors on the use of microbial biomass as an indicator of changes in soil organic matter. In: *Soil Organic Matter Dynamics and Sustainability of Tropical Agriculture* (eds K. Mulongoy and R. Merckx) pp. 57-63. John Wiley & Sons, Chichester, UK.
- Mele P. M. & Crowley D. E. (2008) Application of self-organizing maps for assessing soil biological quality. *Agriculture, Ecosystems & Environment* **126**, 139-52.
- Michalk D. L., Dowling P. M., Kemp D. R., King W. M., Packer I. J., Holst P. J., Jones R. A. E., Priest S. M., Millar G. D., Brisbane S. & Stanley D. F. (2003) Sustainable grazing systems for the Central Tablelands, New South Wales. *Australian Journal of Experimental Agriculture* **43**, 861-74.
- Millar G. D. & Badgery W. B. (2009) Pasture cropping: a new approach to integrate crop and livestock farming systems. *Animal Production Science* **49**, 777-87.
- Moore G., Findlater P. & Carter D. (2001) 7.1 Wind erosion. In: *Soilguide: A Handbook for Understanding and Managing Agricultural Soils. Bulletin 4343* (ed G. Moore) pp. 209-22. Agriculture Western Australia, Perth, WA.
- Moran C. J. (1998) Land degradation processes and water quality effects: decline in soil structure. In: *Farming Action Catchment Reaction: The Effect of Dryland Farming on the Natural Environment* (eds J. Williams, R. A. Hook and H. L. Gascoigne) pp. 141-57. CSIRO Publishing, Collingwood, Victoria, Australia.
- Morin J. & Van Winkel J. (1996) The effect of raindrop impact and sheet erosion on infiltration rate and crust formation. *Soil Science Society of America Journal* **60**, 1223-7.
- Murphy D. V., Cookson W. R., Braimbridge M., Marschner P., Jones D. L., Stockdale E. A. & Abbott L. K. (2011) Relationships between soil organic matter and the soil microbial biomass (size, functional diversity, and community structure) in crop and pasture systems in a semi-arid environment. *Soil Research* **49**, 582-94.
- Murray R. S. (2007) A Review of Dr Bruce Cockroft's Work for Australian Irrigated Horticulture Milestone Report 2. National Program for Sustainable Agriculture, Canberra. <<http://npsi.gov.au/files/products/national-program-sustainable-irrigation/pn21945/pn21945.pdf>>.
- Nahlik A. M., Kentula M. E. & Fennessy M. S. (2012) Where is the consensus? A proposed foundation for moving ecosystem service concepts into practice. *Ecological Economics* **77**, 27-35.
- Namoi Catchment Management Authority. (2010) Namoi Catchment Action Plan 2010-2020. Draft for Public Comment July 2010. Namoi Catchment Management Authority, Tamworth, Australia. <http://www.namoi.cma.nsw.gov.au/2010_cma_cap_web_2.pdf>.
- Nash D., Clemow L., Hannah M., Barlow K. & Gangaiya P. (2005) Modelling phosphorus exports from rain-fed and irrigated pastures in southern Australia. *Australian Journal of Soil Research* **43**, 745.
- Nash D. & Murdoch C. (1997) Phosphorus in runoff from a fertile dairy pasture. *Australian Journal of Soil Research* **35**, 419-29.
- Neil D. T. & Fogarty P. (1991) Land use and sediment yield on the Southern Tablelands of New South Wales. *Australian*

- Journal of Soil and Water Conservation* **4**, 33-9.
- Neil D. T. & Galloway R. W. (1989) Estimation of sediment yields from catchments farm dam. *Australian Journal of Soil and Water Conservation* **2**, 46-51.
- Nelson D. R. & Mele P. M. (2006) The impact of crop residue amendments and lime on microbial community structure and nitrogen-fixing bacteria in the wheat rhizosphere. *Australian Journal of Soil Research* **44**, 319-29.
- NLWRA. (2001) Australian Agricultural Assessment 2001. Australian Government, Canberra, <http://www.anra.gov.au/topics/agriculture/pubs/national/agriculture_contents.html> [Accessed 18 May, 2012].
- NLWRA. (2007) Indicator Protocol: Soil Acidification, Canberra. <<http://lwa.gov.au/files/products/national-land-and-water-resources-audit/pn21220/pn21220.pdf>>.
- NLWRA. (2008) Signposts for Australian agriculture: The Australian horticulture industry. NLWRA, Canberra. <<http://lwa.gov.au/files/products/national-land-and-water-resources-audit/pn21405/pn21405.pdf>>.
- O'Reagain P., Bushell J. & Holmes B. (2011) Managing for rainfall variability: long-term profitability of different grazing strategies in a northern Australian tropical savanna. *Animal Production Science* **51**, 210.
- OECD/FAO. (2011) OECD-FAO Agricultural Outlook 2011-2020. OECD Publishing and FAO, Paris.
- Palm C., Sanchez P., Ahamed S. & Awiti A. (2007) Soils: A contemporary perspective. *Annual Review of Environmental Resources* **32**, 99-129.
- Pannell D. & Vanclay F. (2011) *Changing Land Management: Adoption of New Practices by Rural Landholders*. CSIRO Publishing, Collingwood.
- Parton W. J., Schimel D. S., Cole C. V. & Ojima D. S. (1987) Analysis of factors controlling organic matter in Great Plains Grasslands. *Soil Science Society of America Journal* **51**, 1173-79.
- Passioura J., Kirkby C., Baldock J., Kirkegaard J. & Peoples M. (2008) The hidden costs of carbon sequestration. *GroundCover* **76**, [on line]
<http://www.grdc.com.au/director/events/groundcover?item_id=409DFCDF9E9F1ECE7C574ACD7A028479&article_id=B6DA574BC1352B85C7E842C6C22237D8>.
- Pattison T., Moody P. & Bagshaw J. (2010) Soil Health for Vegetable Production in Australia. Queensland Government, Brisbane. <http://www.daff.qld.gov.au/26_17025.htm>.
- Paustian K., Collins H. P. & Paul E. A. (1997) Management controls on Soil Carbon. In: *Soil Organic Matter in Temperate Agroecosystems* (eds E. A. Paul, K. A. Paustian, E. T. Elliot and C. V. Cole) pp. 15-49. CRC Press, Boca Raton.
- Phillips C., Ekanayake J., Marden M. & Watson A. (2000) Stabilising parameters of vegetation: a critical look down-under. In: *Landscape 2000, Proceedings of International Landscape Conference*.
- Pimentel D., Harvey C., Resosudarmo P. & Sinclair K. (1995) Environmental and economic costs of soil erosion and conservation benefits. *Science* **267**, 1117-23.
- Pimentel D. & Kounang N. (1998) Ecology of Soil Erosion in Ecosystems. *Ecosystems* **1**, 416-26.
- Post W. M., Emanuel W. R., Zinke P. J. & Stangenberger A. G. (1982) Soil carbon pools and world life zones. *Nature* **298**, 156-9.
- Potts D. L., Huxman T. E., Enquist B. J., Weltzin J. F. & Williams D. G. (2006) Resilience and resistance of ecosystem functional response to a precipitation pulse in a semi-arid grassland. *Journal of Ecology* **94**, 23-30.
- Pretty J. (2008) Agricultural sustainability: concepts, principles and evidence. *Philosophical transactions of the Royal Society of London B* **363**, 447-65.
- Radford B. J., Thompson J. P. & Thomas G. A. (1993) Tillage and stubble management. In: *Cropping Strategies for the Next Decade – Queensland Crop Production Conference Proceedings* (eds W. M. Strong and B. J. Radford) pp. 149-62. Queensland Department of Primary Industries.
- Raupach M., McTainsh G. & Leys J. (1994) Estimates of dust mass in recent major Australian dust storms. *Australian Journal of Soil and Water Conservation* **7**, 20-4.
- Raya A. M., Zuazo V. H. D. & Martinez J. R. F. (2006) Soil erosion and runoff response to plant-cover strips on semiarid slopes (SE Spain). *Land Degradation & Development* **17**, 1-11.

- Reid G. (2002) Soil and Nutrient Loss in Macadamia Lands. Horticulture Australia Ltd, Sydney.
- Ridley A. M. (1995) A study of the factors governing the rate of soil acidification under subterranean clover based annual and perennial pastures. PhD Thesis, University of Melbourne Melbourne.
- Ridley A. M. & Coventry D. R. (1995) Soil Acidification Under Pastures of North-Eastern Victoria Five Years After Liming. In: *Plant-soil interactions at low pH: principles and management. Proceedings of the 3rd International Symposium of Plant-Soil Interactions at Low pH. Brisbane, Qld, 12-16 September 1993* (eds R. A. Date, N. J. Grundon, G. E. Rayment and M. E. Probert) pp. 473-7. Kluwer Academic, The Netherlands.
- Ridley A. M. & Windsor S. M. (1992) Persistence and tolerance to soil acidity of phalaris and cocksfoot in north-eastern Victoria. *Australian Journal of Experimental Agriculture* **32**, 1069-75.
- Ringold P. L., Boyd J., Landers D. & Weber M. (2009) Report from the Workshop on Indicators of Final Ecosystem Services for Streams. U.S. Environmental Protection Agency, Corvallis, OR, USA.
- Ringold P. L., Nahlik A. M., Boyd J. W. & Bernard D. (2011) Report from the Workshop on Indicators of Final Ecosystem Goods and Services for Wetlands and Estuaries. US Environmental Protection Agency.
- Robertson G. A. (1987) Soil Management for Sustainable Agriculture. Government of Western Australia, Perth.
- Robinson D. A., Hockley N., Dominati E., Lebron I., Scow K. M., Reynolds B., Emmett B. A., Keith A. M., de Jonge L. W., Schjonning P., Moldrup P., Jones S. B. & Tuller M. (2012) Natural Capital, Ecosystem Services, and Soil Change: Why Soil Science Must Embrace an Ecosystems Approach. *Vadose Zone Journal* **11**, vzj2011.0051.
- Roget D. K. (1995) Decline of root rot (*Rhizoctonia solani* AG-8) in wheat in a tillage and rotation experiment at Avon South Australia. *Australian Journal of Experimental Agriculture* **35**, 1009-13.
- Rolfe J., McCosker J. & Windle J. (2008) Identifying the incentives that graziers in central-western Queensland need to manage for conservation outcomes. *Rangeland Journal* **30**, 297 – 307.
- Rosewell C. J. (1993) Soilloss. A Program to Assist in the Selection of Management Practices to Reduce Erosion. Department of Conservation and Land Management, Sydney.
- Rotstayn L. D., Collier M. A., Mitchell R. M., Qin Y., Campbell S. K. & Dravitzki S. M. (2011) Simulated enhancement of ENSO-related rainfall variability due to Australian dust. *Atmospheric Chemistry and Physics* **11**, 6575-92
- Sanderman J., Baldock J., Hawke B., Macdonald L., Massis-Puccini A. & Szarvas S. (2011) National Soil Carbon Research Programme: Field and Laboratory Protocols. CSIRO publishing, Adelaide.
- Sanderman J., Farquharson R. & Baldock J. (2010) Soil Carbon Sequestration potential: A review for Australian agriculture. Department of Climate Change and Energy Efficiency, Adelaide.
- Sandhu H. S., Wratten S. D., Cullen R. & Case B. (2008) The future of farming: The value of ecosystem services in conventional and organic arable land. An experimental approach. *Ecological Economics* **64**, 835-48.
- Sangha K. K., Midmore D. J., Rolfe J. & Jalota R. K. (2005) Tradeoffs between pasture production and plant diversity and soil health attributes of pasture systems of central Queensland, Australia. *Agriculture, Ecosystems and Environment* **111**, 93-103.
- Sanjari G., Ghadiri H., Ciesiolka C. A. A. & Yu B. (2008) Comparing the effects of continuous and time-controlled grazing systems on soil characteristics in Southeast Queensland. *Soil Research* **46**, 348-58.
- Sanjari G., Yu B., Ghadiri H., Ciesiolka C. A. A. & Rose C. W. (2009) Effects of time-controlled grazing on runoff and sediment loss. *Soil Research* **47**, 796-808.
- Scanlan J., Pressland A. & Myles D. (1996) Run-Off and Soil Movement on Mid-Slopes in North-East Queensland [Australia] Grazed Woodlands. *The Rangeland Journal* **18**, 33-46.
- Schmidt C. P. & Belford R. K. (1993) A comparison of different tillage-seeding systems: the interaction of tillage and time of sowing on sandplain soils in Western Australia [wheat]. *Australian Journal of Experimental Agriculture* **33**, 895-900.
- Schmidt C. P., Belford R. K. & Tennant D. (1994) Effect of different direct drilling and conventional sowing techniques on soil strength, root growth and grain yield of wheat on sandplain soils in Western Australia. *Australian Journal of Agricultural Research* **45**, 547-64.
- Schmidt M. W. I., Torn M. S., Abiven S., Dittmar T., Guggenberger G., Janssens I. A., Kleber M., Kogel-Knabner I., Lehmann J., Manning D. A. C., Nannipieri P., Rasse D. P., Weiner S. & Trumbore S. E. (2011) Persistence of soil organic

- matter as an ecosystem property. *Nature* **478**, 49-56.
- Schumann B. (1999) The Causes of Soil Acidity. NSW Agriculture, Sydney, <http://www.dpi.nsw.gov.au/__data/assets/pdf_file/0009/167175/acidity-causes.pdf> [Accessed 12 July, 2012].
- Scott B. J., Eberbach P. L., Evans J. & Wade L. J. (2010) Stubble Retention in Cropping Systems in Southern Australia: Benefits and Challenges. New South Wales through Department of Industry and Investment, Wagga Wagga.
- Scott B. J., Ridley A. M. & Conyers M. K. (2000) Management of soil acidity in long-term pastures of south-eastern Australia: a review. *Australian Journal of Experimental Agriculture* **40**, 1173-98.
- Scott J. F. & Farquharson R. J. (2004) An assessment of the economic impacts of NSW agriculture research and extension: conservation farming and reduced tillage in Northern NSW. NSW Department of Primary Industries, Sydney. <http://www.dpi.nsw.gov.au/__data/assets/pdf_file/0009/146655/err-19-An-Assessment-of-the-Economic-Impacts-of-NSW-Agriculture-Research-and-Extension-Conservation-Farming-and-Reduced-Tillage-in-Northern-NSW.pdf>.
- Seybold C. A., Herrick J. E. & Brejda J. J. (1999) Soil resilience: A fundamental component of soil quality. *Soil Science* **164**, 224-34.
- Sherren K., Fischer J. & Fazey I. (2012) Managing the grazing landscape: Insights for agricultural adaptation from a mid-drought photo-elicitation study in the Australian sheep-wheat belt. *Agricultural Systems* **106**, 72-83.
- Silburn D. M., Carroll C., Ciesiolka C. A. A., deVoil R. C. & Burger P. (2011) Hillslope runoff and erosion on duplex soils in grazing lands in semi-arid central Queensland. I. Influences of cover, slope, and soil. *Soil Research* **49**, 105.
- Silburn D. M., Robinson J. B. & Freebairn D. M. (2007) Why restore marginal cropland to permanent pasture? Land resource and environmental issues. *Tropical Grasslands* **41**, 139-53.
- Silver W. L., Ostertag R. & Lugo A. E. (2000) The potential for carbon sequestration through reforestation of abandoned tropical agricultural and pasture lands. *Restoration Ecology* **8**, 394-407.
- Simpfendorfer S., Heenan D. P., Kirkegaard J. & Lindbeck K. D., and Murray, G M (2004) Impact of tillage on lupin growth and the incidence of pathogenic fungi in southern New South Wales. *Australian Journal of Experimental Agriculture* **44**, 53-6.
- Slattery W. J., Conyers M. K. & Aitkin R. L. (1989) Soil ph, aluminium, manganese and lime requirement. In: *Soil Analysis: An interpretation Manual* (eds K. I. Peverill, L. A. Sparrow and D. J. Reuter) pp. 103-28. Academic Press, Sydney.
- Slavich P. G. & Cox J. (2010) Improving soil and orchard floor management in macadamias. Horticulture Australia Ltd, Sydney.
- Smith J. & Leys J. (2009) Identification of areas within Australia for reducing soil loss by wind erosion. Bureau of Rural Sciences, Canberra. <<http://nrmonline.nrm.gov.au/catalog/mql:2246>>.
- Smith M. S. & Belvins R. L. (1987) Effect of conservation tillage on biological and chemical soil conditions: Regional and temporal variability. In: *Effects of Conservation Tillage on Groundwater Quality* (eds T. J. Logan, J. M. Davidson, J. L. Baker and M. K. Overcash) pp. 149-66. Lewis Publishers, Michigan USA.
- Smith P., Davies C. A., Ogle S., Zanchi G., Bellarby J., Bird N., Boddey R. M., McNamara N. P., Powlson D., Cowie A., Noordwijk M. v., Davis S. C., Richter D. D. B., Kryzanowski L., Wijk M. T. v., Stuart J., Kirton A., Eggar D., Newton-Cross G., Adhya T. K. & Braimoh A. K. (2012) Towards an integrated global framework to assess the impacts of land use and management change on soil carbon: current capability and future vision. *Global Change Biology* **18**, 2089-101.
- Soils Research Development and Extension Working Group. (2011) A stocktake of Australia's current investment in soils research, development and extension: A snapshot for 2010-11. Department of Agriculture, Fisheries and Forestry, Canberra.
- Southorn N. & Cattle S. (2004) The dynamics of soil quality in livestock grazing systems. In: SuperSoil 2004: 3rd Australian New Zealand Soils Conference. University of Sydney. <www.regional.org.au/au/asssi/>.
- Sparling G. P. (1997) Soil microbial biomass, activity and nutrient cycling as indicators of soil health. In: *Biological Indicators of Soil Health* (eds C. Pankhurst, B. M. Doube and V. V. S. R. Gupta) pp. 97-119. CAB International, New York.
- Sparrow A. D., Friedel M. H. & Tongway D. J. (2003) Degradation and recovery processes in arid grazing lands of central Australia Part 3: Implications at landscape scale. *Journal of arid environments* **55**, 349-60.
- Stafford Smith D. M., McKeon G. M., Watson I. W., Henry B. K., Stone G. S., Hall W. B. & Howden S. M. (2007) Learning from episodes of degradation and recovery in variable Australian rangelands. *Proceedings of the National Academy of*

- Sciences of the United States of America* **104**, 20690-5.
- Steffen W., Persson A., Deutsch L., Zalasiewicz J., Williams M., Richardson K., Crumley C., Crutzen P., Folke C., Gordon L., Molina M., Ramanathan V., Rockström J., Scheffer M., Schellnhuber H. J. & Svedin U. (2011) The anthropocene: from global change to planetary stewardship. *AMBIO: A Journal of the Human Environment* **40**, 739-61.
- Strahan R. & Hoffman A. (2009) Estimating the economic implications for broad acre cropping farms in the Fitzroy Basin catchments of adoption of best management practices. Queensland Primary Industries and Fisheries, Brisbane. <<https://www.grainsbmp.com.au/images/documents/Economic%20report%202009.pdf>>.
- Swinton S. M., Lupi F., Robertson G. P. & Hamilton S. K. (2007) Ecosystem services and agriculture: cultivating agricultural ecosystems for diverse benefits. *Ecological Economics* **64**, 245–52.
- Swinton S. M., Lupi F., Robertson G. P. & Landis D. A. (2006) Ecosystem services from agriculture: looking beyond the usual suspects. *American Journal of Agricultural Economics*, 1160–6.
- Sylvain Z. A. & Wall D. H. (2011) Linking soil biodiversity and vegetation: Implications for a changing planet. *American journal of botany* **98**, 517-27.
- Teague W. R., Dowhower S. L., Baker S. A., Haile N., DeLaune P. B. & Conover D. M. (2011) Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems & Environment* **141**, 310-22.
- TEEB. (2008) The Economics of Ecosystems and Biodiversity -An Interim Report. TEEB, Geneva.
- TEEB. (2009) TEEB – The Economics of Ecosystems and Biodiversity for National and International Policy Makers – Summary: Responding to the Value of Nature. TEEB, Geneva. <<http://www.teebweb.org/LinkClick.aspx?fileticket=dYhOxrQWffs%3D&tabid=1019&mid=1931>>.
- Thomas G. A., Titmarsh G. W., Freebairn D. M. & Radford B. J. (2007) No-tillage and conservation farming practices in grain growing areas of Queensland - A review of 40 years of development. *Australian Journal of Experimental Agriculture* **47**, 887-98.
- Tozer P. (2012) Economic Impact of Off Site Wind Erosion. University of New England, Armidale, NSW. <<http://ageconsearch.umn.edu/bitstream/124458/2/2012AC%20Tozer%20CP.pdf>>.
- Turbé A., Toni A. D., Benito P., Lavelle P., Lavelle P., Ruiz N., Putten W. H. V. d., Labouze E. & Mudgal S. (2010) Soil biodiversity: functions, threats and tools for policy makers. Bio Intelligence Service, IRD, and NIOO Report for European Commission (DG Environment). European Commission.
- Turner R. K. & Daily G. C. (2007) The ecosystem services framework and natural capital conservation. *Environmental and Resource Economics* **39**, 25-35.
- Ugalde D., Brungs A., Kaebernick M., McGregor A. & Slattery B. (2007) Implications of climate change for tillage practice in Australia. *Soil & Tillage Research* **97**, 318-30.
- UK Government. (2011) The Natural Choice: Securing the Value of Nature. UK Government, London.
- UK National Ecosystem Assessment. (2011) The UK National Ecosystem Assessment: Synthesis of the Key Findings. UNEP-WCMC, Cambridge, UK.
- UK National Ecosystem Assessment. (2011) The UK National Ecosystem Assessment Technical Report. UNEP-WCMC, Cambridge.
- van Buren M. & Price R. (2004) Breaking ground – Key findings from 10 years of the National Dryland Salinity Program. Land & Water Australia, Canberra.
- van de Koppel J., Rietkerk M. & Weissing F. J. (1997) Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends in Ecology & Evolution* **12**, 352-6.
- Waddell P. A., Thomas P. W. E. & Findlater P. A. (2012) A Report on the Gascoyne River Catchment Following the 2010/11 Flood Events. Department of Agriculture and Food, Western Australia, Perth.
- Walker B., Holling C. S., Carpenter S. R. & Kinzig A. (2004) Resilience, adaptability and transformability in socialecological systems. *Ecology and Society* **9**, [online] <http://www.ecologyandsociety.org/vol9/iss2/art5/>.
- Walker B. & Salt D. (2006) *Resilience Thinking: Sustaining Ecosystems and People in a Changing World*. Island Press, Washington DC, USA.

- Wall D. H. & Virginia R. A. (2000) The world beneath our feet: soil biodiversity and ecosystem functioning. In: *Nature and Human Society: The Quest for a Sustainable World* (ed P. H. Raven) pp. 225–41. National Academy Press, Washington DC.
- Wallace K. J. (2007) Classification of ecosystem services: Problems and solutions. *Biological Conservation* **139**, 235–46.
- Walpole S., Sinden J. & Yapp T. (1996) Land quality as an input to production: the case of land degradation and agricultural output. *Economic Analysis and Policy* **26**, 185 - 207.
- Wasson R. J., Olive L. J. & Rosewell C. J. (1996) Rates of erosion and sediment transport in Australia. In: *Erosion and Sediment Yield: Global and Regional Perspectives: Proceedings of an International Symposium Held at Exeter, UK* p. 139. IAHS. <<http://books.google.com/books?hl=en&lr=&id=bZ-ufVQV5yAC&oi=fnd&pg=PA139&dq=Rates+of+erosion+and+sediment+transport+in+Australia+R+J+WASSON&ots=u-NjMYzC5Q&sig=wiPFvo4roD19DSqo6d4yi4wfxYE>>.
- Watson P. (2006) Dairying for Tomorrow Survey of NRM Practices on Dairy Farms. Down to Earth Research, Victoria, Australia.
<<http://www.dairyingfortomorrow.com/uploads/documents/DfT%20survey%202006%20final%20version%2022.9.06.pdf>>.
- Weber J. L. (2007) Accounting for soil in the SEEA. European Environment Agency, Rome.
- Webb, M. J., Baldock, J. A., Unkovich, M., Gregory, L., Jacquier, D., Wilson, P., Grundy, M. (in preparation) Acidity and acidification of Australian soils. CSIRO Sustainable Agriculture Flagship report prepared for Caring for our Country.
- Wells T. & Chan K. (1996) Environmental impact of alternative horticultural production systems in the Hawkesbury-Nepean catchment. Horticulture Australia Ltd, Sydney.
<http://scholar.google.com/scholar?q=related:1OBhf0RFd7MJ:scholar.google.com/&hl=en&num=30&as_sdt=0,5>.
- White P. (1990) The influence of alternative tillage systems on the distribution of nutrients and organic-carbon in some common Western Australian wheatbelt soils. *Australian Journal of Soil Research* **28**, 95.
- White P. F. (1990) The influence of alternative tillage systems on the distribution of nutrients and organic carbon in some common Western Australian wheat belt soils. *Australian Journal of Agricultural Research* **28**, 95-116.
- White R. E. (2009) *Understanding Vineyard Soil*. Oxford University Press, New York.
- Whitten M. G., Wong M. T. F. & Rate A. W. (2000) Amelioration of subsurface acidity in the south-west of Western Australia: downward movement and mass balance of surface-incorporated lime after 2-15 years. *Soil Research* **38**, 711-28.
- Wightman B., Peries R., Bluett C. & Johnston T. (2005) Permanent raised bed cropping in southern Australia: Practical guidelines for implementation. In: *ACIAR Proceedings No. 121 – Evaluation and performance of permanent raised bed cropping systems in Asia, Australia and Mexico* (eds C. H. Roth, R. A. Fischer and C. A. Meisner) pp. 173-90. Australian Centre for International Agricultural Research, Canberra.
<[http://vro.dpi.vic.gov.au/dpi/vro/vrosite.nsf/pages/soil_mgmt_raised_beds_pdf/\\$FILE/Perm_raisedbed_cropping_southern_Australia.pdf](http://vro.dpi.vic.gov.au/dpi/vro/vrosite.nsf/pages/soil_mgmt_raised_beds_pdf/$FILE/Perm_raisedbed_cropping_southern_Australia.pdf)>.
- Williams P. & Young M. (1999) Costing dust: how much does wind erosion cost to the people of South Australia. CSIRO, Adelaide, SA.
- Wilson B., Chapman G. & Koen T. (2007) Report on Indicator Protocol Trial for Soil Carbon Monitoring, NSW: National Monitoring and Evaluation Framework. NLWRA, Canberra.
- Wilson P., Baldock J., Grundy M., Jacquier D., Griffin T., Moody P., Chapman G., Hall J., Maschmedt D., Crwaford D., Hill J. & Kidd D. (2009) Identification of land with a risk of acidification, Brisbane, Qld.
<<http://nrmonline.nrm.gov.au/catalog/mql:2560>>.
- Windle J. & Rolfe J. (2007) Developing a benefit transfer database for environmental values in Queensland. In: *51st Annual Australian Agricultural and Resource Economics Society Conference*, Queenstown, New Zealand pp. 13-6. AARES.
- Wylie P. (1997) Practical and economic considerations. In: *Sustainable Crop Production in the Sub-Tropics: An Australian Perspective* (eds A. L. Clarke and P. B. Wylie) pp. 329–38. Queensland Department of Primary Industries, Brisbane.
- Yapp T. P., Walpole S. C. & Sinden J. A. (1992) Soil conservation and land degradation - finding the balance [Australia]. *Search* **23**, 308-10.

- Yates C. J., Norton D. A. & Hobbs R. J. (2000) Grazing effects on plant cover, soil and microclimate in fragmented woodlands in south- western Australia: implications for restoration. *Austral Ecology* **25**, 36-47.
- Zhang B., Deng H., Wang H.-I., Yin R., Hallett P. D., Griffiths B. S. & Daniell T. J. (2010) Does microbial habitat or community structure drive the functional stability of microbes to stresses following re-vegetation of a severely degraded soil? *Soil Biology and Biochemistry* **42**, 850-9.
- Zhang W., Ricketts T. H., Kremen C., Carney K. & Swinton S. M. (2007) Ecosystem services and dis-services to agriculture. *Ecological Economics* **64**, 253-60.
- Zwieten L. V., Graham D., Cox J., Ayres M., Rust J., Vancov T., Morris S. & Walker B. (2001) Soil health in north coast agriculture: assessment and rehabilitation. In: *Soil Health: The Foundation of Sustainable Agriculture* pp. 88-95. NSW Department of Primary Industries. <<http://www.dpi.nsw.gov.au/agriculture/resources/soils/structure/workshop>>.